

include an attempt to obtain additional evidence or the holding of an informal conference with the claimant. Upon the request of the claimant, the reviewer will conduct a hearing under §3.103(c).

(d) The reviewer may grant a benefit sought in the claim notwithstanding §3.105(b), but, except as provided in paragraph (e) of this section, may not revise the decision in a manner that is less advantageous to the claimant than the decision under review. A review decision made under this section will include a summary of the evidence, a citation to pertinent laws, a discussion of how those laws affect the decision, and a summary of the reasons for the decision.

(e) Notwithstanding any other provisions of this section, the reviewer may reverse or revise (even if disadvantageous to the claimant) prior decisions of an agency of original jurisdiction (including the decision being reviewed or any prior decision that has become final due to failure to timely appeal) on the grounds of clear and unmistakable error (see §3.105(a)).

(f) Review under this section does not limit the appeal rights of a claimant. Unless a claimant withdraws his or her Notice of Disagreement as a result of this review process, VA will proceed with the traditional appellate process by issuing a Statement of the Case.

(g) This section applies to all claims in which a Notice of Disagreement is filed on or after June 1, 2001.

(Authority: 38 U.S.C. 5109A and 7105(d))

[66 FR 21874, May 2, 2001]

PART 4—SCHEDULE FOR RATING DISABILITIES

Subpart A—General Policy in Rating

Sec.

- 4.1 Essentials of evaluative rating.
- 4.2 Interpretation of examination reports.
- 4.3 Resolution of reasonable doubt.
- 4.6 Evaluation of evidence.
- 4.7 Higher of two evaluations.
- 4.9 Congenital or developmental defects.
- 4.10 Functional impairment.
- 4.13 Effect of change of diagnosis.
- 4.14 Avoidance of pyramiding.
- 4.15 Total disability ratings.

- 4.16 Total disability ratings for compensation based on unemployability of the individual.
- 4.17 Total disability ratings for pension based on unemployability and age of the individual.
- 4.17a Misconduct etiology.
- 4.18 Unemployability.
- 4.19 Age in service-connected claims.
- 4.20 Analogous ratings.
- 4.21 Application of rating schedule.
- 4.22 Rating of disabilities aggravated by active service.
- 4.23 Attitude of rating officers.
- 4.24 Correspondence.
- 4.25 Combined ratings table.
- 4.26 Bilateral factor.
- 4.27 Use of diagnostic code numbers.
- 4.28 Prestabilization rating from date of discharge from service.
- 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.
- 4.30 Convalescent ratings.
- 4.31 Zero percent evaluations.

Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

- 4.40 Functional loss.
- 4.41 History of injury.
- 4.42 Complete medical examination of injury cases.
- 4.43 Osteomyelitis.
- 4.44 The bones.
- 4.45 The joints.
- 4.46 Accurate measurement.
- 4.47–4.54 [Reserved]
- 4.55 Principles of combined ratings for muscle injuries.
- 4.56 Evaluation of muscle disabilities.
- 4.57 Static foot deformities.
- 4.58 Arthritis due to strain.
- 4.59 Painful motion.
- 4.60 [Reserved]
- 4.61 Examination.
- 4.62 Circulatory disturbances.
- 4.63 Loss of use of hand or foot.
- 4.64 Loss of use of both buttocks.
- 4.65 [Reserved]
- 4.66 Sacroiliac joint.
- 4.67 Pelvic bones.
- 4.68 Amputation rule.
- 4.69 Dominant hand.
- 4.70 Inadequate examinations.
- 4.71 Measurement of ankylosis and joint motion.
- 4.71a Schedule of ratings—musculoskeletal system.
- 4.72 [Reserved]
- 4.73 Schedule of ratings—muscle injuries.

THE ORGANS OF SPECIAL SENSE

- 4.75 Examination of visual acuity.
- 4.76 Examination of field vision.
- 4.76a Computation of average concentric contraction of visual fields.

§ 4.1

- 4.77 Examination of muscle function.
- 4.78 Computing aggravation.
- 4.79 Loss of use of one eye, having only light perception.
- 4.80 Rating of one eye.
- 4.81–4.82 [Reserved]
- 4.83 Ratings at scheduled steps and distances.
- 4.83a Impairment of central visual acuity.
- 4.84 Differences between distant and near visual acuity.
- 4.84a Schedule of ratings—eye.

IMPAIRMENT OF AUDITORY ACUITY

- 4.85 Evaluation of hearing impairment.
- 4.86 Exceptional patterns of hearing impairment.
- 4.87 Schedule of ratings—ear.
- 4.87a Schedule of ratings—other sense organs.

INFECTIOUS DISEASES, IMMUNE DISORDERS AND NUTRITIONAL DEFICIENCIES

- 4.88 [Reserved]
- 4.88a Chronic fatigue syndrome.
- 4.88b Schedule of ratings—infectious diseases, immune disorders and nutritional deficiencies.
- 4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19, 1968.
- 4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

THE RESPIRATORY SYSTEM

- 4.96 Special provisions regarding evaluation of respiratory conditions.
- 4.97 Schedule of ratings—respiratory system.

THE CARDIOVASCULAR SYSTEM

- 4.100–4.103 [Reserved]
- 4.104 Schedule of ratings—cardiovascular system.

THE DIGESTIVE SYSTEM

- 4.110 Ulcers.
- 4.111 Postgastroctomy syndromes.
- 4.112 Weight loss.
- 4.113 Coexisting abdominal conditions.
- 4.114 Schedule of ratings—digestive system.

THE GENITOURINARY SYSTEM

- 4.115 Nephritis.
- 4.115a Ratings of the genitourinary system—dysfunctions.
- 4.115b Ratings of the genitourinary system—diagnoses.

GYNCOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

- 4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

38 CFR Ch. I (7–1–01 Edition)

THE HEMIC AND LYMPHATIC SYSTEMS

- 4.117 Schedule of ratings—hemic and lymphatic systems.

THE SKIN

- 4.118 Schedule of ratings—skin.

THE ENDOCRINE SYSTEM

- 4.119 Schedule of ratings—endocrine system.

NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

- 4.120 Evaluations by comparison.
- 4.121 Identification of epilepsy.
- 4.122 Psychomotor epilepsy.
- 4.123 Neuritis, cranial or peripheral.
- 4.124 Neuralgia, cranial or peripheral.
- 4.124a Schedule of ratings—neurological conditions and convulsive disorders.

MENTAL DISORDERS

- 4.125 Diagnosis of mental disorders.
- 4.126 Evaluation of disability from mental disorders.
- 4.127 Mental retardation and personality disorders.
- 4.128 Convalescence ratings following extended hospitalization.
- 4.129 Mental disorders due to traumatic stress.
- 4.130 Schedule of ratings—mental disorders.

DENTAL AND ORAL CONDITIONS

- 4.149 [Reserved]
- 4.150 Schedule of ratings—dental and oral conditions.

APPENDIX A TO PART 4—TABLE OF AMENDMENTS AND EFFECTIVE DATES SINCE 1946

APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES

APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

AUTHORITY: 38 U.S.C. 1155, unless otherwise noted.

SOURCE: 29 FR 6718, May 22, 1964, unless otherwise noted.

Subpart A—General Policy in Rating

§ 4.1 Essentials of evaluative rating.

This rating schedule is primarily a guide in the evaluation of disability resulting from all types of diseases and injuries encountered as a result of or incident to military service. The percentage ratings represent as far as can practicably be determined the average impairment in earning capacity resulting from such diseases and injuries and

Department of Veterans Affairs

§ 4.10

their residual conditions in civil occupations. Generally, the degrees of disability specified are considered adequate to compensate for considerable loss of working time from exacerbations or illnesses proportionate to the severity of the several grades of disability. For the application of this schedule, accurate and fully descriptive medical examinations are required, with emphasis upon the limitation of activity imposed by the disabling condition. Over a period of many years, a veteran's disability claim may require reratings in accordance with changes in laws, medical knowledge and his or her physical or mental condition. It is thus essential, both in the examination and in the evaluation of disability, that each disability be viewed in relation to its history.

[41 FR 11292, Mar. 18, 1976]

§ 4.2 Interpretation of examination reports.

Different examiners, at different times, will not describe the same disability in the same language. Features of the disability which must have persisted unchanged may be overlooked or a change for the better or worse may not be accurately appreciated or described. It is the responsibility of the rating specialist to interpret reports of examination in the light of the whole recorded history, reconciling the various reports into a consistent picture so that the current rating may accurately reflect the elements of disability present. Each disability must be considered from the point of view of the veteran working or seeking work. If a diagnosis is not supported by the findings on the examination report or if the report does not contain sufficient detail, it is incumbent upon the rating board to return the report as inadequate for evaluation purposes.

[41 FR 11292, Mar. 18, 1976]

§ 4.3 Resolution of reasonable doubt.

It is the defined and consistently applied policy of the Department of Veterans Affairs to administer the law under a broad interpretation, consistent, however, with the facts shown in every case. When after careful consideration of all procurable and assem-

bled data, a reasonable doubt arises regarding the degree of disability such doubt will be resolved in favor of the claimant. See § 3.102 of this chapter.

[40 FR 42535, Sept. 15, 1975]

§ 4.6 Evaluation of evidence.

The element of the weight to be accorded the character of the veteran's service is but one factor entering into the considerations of the rating boards in arriving at determinations of the evaluation of disability. Every element in any way affecting the probative value to be assigned to the evidence in each individual claim must be thoroughly and conscientiously studied by each member of the rating board in the light of the established policies of the Department of Veterans Affairs to the end that decisions will be equitable and just as contemplated by the requirements of the law.

§ 4.7 Higher of two evaluations.

Where there is a question as to which of two evaluations shall be applied, the higher evaluation will be assigned if the disability picture more nearly approximates the criteria required for that rating. Otherwise, the lower rating will be assigned.

§ 4.9 Congenital or developmental defects.

Mere congenital or developmental defects, absent, displaced or supernumerary parts, refractive error of the eye, personality disorder and mental deficiency are not diseases or injuries in the meaning of applicable legislation for disability compensation purposes.

[41 FR 11292, Mar. 18, 1976]

§ 4.10 Functional impairment.

The basis of disability evaluations is the ability of the body as a whole, or of the psyche, or of a system or organ of the body to function under the ordinary conditions of daily life including employment. Whether the upper or lower extremities, the back or abdominal wall, the eyes or ears, or the cardiovascular, digestive, or other system, or psyche are affected, evaluations are based upon lack of usefulness, of these

§ 4.13

parts or systems, especially in self-support. This imposes upon the medical examiner the responsibility of furnishing, in addition to the etiological, anatomical, pathological, laboratory and prognostic data required for ordinary medical classification, full description of the effects of disability upon the person's ordinary activity. In this connection, it will be remembered that a person may be too disabled to engage in employment although he or she is up and about and fairly comfortable at home or upon limited activity.

[41 FR 11292, Mar. 18, 1976]

§ 4.13 Effect of change of diagnosis.

The repercussion upon a current rating of service connection when change is made of a previously assigned diagnosis or etiology must be kept in mind. The aim should be the reconciliation and continuance of the diagnosis or etiology upon which service connection for the disability had been granted. The relevant principle enunciated in § 4.125, entitled "Diagnosis of mental disorders," should have careful attention in this connection. When any change in evaluation is to be made, the rating agency should assure itself that there has been an actual change in the conditions, for better or worse, and not merely a difference in thoroughness of the examination or in use of descriptive terms. This will not, of course, preclude the correction of erroneous ratings, nor will it preclude assignment of a rating in conformity with § 4.7.

[29 FR 6718, May 22, 1964, as amended at 61 FR 52700, Oct. 8, 1996]

§ 4.14 Avoidance of pyramiding.

The evaluation of the same disability under various diagnoses is to be avoided. Disability from injuries to the muscles, nerves, and joints of an extremity may overlap to a great extent, so that special rules are included in the appropriate bodily system for their evaluation. Dyspnea, tachycardia, nervousness, fatigability, etc., may result from many causes; some may be service connected, others, not. Both the use of manifestations not resulting from service-connected disease or injury in establishing the service-connected eval-

38 CFR Ch. I (7-1-01 Edition)

uation, and the evaluation of the same manifestation under different diagnoses are to be avoided.

§ 4.15 Total disability ratings.

The ability to overcome the handicap of disability varies widely among individuals. The rating, however, is based primarily upon the average impairment in earning capacity, that is, upon the economic or industrial handicap which must be overcome and not from individual success in overcoming it. However, full consideration must be given to unusual physical or mental effects in individual cases, to peculiar effects of occupational activities, to defects in physical or mental endowment preventing the usual amount of success in overcoming the handicap of disability and to the effect of combinations of disability. Total disability will be considered to exist when there is present any impairment of mind or body which is sufficient to render it impossible for the average person to follow a substantially gainful occupation; *Provided*, That permanent total disability shall be taken to exist when the impairment is reasonably certain to continue throughout the life of the disabled person. The following will be considered to be permanent total disability: the permanent loss of the use of both hands, or of both feet, or of one hand and one foot, or of the sight of both eyes, or becoming permanently helpless or permanently bedridden. Other total disability ratings are scheduled in the various bodily systems of this schedule.

§ 4.16 Total disability ratings for compensation based on unemployability of the individual.

(a) Total disability ratings for compensation may be assigned, where the schedular rating is less than total, when the disabled person is, in the judgment of the rating agency, unable to secure or follow a substantially gainful occupation as a result of service-connected disabilities: *Provided* That, if there is only one such disability, this disability shall be ratable at 60 percent or more, and that, if there are two or more disabilities, there shall be at least one disability ratable at 40

percent or more, and sufficient additional disability to bring the combined rating to 70 percent or more. For the above purpose of one 60 percent disability, or one 40 percent disability in combination, the following will be considered as one disability: (1) Disabilities of one or both upper extremities, or of one or both lower extremities, including the bilateral factor, if applicable, (2) disabilities resulting from common etiology or a single accident, (3) disabilities affecting a single body system, e.g. orthopedic, digestive, respiratory, cardiovascular-renal, neuropsychiatric, (4) multiple injuries incurred in action, or (5) multiple disabilities incurred as a prisoner of war. It is provided further that the existence or degree of nonservice-connected disabilities or previous unemployability status will be disregarded where the percentages referred to in this paragraph for the service-connected disability or disabilities are met and in the judgment of the rating agency such service-connected disabilities render the veteran unemployable. Marginal employment shall not be considered substantially gainful employment. For purposes of this section, marginal employment generally shall be deemed to exist when a veteran's earned annual income does not exceed the amount established by the U.S. Department of Commerce, Bureau of the Census, as the poverty threshold for one person. Marginal employment may also be held to exist, on a facts found basis (includes but is not limited to employment in a protected environment such as a family business or sheltered workshop), when earned annual income exceeds the poverty threshold. Consideration shall be given in all claims to the nature of the employment and the reason for termination.

(Authority: 38 U.S.C. 501)

(b) It is the established policy of the Department of Veterans Affairs that all veterans who are unable to secure and follow a substantially gainful occupation by reason of service-connected disabilities shall be rated totally disabled. Therefore, rating boards should submit to the Director, Compensation and Pension Service, for extra-sched-

ular consideration all cases of veterans who are unemployable by reason of service-connected disabilities, but who fail to meet the percentage standards set forth in paragraph (a) of this section. The rating board will include a full statement as to the veteran's service-connected disabilities, employment history, educational and vocational attainment and all other factors having a bearing on the issue.

[40 FR 42535, Sept. 15, 1975, as amended at 54 FR 4281, Jan. 30, 1989; 55 FR 31580, Aug. 3, 1990; 58 FR 39664, July 26, 1993; 61 FR 52700, Oct. 8, 1996]

§ 4.17 Total disability ratings for pension based on unemployability and age of the individual.

All veterans who are basically eligible and who are unable to secure and follow a substantially gainful occupation by reason of disabilities which are likely to be permanent shall be rated as permanently and totally disabled. For the purpose of pension, the permanence of the percentage requirements of § 4.16 is a requisite. When the percentage requirements are met, and the disabilities involved are of a permanent nature, a rating of permanent and total disability will be assigned if the veteran is found to be unable to secure and follow substantially gainful employment by reason of such disability. Prior employment or unemployment status is immaterial if in the judgment of the rating board the veteran's disabilities render him or her unemployable. In making such determinations, the following guidelines will be used:

(a) Marginal employment, for example, as a self-employed farmer or other person, while employed in his or her own business, or at odd jobs or while employed at less than half the usual remuneration will not be considered incompatible with a determination of unemployability, if the restriction, as to securing or retaining better employment, is due to disability.

(b) Claims of all veterans who fail to meet the percentage standards but who meet the basic entitlement criteria and are unemployable, will be referred by the rating board to the Adjudication

§ 4.17a

Officer under § 3.321(b)(2) of this chapter.

(Authority: 38 U.S.C. 1155; 38 U.S.C. 3102)

[43 FR 45348, Oct. 2, 1978, as amended at 56 FR 57985, Nov. 15, 1991]

§ 4.17a Misconduct etiology.

A permanent and total disability rating under the provisions of §§ 4.15, 4.16 and 4.17 will not be precluded by reason of the coexistence of misconduct disability when:

(a) A veteran, regardless of employment status, also has innocently acquired 100 percent disability, or

(b) Where unemployable, the veteran has other disabilities innocently acquired which meet the percentage requirements of §§ 4.16 and 4.17 and would render, in the judgment of the rating agency, the average person unable to secure or follow a substantially gainful occupation.

[40 FR 42536, Sept. 15, 1975, as amended at 43 FR 45349, Oct. 2, 1978]

§ 4.18 Unemployability.

A veteran may be considered as unemployable upon termination of employment which was provided on account of disability, or in which special consideration was given on account of the same, when it is satisfactorily shown that he or she is unable to secure further employment. With amputations, sequelae of fractures and other residuals of traumatism shown to be of static character, a showing of continuous unemployability from date of incurrence, or the date the condition reached the stabilized level, is a general requirement in order to establish the fact that present unemployability is the result of the disability. However, consideration is to be given to the circumstances of employment in individual claims, and, if the employment was only occasional, intermittent, try-out or unsuccessful, or eventually terminated on account of the disability, present unemployability may be attributed to the static disability. Where unemployability for pension previously has been established on the basis of combined service-connected and non-service-connected disabilities and the service-connected disability or disabili-

38 CFR Ch. I (7-1-01 Edition)

ties have increased in severity, § 4.16 is for consideration.

[40 FR 42536, Sept. 15, 1975, as amended at 43 FR 45349, Oct. 2, 1978]

§ 4.19 Age in service-connected claims.

Age may not be considered as a factor in evaluating service-connected disability; and unemployability, in service-connected claims, associated with advancing age or intercurrent disability, may not be used as a basis for a total disability rating. Age, as such, is a factor only in evaluations of disability not resulting from service, i.e., for the purposes of pension.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

§ 4.20 Analogous ratings.

When an unlisted condition is encountered it will be permissible to rate under a closely related disease or injury in which not only the functions affected, but the anatomical localization and symptomatology are closely analogous. Conjectural analogies will be avoided, as will the use of analogous ratings for conditions of doubtful diagnosis, or for those not fully supported by clinical and laboratory findings. Nor will ratings assigned to organic diseases and injuries be assigned by analogy to conditions of functional origin.

§ 4.21 Application of rating schedule.

In view of the number of atypical instances it is not expected, especially with the more fully described grades of disabilities, that all cases will show all the findings specified. Findings sufficiently characteristic to identify the disease and the disability therefrom, and above all, coordination of rating with impairment of function will, however, be expected in all instances.

[41 FR 11293, Mar. 18, 1976]

§ 4.22 Rating of disabilities aggravated by active service.

In cases involving aggravation by active service, the rating will reflect only the degree of disability over and above the degree existing at the time of entrance into the active service, whether the particular condition was noted at the time of entrance into the active service, or it is determined upon the

evidence of record to have existed at that time. It is necessary therefore, in all cases of this character to deduct from the present degree of disability the degree, if ascertainable, of the disability existing at the time of entrance into active service, in terms of the rating schedule, except that if the disability is total (100 percent) no deduction will be made. The resulting difference will be recorded on the rating sheet. If the degree of disability at the time of entrance into the service is not ascertainable in terms of the schedule, no deduction will be made.

§ 4.23 Attitude of rating officers.

It is to be remembered that the majority of applicants are disabled persons who are seeking benefits of law to which they believe themselves entitled. In the exercise of his or her functions, rating officers must not allow their personal feelings to intrude; an antagonistic, critical, or even abusive attitude on the part of a claimant should not in any instance influence the officers in the handling of the case. Fairness and courtesy must at all times be shown to applicants by all employees whose duties bring them in contact, directly or indirectly, with the Department's claimants.

[41 FR 11292, Mar. 18, 1976]

§ 4.24 Correspondence.

All correspondence relative to the interpretation of the schedule for rating disabilities, requests for advisory opinions, questions regarding lack of clarity or application to individual cases involving unusual difficulties, will be addressed to the Director, Compensation and Pension Service. A clear statement will be made of the point or points upon which information is desired, and the complete case file will be simultaneously forwarded to Central Office. Rating agencies will assure themselves that the recent report of physical examination presents an adequate picture of the claimant's condition. Claims in regard to which the schedule evaluations are considered inadequate or excessive, and errors in the schedule will be similarly brought to attention.

[41 FR 11292, Mar. 18, 1976]

§ 4.25 Combined ratings table.

Table I, Combined Ratings Table, results from the consideration of the efficiency of the individual as affected first by the most disabling condition, then by the less disabling condition, then by other less disabling conditions, if any, in the order of severity. Thus, a person having a 60 percent disability is considered 40 percent efficient. Proceeding from this 40 percent efficiency, the effect of a further 30 percent disability is to leave only 70 percent of the efficiency remaining after consideration of the first disability, or 28 percent efficiency altogether. The individual is thus 72 percent disabled, as shown in table I opposite 60 percent and under 30 percent.

(a) To use table I, the disabilities will first be arranged in the exact order of their severity, beginning with the greatest disability and then combined with use of table I as hereinafter indicated. For example, if there are two disabilities, the degree of one disability will be read in the left column and the degree of the other in the top row, whichever is appropriate. The figures appearing in the space where the column and row intersect will represent the combined value of the two. This combined value will then be converted to the nearest number divisible by 10, and combined values ending in 5 will be adjusted upward. Thus, with a 50 percent disability and a 30 percent disability, the combined value will be found to be 65 percent, but the 65 percent must be converted to 70 percent to represent the final degree of disability. Similarly, with a disability of 40 percent, and another disability of 20 percent, the combined value is found to be 52 percent, but the 52 percent must be converted to the nearest degree divisible by 10, which is 50 percent. If there are more than two disabilities, the disabilities will also be arranged in the exact order of their severity and the combined value for the first two will be found as previously described for two disabilities. The combined value, exactly as found in table I, will be combined with the degree of the third disability (in order of severity). The combined value for the three disabilities will be found in the space where the column and row intersect, and if there

§ 4.25

38 CFR Ch. I (7-1-01 Edition)

are only three disabilities will be converted to the nearest degree divisible by 10, adjusting final 5's upward. Thus, if there are three disabilities ratable at 60 percent, 40 percent, and 20 percent, respectively, the combined value for the first two will be found opposite 60 and under 40 and is 76 percent. This 76 will be combined with 20 and the combined value for the three is 81 percent. This combined value will be converted to the nearest degree divisible by 10 which is 80 percent. The same procedure will be employed when there are four or more disabilities. (See table I).

(b) Except as otherwise provided in this schedule, the disabilities arising from a single disease entity, e.g., arthritis, multiple sclerosis, cerebrovascular accident, etc., are to be rated separately as are all other disabling conditions, if any. All disabilities are then to be combined as described in paragraph (a) of this section. The conversion to the nearest degree divisible by 10 will be done only once per rating decision, will follow the combining of all disabilities, and will be the last procedure in determining the combined degree of disability.

TABLE I—COMBINED RATINGS TABLE
[10 combined with 10 is 19]

| | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 | 90 |
|----------|----|----|----|----|----|----|----|----|----|
| 19 | 27 | 35 | 43 | 51 | 60 | 68 | 76 | 84 | 92 |
| 20 | 28 | 36 | 44 | 52 | 60 | 68 | 76 | 84 | 92 |
| 21 | 29 | 37 | 45 | 53 | 61 | 68 | 76 | 84 | 92 |
| 22 | 30 | 38 | 45 | 53 | 61 | 69 | 77 | 84 | 92 |
| 23 | 31 | 38 | 46 | 54 | 62 | 69 | 77 | 85 | 92 |
| 24 | 32 | 39 | 47 | 54 | 62 | 70 | 77 | 85 | 92 |
| 25 | 33 | 40 | 48 | 55 | 63 | 70 | 78 | 85 | 93 |
| 26 | 33 | 41 | 48 | 56 | 63 | 70 | 78 | 85 | 93 |
| 27 | 34 | 42 | 49 | 56 | 64 | 71 | 78 | 85 | 93 |
| 28 | 35 | 42 | 50 | 57 | 64 | 71 | 78 | 86 | 93 |
| 29 | 36 | 43 | 50 | 57 | 65 | 72 | 79 | 86 | 93 |
| 30 | 37 | 44 | 51 | 58 | 65 | 72 | 79 | 86 | 93 |
| 31 | 38 | 45 | 52 | 59 | 66 | 72 | 79 | 86 | 93 |
| 32 | 39 | 46 | 52 | 59 | 66 | 73 | 80 | 86 | 93 |
| 33 | 40 | 46 | 53 | 60 | 67 | 73 | 80 | 87 | 93 |
| 34 | 41 | 47 | 54 | 60 | 67 | 74 | 80 | 87 | 93 |
| 35 | 42 | 48 | 55 | 61 | 68 | 74 | 81 | 87 | 94 |
| 36 | 42 | 49 | 55 | 62 | 68 | 74 | 81 | 87 | 94 |
| 37 | 43 | 50 | 56 | 62 | 69 | 75 | 81 | 87 | 94 |
| 38 | 44 | 50 | 57 | 63 | 69 | 75 | 81 | 88 | 94 |
| 39 | 45 | 51 | 57 | 63 | 70 | 76 | 82 | 88 | 94 |
| 40 | 46 | 52 | 58 | 64 | 70 | 76 | 82 | 88 | 94 |
| 41 | 47 | 53 | 59 | 65 | 71 | 76 | 82 | 88 | 94 |
| 42 | 48 | 54 | 59 | 65 | 71 | 77 | 83 | 88 | 94 |
| 43 | 49 | 54 | 60 | 66 | 72 | 77 | 83 | 89 | 94 |
| 44 | 50 | 55 | 61 | 66 | 72 | 78 | 83 | 89 | 94 |
| 45 | 51 | 56 | 62 | 67 | 73 | 78 | 84 | 89 | 95 |
| 46 | 51 | 57 | 62 | 68 | 73 | 78 | 84 | 89 | 95 |
| 47 | 52 | 58 | 63 | 68 | 74 | 79 | 84 | 89 | 95 |
| 48 | 53 | 58 | 64 | 69 | 74 | 79 | 84 | 90 | 95 |
| 49 | 54 | 59 | 64 | 69 | 75 | 80 | 85 | 90 | 95 |
| 50 | 55 | 60 | 65 | 70 | 75 | 80 | 85 | 90 | 95 |
| 51 | 56 | 61 | 66 | 71 | 76 | 80 | 85 | 90 | 95 |
| 52 | 57 | 62 | 66 | 71 | 76 | 81 | 86 | 90 | 95 |
| 53 | 58 | 62 | 67 | 72 | 77 | 81 | 86 | 91 | 95 |
| 54 | 59 | 63 | 68 | 72 | 77 | 82 | 86 | 91 | 95 |
| 55 | 60 | 64 | 69 | 73 | 78 | 82 | 87 | 91 | 96 |
| 56 | 60 | 65 | 69 | 74 | 78 | 82 | 87 | 91 | 96 |
| 57 | 61 | 66 | 70 | 74 | 79 | 83 | 87 | 91 | 96 |
| 58 | 62 | 66 | 71 | 75 | 79 | 83 | 87 | 92 | 96 |
| 59 | 63 | 67 | 71 | 75 | 80 | 84 | 88 | 92 | 96 |
| 60 | 64 | 68 | 72 | 76 | 80 | 84 | 88 | 92 | 96 |
| 61 | 65 | 69 | 73 | 77 | 81 | 84 | 88 | 92 | 96 |
| 62 | 66 | 70 | 73 | 77 | 81 | 85 | 89 | 92 | 96 |
| 63 | 67 | 70 | 74 | 78 | 82 | 85 | 89 | 93 | 96 |
| 64 | 68 | 71 | 75 | 78 | 82 | 86 | 89 | 93 | 96 |
| 65 | 69 | 72 | 76 | 79 | 83 | 86 | 90 | 93 | 97 |
| 66 | 69 | 73 | 76 | 80 | 83 | 86 | 90 | 93 | 97 |
| 67 | 70 | 74 | 77 | 80 | 84 | 87 | 90 | 93 | 97 |
| 68 | 71 | 74 | 78 | 81 | 84 | 87 | 90 | 94 | 97 |
| 69 | 72 | 75 | 78 | 81 | 85 | 88 | 91 | 94 | 97 |

TABLE I—COMBINED RATINGS TABLE—Continued
 [10 combined with 10 is 19]

| | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 | 90 |
|----------|----|----|----|----|----|----|----|----|----|
| 70 | 73 | 76 | 79 | 82 | 85 | 88 | 91 | 94 | 97 |
| 71 | 74 | 77 | 80 | 83 | 86 | 88 | 91 | 94 | 97 |
| 72 | 75 | 78 | 80 | 83 | 86 | 89 | 92 | 94 | 97 |
| 73 | 76 | 78 | 81 | 84 | 87 | 89 | 92 | 95 | 97 |
| 74 | 77 | 79 | 82 | 84 | 87 | 90 | 92 | 95 | 97 |
| 75 | 78 | 80 | 83 | 85 | 88 | 90 | 93 | 95 | 98 |
| 76 | 78 | 81 | 83 | 86 | 88 | 90 | 93 | 95 | 98 |
| 77 | 79 | 82 | 84 | 86 | 89 | 91 | 93 | 95 | 98 |
| 78 | 80 | 82 | 85 | 87 | 89 | 91 | 93 | 96 | 98 |
| 79 | 81 | 83 | 85 | 87 | 90 | 92 | 94 | 96 | 98 |
| 80 | 82 | 84 | 86 | 88 | 90 | 92 | 94 | 96 | 98 |
| 81 | 83 | 85 | 87 | 89 | 91 | 92 | 94 | 96 | 98 |
| 82 | 84 | 86 | 87 | 89 | 91 | 93 | 95 | 96 | 98 |
| 83 | 85 | 86 | 88 | 90 | 92 | 93 | 95 | 97 | 98 |
| 84 | 86 | 87 | 89 | 90 | 92 | 94 | 95 | 97 | 98 |
| 85 | 87 | 88 | 90 | 91 | 93 | 94 | 96 | 97 | 99 |
| 86 | 87 | 89 | 90 | 92 | 93 | 94 | 96 | 97 | 99 |
| 87 | 88 | 90 | 91 | 92 | 94 | 95 | 96 | 97 | 99 |
| 88 | 89 | 90 | 92 | 93 | 94 | 95 | 96 | 98 | 99 |
| 89 | 90 | 91 | 92 | 93 | 95 | 96 | 87 | 38 | 99 |
| 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 |
| 91 | 92 | 93 | 94 | 95 | 96 | 96 | 97 | 98 | 99 |
| 92 | 93 | 94 | 94 | 95 | 96 | 97 | 98 | 98 | 99 |
| 93 | 94 | 94 | 95 | 96 | 97 | 97 | 98 | 99 | 99 |
| 94 | 95 | 95 | 96 | 96 | 97 | 98 | 98 | 99 | 99 |

(Authority: 38 U.S.C. 1155)

[41 FR 11293, Mar. 18, 1976, as amended at 54 FR 27161, June 28, 1989; 54 FR 36029, Aug. 31, 1989]

§ 4.26 Bilateral factor.

When a partial disability results from disease or injury of both arms, or of both legs, or of paired skeletal muscles, the ratings for the disabilities of the right and left sides will be combined as usual, and 10 percent of this value will be added (i.e., not combined) before proceeding with further combinations, or converting to degree of disability. The bilateral factor will be applied to such bilateral disabilities before other combinations are carried out and the rating for such disabilities including the bilateral factor in this section will be treated as 1 disability for the purpose of arranging in order of severity and for all further combinations. For example, with disabilities evaluated at 60 percent, 20 percent, 10 percent and 10 percent (the two 10's representing bilateral disabilities), the order of severity would be 60, 21 and 20. The 60 and 21 combine to 68 percent and the 68 and 20 to 74 percent, converted to

70 percent as the final degree of disability.

(a) The use of the terms “arms” and “legs” is not intended to distinguish between the arm, forearm and hand, or the thigh, leg, and foot, but relates to the upper extremities and lower extremities as a whole. Thus with a compensable disability of the right thigh, for example, amputation, and one of the left foot, for example, pes planus, the bilateral factor applies, and similarly whenever there are compensable disabilities affecting use of paired extremities regardless of location or specified type of impairment.

(b) The correct procedure when applying the bilateral factor to disabilities affecting both upper extremities and both lower extremities is to combine the ratings of the disabilities affecting the 4 extremities in the order of their individual severity and apply the bilateral factor by adding, not combining, 10 percent of the combined value thus attained.

(c) The bilateral factor is not applicable unless there is partial disability of compensable degree in each of 2 paired extremities, or paired skeletal muscles.

§ 4.27

§ 4.27 Use of diagnostic code numbers.

The diagnostic code numbers appearing opposite the listed ratable disabilities are arbitrary numbers for the purpose of showing the basis of the evaluation assigned and for statistical analysis in the Department of Veterans Affairs, and as will be observed, extend from 5000 to a possible 9999. Great care will be exercised in the selection of the applicable code number and in its citation on the rating sheet. No other numbers than these listed or hereafter furnished are to be employed for rating purposes, with an exception as described in this section, as to unlisted conditions. When an unlisted disease, injury, or residual condition is encountered, requiring rating by analogy, the diagnostic code number will be "built-up" as follows: The first 2 digits will be selected from that part of the schedule most closely identifying the part, or system, of the body involved; the last 2 digits will be "99" for all unlisted conditions. This procedure will facilitate a close check of new and unlisted conditions, rated by analogy. In the selection of code numbers, injuries will generally be represented by the number assigned to the residual condition on the basis of which the rating is determined. With diseases, preference is to be given to the number assigned to the disease itself; if the rating is determined on the basis of residual conditions, the number appropriate to the residual condition will be added, preceded by a hyphen. Thus, rheumatoid (atrophic) arthritis rated as ankylosis of the lumbar spine should be coded "5002-5289." In this way, the exact source of each rating can be easily identified. In the citation of disabilities on rating sheets, the diagnostic terminology will be that of the medical examiner, with no attempt to translate the terms into schedule nomenclature. Residuals of diseases or therapeutic procedures will not be cited without reference to the basic disease.

[41 FR 11293, Mar. 18, 1976]

§ 4.28 Prestabilization rating from date of discharge from service.

The following ratings may be assigned, in lieu of ratings prescribed elsewhere, under the conditions stated

for disability from any disease or injury. The prestabilization rating is not to be assigned in any case in which a total rating is immediately assignable under the regular provisions of the schedule or on the basis of individual unemployability. The prestabilization 50-percent rating is not to be used in any case in which a rating of 50 percent or more is immediately assignable under the regular provisions.

| | Rating |
|---|--------|
| Unstabilized condition with severe disability— Substantially gainful employment is not feasible or advisable | 100 |
| Unhealed or incompletely healed wounds or injuries— Material impairment of employability likely .. | 50 |

NOTE (1): Department of Veterans Affairs examination is not required prior to assignment of prestabilization ratings; however, the fact that examination was accomplished will not preclude assignment of these benefits. Prestabilization ratings are for assignment in the immediate postdischarge period. They will continue for a 12-month period following discharge from service. However, prestabilization ratings may be changed to a regular schedular total rating or one authorizing a greater benefit at any time. In each prestabilization rating an examination will be requested to be accomplished not earlier than 6 months nor more than 12 months following discharge. In those prestabilization ratings in which following examination reduction in evaluation is found to be warranted, the higher evaluation will be continued to the end of the 12th month following discharge or to the end of the period provided under § 3.105(e) of this chapter, whichever is later. Special monthly compensation should be assigned concurrently in these cases whenever records are adequate to establish entitlement.

NOTE (2): Diagnosis of disease, injury, or residuals will be cited, with diagnostic code number assigned from this rating schedule for conditions listed therein.

[35 FR 11906, July 24, 1970]

§ 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established that a service-connected disability has required hospital treatment in a Department of Veterans Affairs or an approved hospital for a period in excess of 21 days or

Department of Veterans Affairs

§ 4.30

hospital observation at Department of Veterans Affairs expense for a service-connected disability for a period in excess of 21 days.

(a) Subject to the provisions of paragraphs (d), (e), and (f) of this section this increased rating will be effective the first day of continuous hospitalization and will be terminated effective the last day of the month of hospital discharge (regular discharge or release to non-bed care) or effective the last day of the month of termination of treatment or observation for the service-connected disability. A temporary release which is approved by an attending Department of Veterans Affairs physician as part of the treatment plan will not be considered an absence.

(1) An authorized absence in excess of 4 days which begins during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the first day of such authorized absence. An authorized absence of 4 days or less which results in a total of more than 8 days of authorized absence during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the ninth day of authorized absence.

(2) Following a period of hospitalization in excess of 21 days, an authorized absence in excess of 14 days or a third consecutive authorized absence of 14 days will be regarded as the equivalent of hospital discharge and will interrupt hospitalization effective on the last day of the month in which either the authorized absence in excess of 14 days or the third 14 day period begins, except where there is a finding that convalescence is required as provided by paragraph (e) or (f) of this section. The termination of these total ratings will not be subject to § 3.105(e) of this chapter.

(b) Notwithstanding that hospital admission was for disability not connected with service, if during such hospitalization, hospital treatment for a service-connected disability is instituted and continued for a period in excess of 21 days, the increase to a total rating will be granted from the first day of such treatment. If service connection for the disability under treatment is granted after hospital admission, the rating will be from the first

day of hospitalization if otherwise in order.

(c) The assignment of a total disability rating on the basis of hospital treatment or observation will not preclude the assignment of a total disability rating otherwise in order under other provisions of the rating schedule, and consideration will be given to the propriety of such a rating in all instances and to the propriety of its continuance after discharge. Particular attention, with a view to proper rating under the rating schedule, is to be given to the claims of veterans discharged from hospital, regardless of length of hospitalization, with indications on the final summary of expected confinement to bed or house, or to inability to work with requirement of frequent care of physician or nurse at home.

(d) On these total ratings Department of Veterans Affairs regulations governing effective dates for increased benefits will control.

(e) The total hospital rating if convalescence is required may be continued for periods of 1, 2, or 3 months in addition to the period provided in paragraph (a) of this section.

(f) Extension of periods of 1, 2 or 3 months beyond the initial 3 months may be made upon approval of the Adjudication Officer.

(g) Meritorious claims of veterans who are discharged from the hospital with less than the required number of days but need post-hospital care and a prolonged period of convalescence will be referred to the Director, Compensation and Pension Service, under § 3.321(b)(1) of this chapter.

[29 FR 6718, May 22, 1964, as amended at 41 FR 11294, Mar. 18, 1976; 41 FR 34256, Aug. 13, 1976; 54 FR 4281, Jan. 30, 1989; 54 FR 34981, Aug. 23, 1989]

§ 4.30 Convalescent ratings.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established by report at hospital discharge (regular discharge or release to non-bed care) or outpatient release that entitlement is warranted under paragraph (a) (1), (2) or (3) of this section effective the date of hospital admission or outpatient treatment and

§4.31

continuing for a period of 1, 2, or 3 months from the first day of the month following such hospital discharge or outpatient release. The termination of these total ratings will not be subject to §3.105(e) of this chapter. Such total rating will be followed by appropriate schedular evaluations. When the evidence is inadequate to assign a schedular evaluation, a physical examination will be scheduled and considered prior to the termination of a total rating under this section.

(a) Total ratings will be assigned under this section if treatment of a service-connected disability resulted in:

(1) Surgery necessitating at least one month of convalescence (Effective as to outpatient surgery March 1, 1989.)

(2) Surgery with severe postoperative residuals such as incompletely healed surgical wounds, stumps of recent amputations, therapeutic immobilization of one major joint or more, application of a body cast, or the necessity for house confinement, or the necessity for continued use of a wheelchair or crutches (regular weight-bearing prohibited). (Effective as to outpatient surgery March 1, 1989.)

(3) Immobilization by cast, without surgery, of one major joint or more. (Effective as to outpatient treatment March 10, 1976.)

A reduction in the total rating will not be subject to §3.105(e) of this chapter. The total rating will be followed by an open rating reflecting the appropriate schedular evaluation; where the evidence is inadequate to assign the schedular evaluation, a physical examination will be scheduled prior to the end of the total rating period.

(b) A total rating under this section will require full justification on the rating sheet and may be extended as follows:

(1) Extensions of 1, 2 or 3 months beyond the initial 3 months may be made under paragraph (a) (1), (2) or (3) of this section.

(2) Extensions of 1 or more months up to 6 months beyond the initial 6 months period may be made under paragraph (a) (2) or (3) of this section

38 CFR Ch. I (7-1-01 Edition)

upon approval of the Adjudication Officer.

[41 FR 34256, Aug. 13, 1976, as amended at 54 FR 4281, Jan. 30, 1989]

§4.31 Zero percent evaluations.

In every instance where the schedule does not provide a zero percent evaluation for a diagnostic code, a zero percent evaluation shall be assigned when the requirements for a compensable evaluation are not met.

[58 FR 52018, Oct. 6, 1993]

Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

§4.40 Functional loss.

Disability of the musculoskeletal system is primarily the inability, due to damage or infection in parts of the system, to perform the normal working movements of the body with normal excursion, strength, speed, coordination and endurance. It is essential that the examination on which ratings are based adequately portray the anatomical damage, and the functional loss, with respect to all these elements. The functional loss may be due to absence of part, or all, of the necessary bones, joints and muscles, or associated structures, or to deformity, adhesions, defective innervation, or other pathology, or it may be due to pain, supported by adequate pathology and evidenced by the visible behavior of the claimant undertaking the motion. Weakness is as important as limitation of motion, and a part which becomes painful on use must be regarded as seriously disabled. A little used part of the musculoskeletal system may be expected to show evidence of disuse, either through atrophy, the condition of the skin, absence of normal callosity or the like.

§4.41 History of injury.

In considering the residuals of injury, it is essential to trace the medical-industrial history of the disabled person from the original injury, considering the nature of the injury and the attendant circumstances, and the requirements for, and the effect of, treatment over past periods, and the course

of the recovery to date. The duration of the initial, and any subsequent, period of total incapacity, especially periods reflecting delayed union, inflammation, swelling, drainage, or operative intervention, should be given close attention. This consideration, or the absence of clear cut evidence of injury, may result in classifying the disability as not of traumatic origin, either reflecting congenital or developmental etiology, or the effects of healed disease.

§ 4.42 Complete medical examination of injury cases.

The importance of complete medical examination of injury cases at the time of first medical examination by the Department of Veterans Affairs cannot be overemphasized. When possible, this should include complete neurological and psychiatric examination, and other special examinations indicated by the physical condition, in addition to the required general and orthopedic or surgical examinations. When complete examinations are not conducted covering all systems of the body affected by disease or injury, it is impossible to visualize the nature and extent of the service connected disability. Incomplete examination is a common cause of incorrect diagnosis, especially in the neurological and psychiatric fields, and frequently leaves the Department of Veterans Affairs in doubt as to the presence or absence of disabling conditions at the time of the examination.

§ 4.43 Osteomyelitis.

Chronic, or recurring, suppurative osteomyelitis, once clinically identified, including chronic inflammation of bone marrow, cortex, or periosteum, should be considered as a continuously disabling process, whether or not an actively discharging sinus or other obvious evidence of infection is manifest from time to time, and unless the focus is entirely removed by amputation will entitle to a permanent rating to be combined with other ratings for residual conditions, however, not exceeding amputation ratings at the site of election.

§ 4.44 The bones.

The osseous abnormalities incident to trauma or disease, such as malunion with deformity throwing abnormal stress upon, and causing malalignment of joint surfaces, should be depicted from study and observation of all available data, beginning with inception of injury or disease, its nature, degree of prostration, treatment and duration of convalescence, and progress of recovery with development of permanent residuals. With shortening of a long bone, some degree of angulation is to be expected; the extent and direction should be brought out by X-ray and observation. The direction of angulation and extent of deformity should be carefully related to strain on the neighboring joints, especially those connected with weight-bearing.

§ 4.45 The joints.

As regards the joints the factors of disability reside in reductions of their normal excursion of movements in different planes. Inquiry will be directed to these considerations:

(a) Less movement than normal (due to ankylosis, limitation or blocking, adhesions, tendon-tie-up, contracted scars, etc.).

(b) More movement than normal (from flail joint, resections, nonunion of fracture, relaxation of ligaments, etc.).

(c) Weakened movement (due to muscle injury, disease or injury of peripheral nerves, divided or lengthened tendons, etc.).

(d) Excess fatigability.

(e) Incoordination, impaired ability to execute skilled movements smoothly.

(f) Pain on movement, swelling, deformity or atrophy of disuse. Instability of station, disturbance of locomotion, interference with sitting, standing and weight-bearing are related considerations. For the purpose of rating disability from arthritis, the shoulder, elbow, wrist, hip, knee, and ankle are considered major joints; multiple involvements of the interphalangeal, metacarpal and carpal joints of the upper extremities, the interphalangeal, metatarsal and tarsal joints of the lower extremities, the cervical vertebrae, the dorsal vertebrae, and the

§4.46

lumbar vertebrae, are considered groups of minor joints, ratable on a parity with major joints. The lumbosacral articulation and both sacroiliac joints are considered to be a group of minor joints, ratable on disturbance of lumbar spine functions.

§4.46 Accurate measurement.

Accurate measurement of the length of stumps, excursion of joints, dimensions and location of scars with respect to landmarks, should be insisted on. The use of a goniometer in the measurement of limitation of motion is indispensable in examinations conducted within the Department of Veterans Affairs. Muscle atrophy must also be accurately measured and reported.

[41 FR 11294, Mar. 18, 1976]

§§ 4.47–4.54 [Reserved]

§4.55 Principles of combined ratings for muscle injuries.

(a) A muscle injury rating will not be combined with a peripheral nerve paralysis rating of the same body part, unless the injuries affect entirely different functions.

(b) For rating purposes, the skeletal muscles of the body are divided into 23 muscle groups in 5 anatomical regions: 6 muscle groups for the shoulder girdle and arm (diagnostic codes 5301 through 5306); 3 muscle groups for the forearm and hand (diagnostic codes 5307 through 5309); 3 muscle groups for the foot and leg (diagnostic codes 5310 through 5312); 6 muscle groups for the pelvic girdle and thigh (diagnostic codes 5313 through 5318); and 5 muscle groups for the torso and neck (diagnostic codes 5319 through 5323).

(c) There will be no rating assigned for muscle groups which act upon an ankylosed joint, with the following exceptions:

(1) In the case of an ankylosed knee, if muscle group XIII is disabled, it will be rated, but at the next lower level than that which would otherwise be assigned.

(2) In the case of an ankylosed shoulder, if muscle groups I and II are severely disabled, the evaluation of the shoulder joint under diagnostic code 5200 will be elevated to the level for unfavorable ankylosis, if not already as-

38 CFR Ch. I (7–1–01 Edition)

signed, but the muscle groups themselves will not be rated.

(d) The combined evaluation of muscle groups acting upon a single unankylosed joint must be lower than the evaluation for unfavorable ankylosis of that joint, except in the case of muscle groups I and II acting upon the shoulder.

(e) For compensable muscle group injuries which are in the same anatomical region but do not act on the same joint, the evaluation for the most severely injured muscle group will be increased by one level and used as the combined evaluation for the affected muscle groups.

(f) For muscle group injuries in different anatomical regions which do not act upon ankylosed joints, each muscle group injury shall be separately rated and the ratings combined under the provisions of §4.25.

(Authority: 38 U.S.C. 1155)

[62 FR 30237, June 3, 1997]

§4.56 Evaluation of muscle disabilities.

(a) An open comminuted fracture with muscle or tendon damage will be rated as a severe injury of the muscle group involved unless, for locations such as in the wrist or over the tibia, evidence establishes that the muscle damage is minimal.

(b) A through-and-through injury with muscle damage shall be evaluated as no less than a moderate injury for each group of muscles damaged.

(c) For VA rating purposes, the cardinal signs and symptoms of muscle disability are loss of power, weakness, lowered threshold of fatigue, fatigue-pain, impairment of coordination and uncertainty of movement.

(d) Under diagnostic codes 5301 through 5323, disabilities resulting from muscle injuries shall be classified as slight, moderate, moderately severe or severe as follows:

(1) *Slight disability of muscles*—(i) *Type of injury*. Simple wound of muscle without debridement or infection.

(ii) *History and complaint*. Service department record of superficial wound with brief treatment and return to duty. Healing with good functional results. No cardinal signs or symptoms of

muscle disability as defined in paragraph (c) of this section.

(iii) *Objective findings.* Minimal scar. No evidence of fascial defect, atrophy, or impaired tonus. No impairment of function or metallic fragments retained in muscle tissue.

(2) *Moderate disability of muscles—(i) Type of injury.* Through and through or deep penetrating wound of short track from a single bullet, small shell or shrapnel fragment, without explosive effect of high velocity missile, residuals of debridement, or prolonged infection.

(ii) *History and complaint.* Service department record or other evidence of in-service treatment for the wound. Record of consistent complaint of one or more of the cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, particularly lowered threshold of fatigue after average use, affecting the particular functions controlled by the injured muscles.

(iii) *Objective findings.* Entrance and (if present) exit scars, small or linear, indicating short track of missile through muscle tissue. Some loss of deep fascia or muscle substance or impairment of muscle tonus and loss of power or lowered threshold of fatigue when compared to the sound side.

(3) *Moderately severe disability of muscles—(i) Type of injury.* Through and through or deep penetrating wound by small high velocity missile or large low-velocity missile, with debridement, prolonged infection, or sloughing of soft parts, and intermuscular scarring.

(ii) *History and complaint.* Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section and, if present, evidence of inability to keep up with work requirements.

(iii) *Objective findings.* Entrance and (if present) exit scars indicating track of missile through one or more muscle groups. Indications on palpation of loss of deep fascia, muscle substance, or normal firm resistance of muscles compared with sound side. Tests of strength and endurance compared with

sound side demonstrate positive evidence of impairment.

(4) *Severe disability of muscles—(i) Type of injury.* Through and through or deep penetrating wound due to high-velocity missile, or large or multiple low velocity missiles, or with shattering bone fracture or open comminuted fracture with extensive debridement, prolonged infection, or sloughing of soft parts, intermuscular binding and scarring.

(ii) *History and complaint.* Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, worse than those shown for moderately severe muscle injuries, and, if present, evidence of inability to keep up with work requirements.

(iii) *Objective findings.* Ragged, depressed and adherent scars indicating wide damage to muscle groups in missile track. Palpation shows loss of deep fascia or muscle substance, or soft flabby muscles in wound area. Muscles swell and harden abnormally in contraction. Tests of strength, endurance, or coordinated movements compared with the corresponding muscles of the uninjured side indicate severe impairment of function. If present, the following are also signs of severe muscle disability:

(A) X-ray evidence of minute multiple scattered foreign bodies indicating intermuscular trauma and explosive effect of the missile.

(B) Adhesion of scar to one of the long bones, scapula, pelvic bones, sacrum or vertebrae, with epithelial sealing over the bone rather than true skin covering in an area where bone is normally protected by muscle.

(C) Diminished muscle excitability to pulsed electrical current in electrodiagnostic tests.

(D) Visible or measurable atrophy.

(E) Adaptive contraction of an opposing group of muscles.

(F) Atrophy of muscle groups not in the track of the missile, particularly of the trapezius and serratus in wounds of the shoulder girdle.

§ 4.57

(G) Induration or atrophy of an entire muscle following simple piercing by a projectile.

(Authority: 38 U.S.C. 1155

[62 FR 30238, June 3, 1997]

§ 4.57 Static foot deformities.

It is essential to make an initial distinction between bilateral flatfoot as a congenital or as an acquired condition. The congenital condition, with depression of the arch, but no evidence of abnormal callosities, areas of pressure, strain or demonstrable tenderness, is a congenital abnormality which is not compensable or pensionable. In the acquired condition, it is to be remembered that depression of the longitudinal arch, or the degree of depression, is not the essential feature. The attention should be given to anatomical changes, as compared to normal, in the relationship of the foot and leg, particularly to the inward rotation of the superior portion of the os calcis, medial deviation of the insertion of the Achilles tendon, the medial tilting of the upper border of the astragalus. This is an unfavorable mechanical relationship of the parts. A plumb line dropped from the middle of the patella falls inside of the normal point. The forepart of the foot is abducted, and the foot everted. The plantar surface of the foot is painful and shows demonstrable tenderness, and manipulation of the foot produces spasm of the Achilles tendon, peroneal spasm due to adhesion about the peroneal sheaths, and other evidence of pain and limited motion. The symptoms should be apparent without regard to exercise. In severe cases there is gaping of bones on the inner border of the foot, and rigid valgus position with loss of the power of inversion and adduction. Exercise with undeveloped or unbalanced musculature, producing chronic irritation, can be an aggravating factor. In the absence of trauma or other definite evidence of aggravation, service connection is not in order for pes cavus which is a typically congenital or juvenile disease.

§ 4.58 Arthritis due to strain.

With service incurred lower extremity amputation or shortening, a dis-

38 CFR Ch. I (7-1-01 Edition)

abling arthritis, developing in the same extremity, or in both lower extremities, with indications of earlier, or more severe, arthritis in the injured extremity, including also arthritis of the lumbosacral joints and lumbar spine, if associated with the leg amputation or shortening, will be considered as service incurred, provided, however, that arthritis affecting joints not directly subject to strain as a result of the service incurred amputation will not be granted service connection. This will generally require separate evaluation of the arthritis in the joints directly subject to strain. Amputation, or injury to an upper extremity, is not considered as a causative factor with subsequently developing arthritis, except in joints subject to direct strain or actually injured.

§ 4.59 Painful motion.

With any form of arthritis, painful motion is an important factor of disability, the facial expression, wincing, etc., on pressure or manipulation, should be carefully noted and definitely related to affected joints. Muscle spasm will greatly assist the identification. Sciatic neuritis is not uncommonly caused by arthritis of the spine. The intent of the schedule is to recognize painful motion with joint or periarticular pathology as productive of disability. It is the intention to recognize actually painful, unstable, or malaligned joints, due to healed injury, as entitled to at least the minimum compensable rating for the joint. Crepitation either in the soft tissues such as the tendons or ligaments, or crepitation within the joint structures should be noted carefully as points of contact which are diseased. Flexion elicits such manifestations. The joints involved should be tested for pain on both active and passive motion, in weight-bearing and nonweight-bearing and, if possible, with the range of the opposite undamaged joint.

§ 4.60 [Reserved]

§ 4.61 Examination.

With any form of arthritis (except traumatic arthritis) it is essential that the examination for rating purposes cover all major joints, with especial

Department of Veterans Affairs

§ 4.68

reference to Heberden's or Haygarth's nodes.

§ 4.62 Circulatory disturbances.

The circulatory disturbances, especially of the lower extremity following injury in the popliteal space, must not be overlooked, and require rating generally as phlebitis.

§ 4.63 Loss of use of hand or foot.

Loss of use of a hand or a foot, for the purpose of special monthly compensation, will be held to exist when no effective function remains other than that which would be equally well served by an amputation stump at the site of election below elbow or knee with use of a suitable prosthetic appliance. The determination will be made on the basis of the actual remaining function of the hand or foot, whether the acts of grasping, manipulation, etc., in the case of the hand, or of balance and propulsion, etc., in the case of the foot, could be accomplished equally well by an amputation stump with prosthesis.

(a) Extremely unfavorable complete ankylosis of the knee, or complete ankylosis of 2 major joints of an extremity, or shortening of the lower extremity of 3½ inches (8.9 cms.) or more, will be taken as loss of use of the hand or foot involved.

(b) Complete paralysis of the external popliteal nerve (common peroneal) and consequent, footdrop, accompanied by characteristic organic changes including trophic and circulatory disturbances and other concomitants confirmatory of complete paralysis of this nerve, will be taken as loss of use of the foot.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

§ 4.64 Loss of use of both buttocks.

Loss of use of both buttocks shall be deemed to exist when there is severe damage to muscle Group XVII, bilateral (diagnostic code number 5317) and additional disability rendering it impossible for the disabled person, without assistance, to rise from a seated position and from a stooped position (fingers to toes position) and to maintain postural stability (the pelvis upon head of femur). The assistance may be

rendered by the person's own hands or arms, and, in the matter of postural stability, by a special appliance.

§ 4.65 [Reserved]

§ 4.66 Sacroiliac joint.

The common cause of disability in this region is arthritis, to be identified in the usual manner. The lumbosacral and sacroiliac joints should be considered as one anatomical segment for rating purposes. X-ray changes from arthritis in this location are decrease or obliteration of the joint space, with the appearance of increased bone density of the sacrum and ilium and sharpening of the margins of the joint. Disability is manifest from erector spinae spasm (not accounted for by other pathology), tenderness on deep palpation and percussion over these joints, loss of normal quickness of motion and resiliency, and postural defects often accompanied by limitation of flexion and extension of the hip. Traumatism is a rare cause of disability in this connection, except when superimposed upon congenital defect or upon an existent arthritis; to permit assumption of pure traumatic origin, objective evidence of damage to the joint, and history of trauma sufficiently severe to injure this extremely strong and practically immovable joint is required. There should be careful consideration of lumbosacral sprain, and the various symptoms of pain and paralysis attributable to disease affecting the lumbar vertebrae and the intervertebral disc.

§ 4.67 Pelvic bones.

The variability of residuals following these fractures necessitates rating on specific residuals, faulty posture, limitation of motion, muscle injury, painful motion of the lumbar spine, manifest by muscle spasm, mild to moderate sciatic neuritis, peripheral nerve injury, or limitation of hip motion.

§ 4.68 Amputation rule.

The combined rating for disabilities of an extremity shall not exceed the rating for the amputation at the elective level, were amputation to be performed. For example, the combined evaluations for disabilities below the knee shall not exceed the 40 percent

§ 4.69

evaluation, diagnostic code 5165. This 40 percent rating may be further combined with evaluation for disabilities above the knee but not to exceed the above the knee amputation elective level. Painful neuroma of a stump after amputation shall be assigned the evaluation for the elective site of re-amputation.

§ 4.69 Dominant hand.

Handedness for the purpose of a dominant rating will be determined by the evidence of record, or by testing on VA examination. Only one hand shall be considered dominant. The injured hand, or the most severely injured hand, of an ambidextrous individual will be considered the dominant hand for rating purposes.

(Authority: 38 U.S.C. 1155)

[62 FR 30239, June 3, 1997]

§ 4.70 Inadequate examinations.

If the report of examination is inadequate as a basis for the required consideration of service connection and evaluation, the rating agency may request a supplementary report from the examiner giving further details as to the limitations of the disabled person's ordinary activity imposed by the dis-

38 CFR Ch. I (7-1-01 Edition)

ease, injury, or residual condition, the prognosis for return to, or continuance of, useful work. When the best interests of the service will be advanced by personal conference with the examiner, such conference may be arranged through channels.

§ 4.71 Measurement of ankylosis and joint motion.

Plates I and II provide a standardized description of ankylosis and joint motion measurement. The anatomical position is considered as 0°, with two major exceptions: (a) Shoulder rotation—arm abducted to 90°, elbow flexed to 90° with the position of the forearm reflecting the midpoint 0° between internal and external rotation of the shoulder; and (b) supination and pronation—the arm next to the body, elbow flexed to 90°, and the forearm in midposition 0° between supination and pronation. Motion of the thumb and fingers should be described by appropriate reference to the joints (See Plate III) whose movement is limited, with a statement as to how near, in centimeters, the tip of the thumb can approximate the fingers, or how near the tips of the fingers can approximate the median transverse fold of the palm.




PLATE I




PLATE II

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

§4.71a Schedule of ratings—musculo-skeletal system.

ACUTE, SUBACUTE, OR CHRONIC DISEASES—Continued

| ACUTE, SUBACUTE, OR CHRONIC DISEASES | | Rating |
|---|-----|---|
| 5000 Osteomyelitis, acute, subacute, or chronic: | | |
| Of the pelvis, vertebrae, or extending into major joints, or with multiple localization or with long history of intractability and debility, anemia, amyloid liver changes, or other continuous constitutional symptoms | 100 | NOTE (1): A rating of 10 percent, as an exception to the amputation rule, is to be assigned in any case of active osteomyelitis where the amputation rating for the affected part is no percent. This 10 percent rating and the other partial ratings of 30 percent or less are to be combined with ratings for ankylosis, limited motion, nonunion or malunion, shortening, etc., subject, of course, to the amputation rule. The 60 percent rating, as it is based on constitutional symptoms, is not subject to the amputation rule. A rating for osteomyelitis will not be applied following cure by removal or radical resection of the affected bone. |
| Frequent episodes, with constitutional symptoms | 60 | |
| With definite involucrum or sequestrum, with or without discharging sinus | 30 | |
| With discharging sinus or other evidence of active infection within the past 5 years | 20 | |
| Inactive, following repeated episodes, without evidence of active infection in past 5 years | 10 | |

Department of Veterans Affairs

§ 4.71a

ACUTE, SUBACUTE, OR CHRONIC DISEASES—
Continued

ACUTE, SUBACUTE, OR CHRONIC DISEASES—
Continued

| | Rat- ing |
|--|------------------------------------|
| NOTE (2): The 20 percent rating on the basis of activity within the past 5 years is not assignable following the initial infection of active osteomyelitis with no subsequent reactivation. The prerequisite for this historical rating is an established recurrent osteomyelitis. To qualify for the 10 percent rating, 2 or more episodes following the initial infection are required. This 20 percent rating or the 10 percent rating, when applicable, will be assigned once only to cover disability at all sites of previously active infection with a future ending date in the case of the 20 percent rating. | |
| 5001 Bones and joints, tuberculosis of, active or inactive: Active Inactive: See §§ 4.88b and 4.89. | 100 |
| 5002 Arthritis rheumatoid (atrophic) <i>As an active process:</i> With constitutional manifestations associated with active joint involvement, totally incapacitating Less than criteria for 100% but with weight loss and anemia productive of severe impairment of health or severely incapacitating exacerbations occurring 4 or more times a year or a lesser number over prolonged periods Symptom combinations productive of definite impairment of health objectively supported by examination findings or incapacitating exacerbations occurring 3 or more times a year One or two exacerbations a year in a well-established diagnosis | 100 100 60 40 |
| For chronic residuals: For residuals such as limitation of motion or ankylosis, favorable or unfavorable, rate under the appropriate diagnostic codes for the specific joints involved. Where, however, the limitation of motion of the specific joint or joints involved is noncompensable under the codes a rating of 10 percent is for application for each such major joint or group of minor joints affected by limitation of motion, to be combined, not added under diagnostic code 5002. Limitation of motion must be objectively confirmed by findings such as swelling, muscle spasm, or satisfactory evidence of painful motion. NOTE: The ratings for the active process will not be combined with the residual ratings for limitation of motion or ankylosis. Assign the higher evaluation. | 20 |
| 5003 Arthritis, degenerative (hypertrophic or osteoarthritis): Degenerative arthritis established by X-ray findings will be rated on the basis of limitation of motion under the appropriate diagnostic codes for the specific joint or joints involved (DC 5200 etc.). When however, the limitation of motion of the specific joint or joints involved is noncompensable under the appropriate diagnostic codes, a rating of 10 pct is for application for each such major joint or group of minor joints affected by limitation of motion, to be combined, not added under diagnostic code 5003. Limitation of motion must be objectively confirmed by findings such as swelling, muscle spasm, or satisfactory evidence of painful motion. In the absence of limitation of motion, rate as below: | |

| | Rat- ing |
|---|-------------|
| With X-ray evidence of involvement of 2 or more major joints or 2 or more minor joint groups, with occasional incapacitating exacerbations | 20 |
| With X-ray evidence of involvement of 2 or more major joints or 2 or more minor joint groups | 10 |
| NOTE (1): The 20 pct and 10 pct ratings based on X-ray findings, above, will not be combined with ratings based on limitation of motion. NOTE (2): The 20 pct and 10 pct ratings based on X-ray findings, above, will not be utilized in rating conditions listed under diagnostic codes 5013 to 5024, inclusive. | |
| 5004 Arthritis, gonorrhoeal. | |
| 5005 Arthritis, pneumococcic. | |
| 5006 Arthritis, typhoid. | |
| 5007 Arthritis, syphilitic. | |
| 5008 Arthritis, streptococcic. | |
| 5009 Arthritis, other types (specify). With the types of arthritis, diagnostic codes 5004 through 5009, rate the disability as rheumatoid arthritis. | |
| 5010 Arthritis, due to trauma, substantiated by X-ray findings: Rate as arthritis, degenerative. | |
| 5011 Bones, caisson disease of: Rate as arthritis, cord involvement, or deafness, depending on the severity of disabling manifestations. | |
| 5012 Bones, new growths of, malignant NOTE: The 100 percent rating will be continued for 1 year following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. At this point, if there has been no local recurrence or metastases, the rating will be made on residuals. | 100 |
| 5013 Osteoporosis, with joint manifestations. | |
| 5014 Osteomalacia. | |
| 5015 Bones, new growths of, benign. | |
| 5016 Osteitis deformans. | |
| 5017 Gout. | |
| 5018 Hydrarthrosis, intermittent. | |
| 5019 Bursitis. | |
| 5020 Synovitis. | |
| 5021 Myositis. | |
| 5022 Periostitis. | |
| 5023 Myositis ossificans. | |
| 5024 Tenosynovitis. The diseases under diagnostic codes 5013 through 5024 will be rated on limitation of motion of affected parts, as arthritis, degenerative, except gout which will be rated under diagnostic code 5002. | |
| 5025 Fibromyalgia (fibrositis, primary fibromyalgia syndrome) With widespread musculoskeletal pain and tender points, with or without associated fatigue, sleep disturbance, stiffness, paresthesias, headache, irritable bowel symptoms, depression, anxiety, or Raynaud's-like symptoms: That are constant, or nearly so, and refractory to therapy | 40 |
| That are episodic, with exacerbations often precipitated by environmental or emotional stress or by overexertion, but that are present more than one-third of the time | 20 |
| That require continuous medication for control | 10 |

§4.71a

38 CFR Ch. I (7-1-01 Edition)

ACUTE, SUBACUTE, OR CHRONIC DISEASES—
Continued

| | Rating |
|--|--------|
| NOTE: Widespread pain means pain in both the left and right sides of the body, that is both above and below the waist, and that affects both the axial skeleton (i.e., cervical spine, anterior chest, thoracic spine, or low back) and the extremities. | |

PROSTHETIC IMPLANTS

| | Rating | |
|--|--------|-------|
| | Major | Minor |
| 5051 Shoulder replacement (prosthesis). Prosthetic replacement of the shoulder joint: For 1 year following implantation of prosthesis With chronic residuals consisting of severe, painful motion or weakness in the affected extremity With intermediate degrees of residual weakness, pain or limitation of motion, rate by analogy to diagnostic codes 5200 and 5203. Minimum rating | 100 | 100 |
| 5052 Elbow replacement (prosthesis). Prosthetic replacement of the elbow joint: For 1 year following implantation of prosthesis With chronic residuals consisting of severe painful motion or weakness in the affected extremity With intermediate degrees of residual weakness, pain or limitation of motion rate by analogy to diagnostic codes 5205 through 5208. Minimum evaluation | 100 | 100 |
| 5053 Wrist replacement (prosthesis). Prosthetic replacement of wrist joint: For 1 year following implantation of prosthesis With chronic residuals consisting of severe, painful motion or weakness in the affected extremity With intermediate degrees of residual weakness, pain or limitation of motion, rate by analogy to diagnostic code 5214. Minimum rating | 100 | 100 |
| 5054 Hip replacement (prosthesis). Prosthetic replacement of the head of the femur or of the acetabulum: For 1 year following implantation of prosthesis | 100 | 100 |

PROSTHETIC IMPLANTS—Continued

| | Rating | |
|--|--------|-------|
| | Major | Minor |
| Following implantation of prosthesis with painful motion or weakness such as to require the use of crutches | | 90 |
| Markedly severe residual weakness, pain or limitation of motion following implantation of prosthesis | | 70 |
| Moderately severe residuals of weakness, pain or limitation of motion | | 50 |
| Minimum rating | | 30 |
| 5055 Knee replacement (prosthesis). Prosthetic replacement of knee joint: For 1 year following implantation of prosthesis With chronic residuals consisting of severe painful motion or weakness in the affected extremity With intermediate degrees of residual weakness, pain or limitation of motion rate by analogy to diagnostic codes 5256, 5261, or 5262. Minimum rating | | 100 |
| 5056 Ankle replacement (prosthesis). Prosthetic replacement of ankle joint: For 1 year following implantation of prosthesis With chronic residuals consisting of severe painful motion or weakness With intermediate degrees of residual weakness, pain or limitation of motion rate by analogy to 5270 or 5271. Minimum rating | | 100 |
| NOTE (1): The 100 pct rating for 1 year following implantation of prosthesis will commence after initial grant of the 1-month total rating assigned under §4.30 following hospital discharge. NOTE (2): Special monthly compensation is assignable during the 100 pct rating period the earliest date permanent use of crutches is established. | | 40 |
| | | 20 |
| COMBINATIONS OF DISABILITIES | | |
| 5104 Anatomical loss of one hand and loss of use of one foot | | 100 |
| 5105 Anatomical loss of one foot and loss of use of one hand | | 100 |
| 5106 Anatomical loss of both hands | | 100 |
| 5107 Anatomical loss of both feet | | 100 |
| 5108 Anatomical loss of one hand and one foot | | 100 |
| 5109 Loss of use of both hands | | 100 |
| 5110 Loss of use of both feet | | 100 |
| 5111 Loss of use of one hand and one foot | | 100 |

¹ Also entitled to special monthly compensation.

TABLE II—RATINGS FOR MULTIPLE LOSSES OF EXTREMITIES WITH DICTATOR’S RATING CODE AND 38 CFR CITATION

| Impairment of one extremity | Impairment of other extremity | | | | | |
|--|---|---|---|--|--|---|
| | Anatomical loss or loss of use below elbow | Anatomical loss or loss of use below knee | Anatomical loss or loss of use above elbow (preventing use of prosthesis) | Anatomical loss or loss of use above knee (preventing use of prosthesis) | Anatomical loss near shoulder (preventing use of prosthesis) | Anatomical loss near hip (preventing use of prosthesis) |
| Anatomical loss or loss of use below elbow. | M Codes M-1 a, b, or c, 38 CFR 3.350 (c)(1)(i). | L Codes L-1 d, e, f, or g, 38 CFR 3.350(b). | M½ Code M-5, 38 CFR 3.350 (f)(1)(x). | L½ Code L-2 c, 38 CFR 3.350 (f)(1)(vi). | N Code N-3, 38 CFR 3.350 (f)(1)(xi). | M Code M-3 c, 38 CFR 3.350 (f)(1)(viii) |
| Anatomical loss or loss of use below knee. | | L Codes L-1 a, b, or c, 38 CFR 3.350(b). | L½ Code L-2 b, 38 CFR 3.350 (f)(1)(iii). | L½ Code L-2 a, 38 CFR 3.350 (f)(1)(i). | M Code M-3 b, 38 CFR 3.350 (f)(1)(iv). | M Code M-3 a, 38 CFR 3.350 (f)(1)(ii) |
| Anatomical loss or loss of use above elbow (preventing use of prosthesis). | | | N Code N-1, 38 CFR 3.350 (d)(1). | M Code M-2 a, 38 CFR 3.350 (c)(1)(iii). | N½ Code N-4, 38 CFR 3.350 (f)(1)(ix). | M½ Code M-4 c, 38 CFR 3.350 (f)(1)(xi) |
| Anatomical loss or loss of use above knee (preventing use of prosthesis). | | | | M Code M-2 a, 38 CFR 3.350 (c)(1)(ii). | M½ Code M-4 b, 38 CFR 3.350 (f)(1)(vii). | M½ Code M-4 a, 38 CFR 3.350 (f)(1)(v) |
| Anatomical loss near shoulder (preventing use of prosthesis). | | | | | O Code O-1, 38 CFR 3.350 (e)(1)(i). | N Code N-2 b, 38 CFR 3.350 (d)(3) |
| Anatomical loss near hip (preventing use of prosthesis). | | | | | | N Code N-2 a, 38 CFR 3.350 (d)(2) |

NOTE.—Need for aid attendance or permanently bedridden qualifies for subpar. L. Code L-1 h, i (38 CFR 3.350(b)). Paraplegia with loss of use of both lower extremities and loss of anal and bladder sphincter control qualifies for subpar. O. Code O-2 (38 CFR 3.350(e)(2)). Where there are additional disabilities rated 50% or 100%, or anatomical or loss of use of a third extremity see 38 CFR 3.350(f) (3), (4) or (5).

(Authority: 38 U.S.C. 1115)

§4.71a

38 CFR Ch. I (7-1-01 Edition)

AMPUTATIONS: UPPER EXTREMITY

| | Rating | |
|---|--------|-------|
| | Major | Minor |
| Arm, amputation of: | | |
| 5120 Disarticulation | 190 | 190 |
| 5121 Above insertion of deltoid | 190 | 180 |
| 5122 Below insertion of deltoid | 180 | 170 |
| Forearm, amputation of: | | |
| 5123 Above insertion of pronator teres | 180 | 170 |
| 5124 Below insertion of pronator teres | 170 | 160 |
| 5125 Hand, loss of use of | 170 | 160 |
| MULTIPLE FINGER AMPUTATIONS | | |
| 5126 Five digits of one hand, amputation of | 170 | 160 |
| Four digits of one hand, amputation of: | | |
| 5127 Thumb, index, middle and ring | 170 | 160 |
| 5128 Thumb, index, middle and little | 170 | 160 |
| 5129 Thumb, index, ring and little | 170 | 160 |
| 5130 Thumb, middle, ring and little | 170 | 160 |
| 5131 Index, middle, ring and little | 60 | 50 |
| Three digits of one hand, amputation of: | | |
| 5132 Thumb, index and middle | 60 | 50 |
| 5133 Thumb, index and ring | 60 | 50 |
| 5134 Thumb, index and little | 60 | 50 |
| 5135 Thumb, middle and ring | 60 | 50 |
| 5136 Thumb, middle and little | 60 | 50 |
| 5137 Thumb, ring and little | 60 | 50 |
| 5138 Index, middle and ring | 50 | 40 |
| 5139 Index, middle and little | 50 | 40 |
| 5140 Index, ring and little | 50 | 40 |
| 5141 Middle, ring and little | 40 | 30 |
| Two digits of one hand, amputation of: | | |
| 5142 Thumb and index | 50 | 40 |
| 5143 Thumb and middle | 50 | 40 |
| 5144 Thumb and ring | 50 | 40 |
| 5145 Thumb and little | 50 | 40 |
| 5146 Index and middle | 40 | 30 |
| 5147 Index and ring | 40 | 30 |
| 5148 Index and little | 40 | 30 |
| 5149 Middle and ring | 30 | 20 |
| 5150 Middle and little | 30 | 20 |
| 5151 Ring and little | 30 | 20 |
| (a) The ratings for multiple finger amputations apply to amputations at the proximal interphalangeal joints or through proximal phalanges.. | | |
| (b) Amputation through middle phalanges will be rated as prescribed for unfavorable ankylosis of the fingers.. | | |
| (c) Amputations at distal joints, or through distal phalanges, other than negligible losses, will be rated as prescribed for favorable ankylosis of the fingers.. | | |

AMPUTATIONS: UPPER EXTREMITY—Continued

| | Rating | |
|---|--------|-------|
| | Major | Minor |
| (d) Amputation or resection of metacarpal bones (more than one-half the bone lost) in multiple fingers injuries will require a rating of 10 percent added to (not combined with) the ratings, multiple finger amputations, subject to the amputation rule applied to the forearm. | | |
| (e) Combinations of finger amputations at various levels, or finger amputations with ankylosis or limitation of motion of the fingers will be rated on the basis of the grade of disability; i.e., amputation, unfavorable ankylosis, most representative of the levels or combinations. With an even number of fingers involved, and adjacent grades of disability, select the higher of the two grades. | | |
| (f) Loss of use of the hand will be held to exist when no effective function remains other than that which would be equally well served by an amputation stump with a suitable prosthetic appliance. | | |
| SINGLE FINGER AMPUTATIONS | | |
| 5152 Thumb, amputation of: | | |
| With metacarpal resection | 40 | 30 |
| At metacarpophalangeal joint or through proximal phalanx | 30 | 20 |
| At distal joint or through distal phalanx | 20 | 20 |
| 5153 Index finger, amputation of: | | |
| With metacarpal resection (more than one-half the bone lost) | 30 | 20 |
| Without metacarpal resection, at proximal interphalangeal joint or proximal thereto | 20 | 20 |
| Through middle phalanx or at distal joint | 10 | 10 |
| 5154 Middle finger, amputation of: | | |
| With metacarpal resection (more than one-half the bone lost) | 20 | 20 |
| Without metacarpal resection, at proximal interphalangeal joint or proximal thereto | 10 | 10 |
| 5155 Ring finger, amputation of: | | |
| With metacarpal resection (more than one-half the bone lost) | 20 | 20 |
| Without metacarpal resection, at proximal interphalangeal joint or proximal thereto | 10 | 10 |
| 5156 Little finger, amputation of: | | |
| With metacarpal resection (more than one-half the bone lost) | 20 | 20 |
| Without metacarpal resection, at proximal interphalangeal joint or proximal thereto | 10 | 10 |
| NOTE: The single finger amputation ratings are the only applicable ratings for amputations of whole or part of single fingers. | | |

¹ Entitled to special monthly compensation.

SINGLE FINGER AMPUTATIONS




PLATE III

§4.71a

38 CFR Ch. I (7-1-01 Edition)

AMPUTATIONS: LOWER EXTREMITY

AMPUTATIONS: LOWER EXTREMITY—Continued

| | Rat- ing |
|---|-------------|
| Thigh, amputation of: | |
| 5160 Disarticulation, with loss of extrinsic pelvic gir- dle muscles | 2 90 |
| 5161 Upper third, one-third of the distance from perineum to knee joint measured from perineum ... | 2 80 |
| 5162 Middle or lower thirds | 2 60 |
| Leg, amputation of: | |
| 5163 With defective stump, thigh amputation rec- ommended | 2 60 |
| 5164 Amputation not improvable by prosthesis con- trolled by natural knee action | 2 60 |
| 5165 At a lower level, permitting prosthesis | 2 40 |
| 5166 Forefoot, amputation proximal to metatarsal bones (more than one-half of metatarsal loss) | 2 40 |
| 5167 Foot, loss of use of | 2 40 |

| | Rat- ing |
|---|-------------|
| 5170 Toes, all, amputation of, without metatarsal loss | 30 |
| 5171 Toe, great, amputation of: | |
| With removal of metatarsal head | 30 |
| Without metatarsal involvement | 10 |
| 5172 Toes, other than great, amputation of, with re- moval of metatarsal head: | |
| One or two | 20 |
| Without metatarsal involvement | 0 |
| 5173 Toes, three or four, amputation of, without metatarsal involvement: | |
| Including great toe | 20 |
| Not including great toe | 10 |

² Also entitled to special monthly compensation.

AMPUTATIONS: LOWER EXTREMITY




PLATE IV

§4.71a

38 CFR Ch. I (7-1-01 Edition)

THE SHOULDER AND ARM

| | Rating | |
|---|--------|-------|
| | Major | Minor |
| 5200 Scapulohumeral articulation, ankylosis of: | | |
| NOTE: The scapula and humerus move as one piece. | | |
| Unfavorable, abduction limited to 25° from side | 50 | 40 |
| Intermediate between favorable and unfavorable | 40 | 30 |
| Favorable, abduction to 60°, can reach mouth and head | 30 | 20 |
| 5201 Arm, limitation of motion of: | | |
| To 25° from side | 40 | 30 |
| Midway between side and shoulder level | 30 | 20 |
| At shoulder level | 20 | 20 |
| 5202 Humerus, other impairment of: | | |
| Loss of head of (flail shoulder) | 80 | 70 |
| Nonunion of (false flail joint) | 60 | 50 |
| Fibrous union of | 50 | 40 |
| Recurrent dislocation of at scapulohumeral joint. | | |
| With frequent episodes and guarding of all arm movements | 30 | 20 |
| With infrequent episodes, and guarding of movement only at shoulder level | 20 | 20 |
| Malunion of: | | |
| Marked deformity | 30 | 20 |
| Moderate deformity | 20 | 20 |
| 5203 Clavicle or scapula, impairment of: | | |
| Dislocation of | 20 | 20 |
| Nonunion of: | | |
| With loose movement | 20 | 20 |
| Without loose movement | 10 | 10 |
| Malunion of | 10 | 10 |
| Or rate on impairment of function of contiguous joint. | | |

THE ELBOW AND FOREARM

| | Rating | |
|--|--------|-------|
| | Major | Minor |
| 5205 Elbow, ankylosis of: | | |
| Unfavorable, at an angle of less than 50° or with complete loss of supination or pronation | 60 | 50 |
| Intermediate, at an angle of more than 90°, or between 70° and 50° | 50 | 40 |
| Favorable, at an angle between 90° and 70° | 40 | 30 |
| 5206 Forearm, limitation of flexion of: | | |
| Flexion limited to 45° | 50 | 40 |
| Flexion limited to 55° | 40 | 30 |
| Flexion limited to 70° | 30 | 20 |
| Flexion limited to 90° | 20 | 20 |
| Flexion limited to 100° | 10 | 10 |
| Flexion limited to 110° | 0 | 0 |
| 5207 Forearm, limitation of extension of: | | |
| Extension limited to 110° | 50 | 40 |
| Extension limited to 100° | 40 | 30 |
| Extension limited to 90° | 30 | 20 |
| Extension limited to 75° | 20 | 20 |
| Extension limited to 60° | 10 | 10 |
| Extension limited to 45° | 10 | 10 |
| 5208 Forearm, flexion limited to 100° and extension to 45° | 20 | 20 |
| 5209 Elbow, other impairment of Flail joint | 60 | 50 |

THE ELBOW AND FOREARM—Continued

| | Rating | |
|--|--------|-------|
| | Major | Minor |
| Joint fracture, with marked cubitus varus or cubitus valgus deformity or with ununited fracture of head of radius | 20 | 20 |
| 5210 Radius and ulna, nonunion of, with flail false joint | 50 | 40 |
| 5211 Ulna, impairment of: | | |
| Nonunion in upper half, with false movement: | | |
| With loss of bone substance (1 inch (2.5 cms.) or more) and marked deformity | 40 | 30 |
| Without loss of bone substance or deformity | 30 | 20 |
| Nonunion in lower half | 20 | 20 |
| Malunion of, with bad alignment | 10 | 10 |
| 5212 Radius, impairment of: | | |
| Nonunion in lower half, with false movement: | | |
| With loss of bone substance (1 inch (2.5 cms.) or more) and marked deformity | 40 | 30 |
| Without loss of bone substance or deformity | 30 | 20 |
| Nonunion in upper half | 20 | 20 |
| Malunion of, with bad alignment | 10 | 10 |
| 5213 Supination and pronation, impairment of: | | |
| Loss of (bone fusion): | | |
| The hand fixed in supination or hyperpronation | 40 | 30 |
| The hand fixed in full pronation | 30 | 20 |
| The hand fixed near the middle of the arc or moderate pronation | 20 | 20 |
| Limitation of pronation: | | |
| Motion lost beyond middle of arc ... | 30 | 20 |
| Motion lost beyond last quarter of arc, the hand does not approach full pronation | 20 | 20 |
| Limitation of supination: | | |
| To 30° or less | 10 | 10 |
| NOTE: In all the forearm and wrist injuries, codes 5205 through 5213, multiple impaired finger movements due to tendon tie-up, muscle or nerve injury, are to be separately rated and combined not to exceed rating for loss of use of hand. | | |

THE WRIST

| | Rating | |
|---|--------|-------|
| | Major | Minor |
| 5214 Wrist, ankylosis of: | | |
| Unfavorable, in any degree of palmar flexion, or with ulnar or radial deviation | 50 | 40 |
| Any other position, except favorable | 40 | 30 |
| Favorable in 20° to 30° dorsiflexion | 30 | 20 |
| NOTE: Extremely unfavorable ankylosis will be rated as loss of use of hands under diagnostic code 5125. | | |
| 5215 Wrist, limitation of motion of: | | |
| Dorsiflexion less than 15° | 10 | 10 |
| Palmar flexion limited in line with forearm | 10 | 10 |

Department of Veterans Affairs

§4.71a

MULTIPLE FINGERS: UNFAVORABLE ANKYLOSIS

| | Rating | |
|--|--------|-------|
| | Major | Minor |
| In classifying the severity of ankylosis and limitation of motion of single digits and combinations of digits the following rules will be observed: | | |
| (1) Ankylosis of both the metacarpophalangeal and proximal interphalangeal joints, with either joint in extension or in extreme flexion, will be rated as amputation. | | |
| (2) Ankylosis of both the metacarpophalangeal and proximal interphalangeal joints, even though each is individually in favorable position, will be rated as unfavorable ankylosis. | | |
| (3) With only one joint of a digit ankylosed or limited in its motion, the determination will be made on the basis of whether motion is possible to within 2 inches (5.1 cms.) of the median transverse fold of the palm; when so possible, the rating will be for favorable ankylosis, otherwise unfavorable. | | |
| (4) With the thumb, the carpometacarpal joint is to be regarded as comparable to the metacarpophalangeal joint of other digits. | | |
| 5216 Five digits of one hand, unfavorable ankylosis of | 60 | 50 |
| 5217 Four digits of one hand, unfavorable ankylosis of: | | |
| Thumb, index, middle and ring | 60 | 50 |
| Thumb, index, middle and little | 60 | 50 |
| Thumb, index, ring and little | 60 | 50 |
| Thumb, middle, ring and little | 60 | 50 |
| Index, middle, ring and little | 50 | 40 |
| 5218 Three digits of one hand, unfavorable ankylosis of: | | |
| Thumb, index and middle | 50 | 40 |
| Thumb, index and ring | 50 | 40 |
| Thumb, index and little | 50 | 40 |
| Thumb, middle and ring | 50 | 40 |
| Thumb, middle and little | 50 | 40 |
| Thumb, ring and little | 50 | 40 |
| Index, middle and ring | 40 | 30 |
| Index, middle and little | 40 | 30 |
| Index, ring and little | 40 | 30 |
| Middle, ring and little | 30 | 20 |
| 5219 Two digits of one hand, unfavorable ankylosis of: | | |
| Thumb and index | 40 | 30 |
| Thumb and middle | 40 | 30 |
| Thumb and ring | 40 | 30 |
| Thumb and little | 40 | 30 |
| Index and middle | 30 | 20 |
| Index and ring | 30 | 20 |
| Index and little | 30 | 20 |
| Middle and ring | 20 | 20 |
| Middle and little | 20 | 20 |
| Ring and little | 20 | 20 |

MULTIPLE FINGERS: UNFAVORABLE ANKYLOSIS—
Continued

| | Rating | |
|--|--------|-------|
| | Major | Minor |
| (a) Extremely unfavorable ankylosis of the fingers, all joints in extension or in extreme flexion, or with rotation and angulation of bones, will be rated as amputation. | | |
| (b) The ratings for codes 5216 through 5219 apply to unfavorable ankylosis or limited motion preventing flexion of tips to within 2 inches (5.1 cms.) of median transverse fold of the palm. | | |
| (c) Combinations of finger amputations at various levels, or of finger amputations with ankylosis or limitation of motion of the fingers will be rated on the basis of the grade of disability, i.e., amputation, unfavorable ankylosis, or favorable ankylosis, most representative of the levels or combinations. With an even number of fingers involved, and adjacent grades of disability, select the higher of the two grades. | | |

MULTIPLE FINGERS: FAVORABLE ANKYLOSIS

| | Rating | |
|--|--------|-------|
| | Major | Minor |
| In classifying the severity of ankylosis and limitation of motion of single digits and combinations of digits the following rules will be observed: | | |
| (1) Ankylosis of both the metacarpophalangeal and proximal interphalangeal joints, with either joint in extension or in extreme flexion, will be rated as amputation. | | |
| (2) Ankylosis of both the metacarpophalangeal and proximal interphalangeal joints, even though each is individually in favorable position, will be rated as unfavorable ankylosis. | | |
| (3) With only one joint of a digit ankylosed or limited in its motion, the determination will be made on the basis of whether motion is possible to within 2 inches (5.1 cms.) of the median transverse fold of the palm; when so possible, the rating will be for favorable ankylosis, otherwise unfavorable. | | |
| (4) With the thumb, the carpometacarpal joint is to be regarded as comparable to the metacarpophalangeal joint of other digits. | | |
| 5220 Five digits of one hand, favorable ankylosis of | 50 | 40 |
| 5221 Four digits of one hand, favorable ankylosis of: | | |
| Thumb, index, middle and ring | 50 | 40 |
| Thumb, index, middle and little | 50 | 40 |
| Thumb, index, ring and little | 50 | 40 |
| Thumb, middle, ring and little | 50 | 40 |

§4.71a

38 CFR Ch. I (7-1-01 Edition)

MULTIPLE FINGERS: FAVORABLE ANKYLOSIS—
Continued

| | Rating | |
|---|--------|-------|
| | Major | Minor |
| Index, middle, ring and little | 40 | 30 |
| 5222 Three digits of one hand, favorable ankylosis of: | | |
| Thumb, index and middle | 40 | 30 |
| Thumb, index and ring | 40 | 30 |
| Thumb, index and little | 40 | 30 |
| Thumb, middle and ring | 40 | 30 |
| Thumb, middle and little | 40 | 30 |
| Thumb, ring and little | 40 | 30 |
| Index, middle and ring | 30 | 20 |
| Index, middle and little | 30 | 20 |
| Index, ring and little | 30 | 20 |
| Middle, ring and little | 20 | 20 |
| 5223 Two digits of one hand, favorable ankylosis of: | | |
| Thumb and index | 30 | 20 |
| Thumb and middle | 30 | 20 |
| Thumb and ring | 30 | 20 |
| Thumb and little | 30 | 20 |
| Index and middle | 20 | 20 |
| Index and ring | 20 | 20 |
| Index and little | 20 | 20 |
| Middle and ring | 10 | 10 |
| Middle and little | 10 | 10 |
| Ring and little | 10 | 10 |
| (a) The ratings for codes 5220 through 5223 apply to favorable ankylosis or limited motion permitting flexion of the tips to within 2 inches (5.1 cms.) of the transverse fold of the palm. Limitation of motion of less than 1 inch (2.5 cms.) in either direction is not considered disabling. | | |
| (b) Combination of finger amputations at various levels, or of finger amputations with ankylosis or limitation of motion of the fingers will be rated on the basis of the grade of disability, i.e., amputation, unfavorable ankylosis, or favorable ankylosis, most representative of the levels or combinations. With an even number of fingers involved, and adjacent grades of disability, select the higher of the two grades. | | |

ANKYLOSIS OF INDIVIDUAL FINGERS

| | Rating | |
|---|--------|-------|
| | Major | Minor |
| 5224 Thumb, ankylosis of: | | |
| Unfavorable | 20 | 20 |
| Favorable | 10 | 10 |
| 5225 Index finger, ankylosis of: | | |
| Unfavorable | 10 | 10 |
| Favorable | 10 | 10 |
| 5226 Middle finger, ankylosis of: | | |
| Unfavorable | 10 | 10 |
| Favorable | 10 | 10 |
| 5227 Finger, any other, ankylosis of | 0 | 0 |
| NOTE: Extremely unfavorable ankylosis will be rated as amputation under diagnostic codes 5152 through 5156. | | |

THE HIP AND THIGH

| | Rating |
|---|-----------------|
| 5250 Hip, ankylosis of: | |
| Unfavorable, extremely unfavorable ankylosis, the foot not reaching ground, crutches necessitated | ³ 90 |
| Intermediate | 70 |
| Favorable, in flexion at an angle between 20° and 40°, and slight adduction or abduction | 60 |
| 5251 Thigh, limitation of extension of: | |
| Extension limited to 5° | 10 |
| 5252 Thigh, limitation of flexion of: | |
| Flexion limited to 10° | 40 |
| Flexion limited to 20° | 30 |
| Flexion limited to 30° | 20 |
| Flexion limited to 45° | 10 |
| 5253 Thigh, impairment of: | |
| Limitation of abduction of, motion lost beyond 10° | 20 |
| Limitation of adduction of, cannot cross legs | 10 |
| Limitation of rotation of, cannot toe-out more than 15°, affected leg | 10 |
| 5254 Hip, flail joint | 80 |
| 5255 Femur, impairment of: | |
| Fracture of shaft or anatomical neck of: | |
| With nonunion, with loose motion (spiral or oblique fracture) | 80 |
| With nonunion, without loose motion, weightbearing preserved with aid of brace | 60 |
| Fracture of surgical neck of, with false joint | 60 |
| Malunion of: | |
| With marked knee or hip disability | 30 |
| With moderate knee or hip disability | 20 |
| With slight knee or hip disability | 10 |

³Entitled to special monthly compensation.

THE KNEE AND LEG

| | Rating |
|---|--------|
| 5256 Knee, ankylosis of: | |
| Extremely unfavorable, in flexion at an angle of 45° or more | 60 |
| In flexion between 20° and 45° | 50 |
| In flexion between 10° and 20° | 40 |
| Favorable angle in full extension, or in slight flexion between 0° and 10° | 30 |
| 5257 Knee, other impairment of: | |
| Recurrent subluxation or lateral instability: | |
| Severe | 30 |
| Moderate | 20 |
| Slight | 10 |
| 5258 Cartilage, semilunar, dislocated, with frequent episodes of "locking," pain, and effusion into the joint | 20 |
| 5259 Cartilage, semilunar, removal of, symptomatic | 10 |
| 5260 Leg, limitation of flexion of: | |
| Flexion limited to 15° | 30 |
| Flexion limited to 30° | 20 |
| Flexion limited to 45° | 10 |
| Flexion limited to 60° | 0 |
| 5261 Leg, limitation of extension of: | |
| Extension limited to 45° | 50 |
| Extension limited to 30° | 40 |
| Extension limited to 20° | 30 |
| Extension limited to 15° | 20 |
| Extension limited to 10° | 10 |
| Extension limited to 5° | 0 |
| 5262 Tibia and fibula, impairment of: | |
| Nonunion of, with loose motion, requiring brace | 40 |
| Malunion of: | |
| With marked knee or ankle disability | 30 |

Department of Veterans Affairs

§4.71a

THE KNEE AND LEG—Continued

| | Rat- ing |
|---|-------------|
| With moderate knee or ankle disability | 20 |
| With slight knee or ankle disability | 10 |
| 5263 Genu recurvatum (acquired, traumatic, with weakness and insecurity in weight-bearing objectively demonstrated) | 10 |

THE ANKLE

| | Rat- ing |
|--|-------------|
| 5270 Ankle, ankylosis of: | |
| In plantar flexion at more than 40°, or in dorsiflexion at more than 10° or with abduction, adduction, inversion or eversion deformity | 40 |
| In plantar flexion, between 30° and 40°, or in dorsiflexion, between 0° and 10° | 30 |
| In plantar flexion, less than 30° | 20 |
| 5271 Ankle, limited motion of: | |
| Marked | 20 |
| Moderate | 10 |
| 5272 Subastragalar or tarsal joint, ankylosis of: | |
| In poor weight-bearing position | 20 |
| In good weight-bearing position | 10 |
| 5273 Os calcis or astragalus, malunion of: | |
| Marked deformity | 20 |
| Moderate deformity | 10 |
| 5274 Astragalectomy | 20 |

SHORTENING OF THE LOWER EXTREMITY

| | Rat- ing |
|--|-----------------|
| 5275 Bones, of the lower extremity, shortening of: | |
| Over 4 inches (10.2 cms.) | ³ 60 |
| 3½ to 4 inches (8.9 cms. to 10.2 cms.) | ³ 50 |
| 3 to 3½ inches (7.6 cms. to 8.9 cms.) | 40 |
| 2½ to 3 inches (6.4 cms. to 7.6 cms.) | 30 |
| 2 to 2½ inches (5.1 cms. to 6.4 cms.) | 20 |
| 1¼ to 2 inches (3.2 cms. to 5.1 cms.) | 10 |
| NOTE: Measure both lower extremities from anterior superior spine of the ilium to the internal malleolus of the tibia. Not to be combined with other ratings for fracture or faulty union in the same extremity. | |

³ Also entitled to special monthly compensation.

THE FOOT

| | Rat- ing |
|--|-------------|
| 5276 Flatfoot, acquired: | |
| Pronounced; marked pronation, extreme tenderness of plantar surfaces of the feet, marked inward displacement and severe spasm of the tendo achillis on manipulation, not improved by orthopedic shoes or appliances. | |
| Bilateral | 50 |
| Unilateral | 30 |
| Severe; objective evidence of marked deformity (pronation, abduction, etc.), pain on manipulation and use accentuated, indication of swelling on use, characteristic callosities: | |
| Bilateral | 30 |
| Unilateral | 20 |

THE FOOT—Continued

| | Rat- ing |
|---|-------------|
| Moderate; weight-bearing line over or medial to great toe, inward bowing of the tendo achillis, pain on manipulation and use of the feet, bilateral or unilateral | 10 |
| Mild; symptoms relieved by built-up shoe or arch support | 0 |
| 5277 Weak foot, bilateral: | |
| A symptomatic condition secondary to many constitutional conditions, characterized by atrophy of the musculature, disturbed circulation, and weakness: | |
| Rate the underlying condition, minimum rating | 10 |
| 5278 Claw foot (pes cavus), acquired: | |
| Marked contraction of plantar fascia with dropped forefoot, all toes hammer toes, very painful callosities, marked varus deformity: | |
| Bilateral | 50 |
| Unilateral | 30 |
| All toes tending to dorsiflexion, limitation of dorsiflexion at ankle to right angle, shortened plantar fascia, and marked tenderness under metatarsal heads: | |
| Bilateral | 30 |
| Unilateral | 20 |
| Great toe dorsiflexed, some limitation of dorsiflexion at ankle, definite tenderness under metatarsal heads: | |
| Bilateral | 10 |
| Unilateral | 10 |
| Slight | 0 |
| 5279 Metatarsalgia, anterior (Morton's disease), unilateral, or bilateral | 10 |
| 5280 Hallux valgus, unilateral: | |
| Operated with resection of metatarsal head | 10 |
| Severe, if equivalent to amputation of great toe .. | 10 |
| 5281 Hallux rigidus, unilateral, severe: | |
| Rate as hallux valgus, severe. | |
| NOTE: Not to be combined with claw foot ratings. | |
| 5282 Hammer toe: | |
| All toes, unilateral without claw foot | 10 |
| Single toes | 0 |
| 5283 Tarsal, or metatarsal bones, malunion of, or nonunion of: | |
| Severe | 30 |
| Moderately severe | 20 |
| Moderate | 10 |
| NOTE: With actual loss of use of the foot, rate 40 percent. | |
| 5284 Foot injuries, other: | |
| Severe | 30 |
| Moderately severe | 20 |
| Moderate | 10 |
| NOTE: With actual loss of use of the foot, rate 40 percent. | |

THE SPINE

| | Rat- ing |
|---|-------------|
| 5285 Vertebra, fracture of, residuals: | |
| With cord involvement, bedridden, or requiring long leg braces | 100 |
| Consider special monthly compensation; with lesser involvements rate for limited motion, nerve paralysis. | |
| Without cord involvement; abnormal mobility requiring neck brace (jury mast) | 60 |

§ 4.72

38 CFR Ch. I (7-1-01 Edition)

THE SPINE—Continued

| | Rat- ing |
|---|-------------|
| In other cases rate in accordance with definite limited motion or muscle spasm, adding 10 percent for demonstrable deformity of vertebral body. NOTE: Both under ankylosis and limited motion, ratings should not be assigned for more than one segment by reason of involvement of only the first or last vertebrae of an adjacent segment. | |
| 5286 Spine, complete bony fixation (ankylosis) of: Unfavorable angle, with marked deformity and involvement of major joints (Marie-Strumpell type) or without other joint involvement (Bechterew type) | 100 |
| Favorable angle | 60 |
| 5287 Spine, ankylosis of, cervical: Unfavorable | 40 |
| Favorable | 30 |
| 5288 Spine, ankylosis of, dorsal: Unfavorable | 30 |
| Favorable | 20 |
| 5289 Spine, ankylosis of, lumbar: Unfavorable | 50 |
| Favorable | 40 |
| 5290 Spine, limitation of motion of, cervical: Severe | 30 |
| Moderate | 20 |
| Slight | 10 |
| 5291 Spine, limitation of motion of, dorsal: Severe | 10 |
| Moderate | 10 |
| Slight | 0 |
| 5292 Spine, limitation of motion of, lumbar: Severe | 40 |
| Moderate | 20 |
| Slight | 10 |
| 5293 Intervertebral disc syndrome: Pronounced; with persistent symptoms compatible with sciatic neuropathy with characteristic pain and demonstrable muscle spasm, absent ankle jerk, or other neurological findings appropriate to site of diseased disc, little intermittent relief | 60 |
| Severe; recurring attacks, with intermittent relief | 40 |
| Moderate; recurring attacks | 20 |
| Mild | 10 |
| Postoperative, cured | 0 |
| 5294 Sacro-iliac injury and weakness: | |
| 5295 Lumbosacral strain: Severe; with listing of whole spine to opposite side, positive Goldthwaite's sign, marked limitation of forward bending in standing position, loss of lateral motion with osteo-arthritic changes, or narrowing or irregularity of joint space, or some of the above with abnormal mobility on forced motion | 40 |
| With muscle spasm on extreme forward bending, loss of lateral spine motion, unilateral, in standing position | 20 |
| With characteristic pain on motion | 10 |
| With slight subjective symptoms only | 0 |

THE SKULL

| | Rat- ing |
|---|-------------|
| 5296 Skull, loss of part of, both inner and outer tables: | |

THE SKULL—Continued

| | Rat- ing |
|---|-------------|
| With brain hernia | 80 |
| Without brain hernia: Area larger than size of a 50-cent piece or 1.140 in ² (7.355 cm ²) | 50 |
| Area intermediate | 30 |
| Area smaller than the size of a 25-cent piece or 0.716 in ² (4.619 cm ²) | 10 |
| NOTE: Rate separately for intracranial complications. | |

THE RIBS

| | Rat- ing |
|--|-------------|
| 5297 Ribs, removal of: More than six | 50 |
| Five or six | 40 |
| Three or four | 30 |
| Two | 20 |
| One or resection of two or more ribs without regeneration | 10 |
| NOTE (1): The rating for rib resection or removal is not to be applied with ratings for purulent pleurisy, lobectomy, pneumonectomy or injuries of pleural cavity. | |
| NOTE (2): However, rib resection will be considered as rib removal in thoracoplasty performed for collapse therapy or to accomplish obliteration of space and will be combined with the rating for lung collapse, or with the rating for lobectomy, pneumonectomy or the graduated ratings for pulmonary tuberculosis. | |

THE COCCYX

| | Rat- ing |
|---|-------------|
| 5298 Coccyx, removal of: Partial or complete, with painful residuals | 10 |
| Without painful residuals | 0 |

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 40 FR 42536, Sept. 15, 1975; 41 FR 11294, Mar. 18, 1976; 43 FR 45350, Oct. 2, 1978; 51 FR 6411, Feb. 24, 1986; 61 FR 20439, May 7, 1996]

§ 4.72 [Reserved]

§ 4.73 Schedule of ratings—muscle injuries.

NOTE: When evaluating any claim involving muscle injuries resulting in loss of use of any extremity or loss of use of both buttocks (diagnostic code 5317, Muscle Group XVII), refer to § 3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation.

Department of Veterans Affairs

§ 4.73

THE SHOULDER GIRDLE AND ARM

THE FOREARM AND HAND

| | Rating | |
|---|----------|--------------|
| | Dominant | Non-dominant |
| 5301 Group I. <i>Function:</i> Upward rotation of scapula; elevation of arm above shoulder level. <i>Extrinsic muscles of shoulder girdle:</i> (1) Trapezius; (2) levator scapulae; (3) serratus magnus. | | |
| Severe | 40 | 30 |
| Moderately Severe | 30 | 20 |
| Moderate | 10 | 10 |
| Slight | 0 | 0 |
| 5302 Group II. <i>Function:</i> Depression of arm from vertical overhead to hanging at side (1, 2); downward rotation of scapula (3, 4); 1 and 2 act with Group III in forward and backward swing of arm. <i>Intrinsic muscles of shoulder girdle:</i> (1) Pectoralis major II (costosternal); (2) latissimus dorsi and teres major (teres major, although technically an intrinsic muscle, is included with latissimus dorsi); (3) pectoralis minor; (4) rhomboid. | | |
| Severe | 40 | 30 |
| Moderately Severe | 30 | 20 |
| Moderate | 20 | 20 |
| Slight | 0 | 0 |
| 5303 Group III. <i>Function:</i> Elevation and abduction of arm to level of shoulder; act with 1 and 2 of Group II in forward and backward swing of arm. <i>Intrinsic muscles of shoulder girdle:</i> (1) Pectoralis major I (clavicular); (2) deltoid. | | |
| Severe | 40 | 30 |
| Moderately Severe | 30 | 20 |
| Moderate | 20 | 20 |
| Slight | 0 | 0 |
| 5304 Group IV. <i>Function:</i> Stabilization of shoulder against injury in strong movements, holding head of humerus in socket; abduction; outward rotation and inward rotation of arm. <i>Intrinsic muscles of shoulder girdle:</i> (1) Supraspinatus; (2) infraspinatus and teres minor; (3) subscapularis; (4) coracobrachialis. | | |
| Severe | 30 | 20 |
| Moderately Severe | 20 | 20 |
| Moderate | 10 | 10 |
| Slight | 0 | 0 |
| 5305 Group V. <i>Function:</i> Elbow supination (1) (long head of biceps is stabilizer of shoulder joint); flexion of elbow (1, 2, 3). <i>Flexor muscles of elbow:</i> (1) Biceps; (2) brachialis; (3) brachioradialis. | | |
| Severe | 40 | 30 |
| Moderately Severe | 30 | 20 |
| Moderate | 10 | 10 |
| Slight | 0 | 0 |
| 5306 Group VI. <i>Function:</i> Extension of elbow (long head of triceps is stabilizer of shoulder joint). <i>Extensor muscles of the elbow:</i> (1) Triceps; (2) anconeus. | | |
| Severe | 40 | 30 |
| Moderately Severe | 30 | 20 |
| Moderate | 10 | 10 |
| Slight | 0 | 0 |

| | Rating | |
|---|----------|--------------|
| | Dominant | Non-dominant |
| 5307 Group VII. <i>Function:</i> Flexion of wrist and fingers. <i>Muscles arising from internal condyle of humerus:</i> Flexors of the carpus and long flexors of fingers and thumb; pronator. | | |
| Severe | 40 | 30 |
| Moderately Severe | 30 | 20 |
| Moderate | 10 | 10 |
| Slight | 0 | 0 |
| 5308 Group VIII. <i>Function:</i> Extension of wrist, fingers, and thumb; abduction of thumb. <i>Muscles arising mainly from external condyle of humerus:</i> Extensors of carpus, fingers, and thumb; supinator. | | |
| Severe | 30 | 20 |
| Moderately Severe | 20 | 20 |
| Moderate | 10 | 10 |
| Slight | 0 | 0 |
| 5309 Group IX. <i>Function:</i> The forearm muscles act in strong grasping movements and are supplemented by the intrinsic muscles in delicate manipulative movements. <i>Intrinsic muscles of hand:</i> Thenar eminence; short flexor, opponens, abductor and adductor of thumb; hypothenar eminence; short flexor, opponens and abductor of little finger; 4 lumbricales; 4 dorsal and 3 palmar interossei. | | |
| NOTE: The hand is so compact a structure that isolated muscle injuries are rare, being nearly always complicated with injuries of bones, joints, tendons, etc. Rate on limitation of motion, minimum 10 percent. | | |

THE FOOT AND LEG

| | Rating |
|---|--------|
| 5310 Group X. <i>Function:</i> Movements of forefoot and toes; propulsion thrust in walking. <i>Intrinsic muscles of the foot: Plantar:</i> (1) Flexor digitorum brevis; (2) abductor hallucis; (3) abductor digiti minimi; (4) quadratus plantae; (5) lumbricales; (6) flexor hallucis brevis; (7) adductor hallucis; (8) flexor digiti minimi brevis; (9) dorsal and plantar interossei. Other important plantar structures: Plantar aponeurosis, long plantar and calcaneonavicular ligament, tendons of posterior tibial, peroneus longus, and long flexors of great and little toes. | |
| Severe | 30 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |
| <i>Dorsal:</i> (1) Extensor hallucis brevis; (2) extensor digitorum brevis. Other important dorsal structures: cruciate, crural, deltoid, and other ligaments; tendons of long extensors of toes and peronei muscles. | |
| Severe | 20 |
| Moderately Severe | 10 |
| Moderate | 10 |
| Slight | 0 |

THE FOOT AND LEG—Continued

| | Rat- ing |
|--|-------------|
| NOTE: Minimum rating for through-and-through wounds of the foot—10. | |
| 5311 Group XI. <i>Function:</i> Propulsion, plantar flexion of foot (1); stabilization of arch (2, 3); flexion of toes (4, 5); Flexion of knee (6). <i>Posterior and lateral crural muscles, and muscles of the calf:</i> (1) Triceps surae (gastrocnemius and soleus); (2) tibialis posterior; (3) peroneus longus; (4) peroneus brevis; (5) flexor hallucis longus; (6) flexor digitorum longus; (7) popliteus; (8) plantaris. | |
| Severe | 30 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |
| 5312 Group XII. <i>Function:</i> Dorsiflexion (1); extension of toes (2); stabilization of arch (3). <i>Anterior muscles of the leg:</i> (1) Tibialis anterior; (2) extensor digitorum longus; (3) extensor hallucis longus; (4) peroneus tertius. | |
| Severe | 30 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |

THE PELVIC GIRDLE AND THIGH

| | Rat- ing |
|---|-------------|
| 5313 Group XIII. <i>Function:</i> Extension of hip and flexion of knee; outward and inward rotation of flexed knee; acting with rectus femoris and sartorius (see XIV, 1, 2) synchronizing simultaneous flexion of hip and knee and extension of hip and knee by belt-over-pulley action at knee joint. <i>Posterior thigh group, Hamstring complex of 2-joint muscles:</i> (1) Biceps femoris; (2) semimembranosus; (3) semitendinosus. | |
| Severe | 40 |
| Moderately Severe | 30 |
| Moderate | 10 |
| Slight | 0 |
| 5314 Group XIV. <i>Function:</i> Extension of knee (2, 3, 4, 5); simultaneous flexion of hip and flexion of knee (1); tension of fascia lata and iliotibial (Maissiat's) band, acting with XVII (1) in postural support of body (6); acting with hamstrings in synchronizing hip and knee (1, 2). <i>Anterior thigh group:</i> (1) Sartorius; (2) rectus femoris; (3) vastus externus; (4) vastus intermedius; (5) vastus internus; (6) tensor vaginae femoris. | |
| Severe | 40 |
| Moderately Severe | 30 |
| Moderate | 10 |
| Slight | 0 |
| 5315 Group XV. <i>Function:</i> Adduction of hip (1, 2, 3, 4); flexion of hip (1, 2); flexion of knee (4). <i>Mesial thigh group:</i> (1) Adductor longus; (2) adductor brevis; (3) adductor magnus; (4) gracilis. | |
| Severe | 30 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |
| 5316 Group XVI. <i>Function:</i> Flexion of hip (1, 2, 3). <i>Pelvic girdle group 1:</i> (1) Psoas; (2) iliacus; (3) pectineus. | |
| Severe | 40 |
| Moderately Severe | 30 |
| Moderate | 10 |
| Slight | 0 |

THE PELVIC GIRDLE AND THIGH—Continued

| | Rat- ing |
|---|-------------|
| 5317 Group XVII. <i>Function:</i> Extension of hip (1); abduction of thigh; elevation of opposite side of pelvis (2, 3); tension of fascia lata and iliotibial (Maissiat's) band, acting with XIV (6) in postural support of body steadying pelvis upon head of femur and condyles of femur on tibia (1). <i>Pelvic girdle group 2:</i> (1) Gluteus maximus; (2) gluteus medius; (3) gluteus minimus. | |
| Severe | *50 |
| Moderately Severe | 40 |
| Moderate | 20 |
| Slight | 0 |
| 5318 Group XVIII. <i>Function:</i> Outward rotation of thigh and stabilization of hip joint. <i>Pelvic girdle group 3:</i> (1) Piriformis; (2) gemellus (superior or inferior); (3) obturator (external or internal); (4) quadratus femoris. | |
| Severe | 30 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |

*If bilateral, see §3.350(a)(3) of this chapter to determine whether the veteran may be entitled to special monthly compensation.

THE TORSO AND NECK

| | Rat- ing |
|---|-------------|
| 5319 Group XIX. <i>Function:</i> Support and compression of abdominal wall and lower thorax; flexion and lateral motions of spine; synergists in strong downward movements of arm (1). <i>Muscles of the abdominal wall:</i> (1) Rectus abdominis; (2) external oblique; (3) internal oblique; (4) transversalis; (5) quadratus lumborum. | |
| Severe | 50 |
| Moderately Severe | 30 |
| Moderate | 10 |
| Slight | 0 |
| 5320 Group XX. <i>Function:</i> Postural support of body; extension and lateral movements of spine. <i>Spinal muscles:</i> Sacrospinalis (erector spinae and its prolongations in thoracic and cervical regions). <i>Cervical and thoracic region:</i> | |
| Severe | 40 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |
| <i>Lumbar region:</i> | |
| Severe | 60 |
| Moderately Severe | 40 |
| Moderate | 20 |
| Slight | 0 |
| 5321 Group XXI. <i>Function:</i> Respiration. <i>Muscles of respiration:</i> Thoracic muscle group. | |
| Severe or Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |
| 5322 Group XXII. <i>Function:</i> Rotary and forward movements of the head; respiration; deglutition. <i>Muscles of the front of the neck:</i> (Lateral, supra- and infrahyoid group.) (1) Trapezius I (clavicular insertion); (2) sternocleidomastoid; (3) the "hyoid" muscles; (4) sternothyroid; (5) digastric. | |
| Severe | 30 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |

Department of Veterans Affairs

§ 4.76a

THE TORSO AND NECK—Continued

| | Rat- ing |
|--|-------------|
| 5323 Group XXIII. <i>Function:</i> Movements of the head; fixation of shoulder movements. <i>Muscles of the side and back of the neck:</i> Suboccipital; lateral vertebral and anterior vertebral muscles. | |
| Severe | 30 |
| Moderately Severe | 20 |
| Moderate | 10 |
| Slight | 0 |

MISCELLANEOUS

| | Rat- ing |
|--|-------------|
| 5324 Diaphragm, rupture of, with herniation. Rate under diagnostic code 7346. | |
| 5325 Muscle injury, facial muscles. Evaluate functional impairment as seventh (facial) cranial nerve neuropathy (diagnostic code 8207), disfiguring scar (diagnostic code 7800), etc. Minimum, if interfering to any extent with mastication—10. | |
| 5326 Muscle hernia, extensive. Without other injury to the muscle—10. | |
| 5327 Muscle, neoplasm of, malignant (excluding soft tissue sarcoma)—100. | |
| NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residual impairment of function. | |
| 5328 Muscle, neoplasm of, benign, postoperative. Rate on impairment of function, i.e., limitation of motion, or scars, diagnostic code 7805, etc. | |
| 5329 Sarcoma, soft tissue (of muscle, fat, or fibrous connective tissue)—100. | |
| NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residual impairment of function. | |

(Authority: 38 U.S.C. 1155)
[62 FR 30239, June 3, 1997]

THE ORGANS OF SPECIAL SENSE

§ 4.75 Examination of visual acuity.

Ratings on account of visual impairments considered for service connection are, when practicable, to be based only on examination by specialists. Such special examinations should include uncorrected and corrected cen-

tral visual acuity for distance and near, with record of the refraction. Snellen's test type or its equivalent will be used. Mydriatics should be routine, except when contraindicated. Fundusoscopic and ophthalmological findings must be recorded. The best distant vision obtainable after best correction by glasses will be the basis of rating, except in cases of keratoconus in which contact lenses are medically required. Also, if there exists a difference of more than 4 diopters of spherical correction between the two eyes, the best possible visual acuity of the poorer eye without glasses, or with a lens of not more than 4 diopters difference from that used with the better eye will be taken as the visual acuity of the poorer eye. When such a difference exists, close attention will be given to the likelihood of congenital origin in mere refractive error.

[40 FR 42537, Sept. 15, 1975]

§ 4.76 Examination of field vision.

Measurement of the visual field will be made when there is disease of the optic nerve or when otherwise indicated. The usual perimetric methods will be employed, using a standard perimeter and 3 mm. white test object. At least 16 meridians 22½ degrees apart will be charted for each eye. (See Figure 1. For the 8 principal meridians, see table III.) The charts will be made a part of the report of examination. Not less than 2 recordings, and when possible, 3 will be made. The minimum limit for this function is established as a concentric central contraction of the visual field to 5°. This type of contraction of the visual field reduces the visual efficiency to zero. Where available the examination for form field should be supplemented, when indicated, by the use of tangent screen or campimeter. This last test is especially valuable in detection of scotoma.

[43 FR 45352, Oct. 2, 1978]

§ 4.76a Computation of average concentric contraction of visual fields.

The extent of contraction of visual field in each eye is determined by recording the extent of the remaining visual fields in each of the eight 45 degree principal meridians. The number

§4.76a

of degrees lost is determined at each meridian by subtracting the remaining degrees from the normal visual fields given in table III. The degrees lost are then added together to determine total degrees lost. This is subtracted from 500. The difference represents the total remaining degrees of visual field. The difference divided by eight represents the average contraction for rating purposes.

38 CFR Ch. I (7-1-01 Edition)

TABLE III—NORMAL VISUAL FIELD EXTENT AT 8 PRINCIPAL MERIDIANS

| Meridian | Normal de- grees |
|-----------------------|---------------------|
| Temporally | 85 |
| Down temporally | 85 |
| Down | 65 |
| Down nasally | 50 |
| Nasally | 60 |
| Up nasally | 55 |
| Up | 45 |
| Up temporally | 55 |
| Total | 500 |




Figure 1. Chart of visual field showing normal field right eye and abnormal contraction visual field left eye.

TS-19

52a

Example of computation of concentric contraction under the schedule with abnormal findings taken from Figure 1.

| Loss | Degrees |
|------------------------|---------|
| Temporarily | 55 |
| Down temporarily | 55 |
| Down | 45 |

§ 4.77

| Loss | Degrees |
|-------------------------|------------|
| Down nasally | 30 |
| Nasally | 40 |
| Up nasally | 35 |
| Up | 25 |
| Up temporally | 35 |
| Total loss | 320 |

Remaining field 500° minus 320°=180°. 180°÷8=22½° average concentric contraction.

[43 FR 45352, Oct. 2, 1978]

§ 4.77 Examination of muscle function.

The measurement of muscle function will be undertaken only when the history and findings reflect disease or injury of the extrinsic muscles of the eye, or of the motor nerves supplying these muscles. The measurement will

be performed using a Goldmann Perimeter Chart as in Figure 2 below. The chart identifies four major quadrants, (upward, downward, and two lateral) plus a central field (20 ° or less). The examiner will chart the areas in which diplopia exists, and such plotted chart will be made a part of the examination report. Muscle function is considered normal (20/40) when diplopia does not exist within 40° in the lateral or downward quadrants, or within 30° in the upward quadrant. Impairment of muscle function is to be supported in each instance by record of actual appropriate pathology. Diplopia which is only occasional or correctable is not considered a disability.




Figure 2. Goldmann Perimeter Chart

52c

Department of Veterans Affairs

§ 4.84a

[53 FR 30262, Aug. 11, 1988]

§ 4.78 Computing aggravation.

In determining the effect of aggravation of visual disability, even though the visual impairment of only one eye is service connected, evaluate the vision of both eyes, before and after suffering the aggravation, and subtract the former evaluation from the latter except when the bilateral vision amounts to total disability. In the event of subsequent increase in the disability of either eye, due to intercurrent disease or injury not associated with the service, the condition of the eyes before suffering the subsequent increase will be taken as the basis of compensation subject to the provisions of § 3.383(a) of this chapter.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45354, Oct. 2, 1978]

§ 4.79 Loss of use of one eye, having only light perception.

Loss of use or blindness of one eye, having only light perception, will be held to exist when there is inability to recognize test letters at 1 foot (.30m.) and when further examination of the eyes reveals that perception of objects, hand movements or counting fingers cannot be accomplished at 3 feet (.91m.), lesser extents of visions, particularly perception of objects, hand movements, or counting fingers at distances less than 3 feet (.91 m.), being considered of negligible utility. With visual acuity 5/200 (1.5/60) or less or the visual field reduced to 5° concentric contraction, in either event in both eyes, the question of entitlement on account of regular aid and attendance will be determined on the facts in the individual case.

[43 FR 45354, Oct. 2, 1978]

§ 4.80 Rating of one eye.

Combined ratings for disabilities of the same eye should not exceed the amount for total loss of vision of that eye unless there is an enucleation or a serious cosmetic defect added to the total loss of vision.

§§ 4.81–4.82 [Reserved]

§ 4.83 Ratings at scheduled steps and distances.

In applying the ratings for impairment of visual acuity, a person not having the ability to read at any one of the scheduled steps or distances, but reading at the next scheduled step or distance, is to be rated as reading at this latter step or distance. That is, a person who can read at 20/100 (6/30) but who cannot at 20/70 (6/21), should be rated as seeing at 20/100 (6/30).

[41 FR 34257, Aug. 13, 1976, as amended at 43 FR 45354, Oct. 2, 1978]

§ 4.83a Impairment of central visual acuity.

The percentage evaluation will be found from table V by intersecting the horizontal row appropriate for the Snellen index for one eye and the vertical column appropriate to the Snellen index of the other eye. For example, if one eye has a Snellen index of 5/200 (1.5/60) and the other eye has a Snellen index of 20/70 (6/21), the percentage evaluation is found in the third horizontal row from the bottom and the fourth vertical column from the left. The evaluation is 50 percent and the diagnostic code 6073.

[41 FR 11297, Mar. 18, 1976, as amended at 43 FR 45354, Oct. 2, 1978]

§ 4.84 Differences between distant and near visual acuity.

Where there is a substantial difference between the near and distant corrected vision, the case should be referred to the Director, Compensation and Pension Service.

[40 FR 42537, Sept. 15, 1975]

§ 4.84a Schedule of ratings—eye.

| DISEASES OF THE EYE | | Rating |
|---------------------|----------------------------------|--------|
| 6000 | Uveitis | |
| 6001 | Keratitis | |
| 6002 | Scleritis | |
| 6003 | Iritis | |
| 6004 | Cyclitis | |
| 6005 | Choroiditis | |
| 6006 | Retinitis | |
| 6007 | Hemorrhage, intra-ocular, recent | |

§ 4.84a

38 CFR Ch. I (7-1-01 Edition)

| DISEASES OF THE EYE—Continued | | Rat- ing |
|-------------------------------|--|-------------|
| 6008 | Retina, detachment of | |
| 6009 | Eye, injury of, unhealed: The above disabilities, in chronic form, are to be rated from 10 percent to 100 percent for impairment of visual acuity or field loss, pain, rest-requirements, or episodic incapacity, combining an additional rating of 10 percent during continuance of active pathology. Minimum rating during active pathology | 10 |
| 6010 | Eye, tuberculosis of, active or inactive: Active | 100 |
| | Inactive: See §§ 4.88b and 4.89. | |
| 6011 | Retina, localized scars, atrophy, or irregularities of, centrally located, with irregular, duplicated enlarged or diminished image: Unilateral or bilateral | 10 |
| 6012 | Glaucoma, congestive or inflammatory: Frequent attacks of considerable duration; during continuance of actual total disability | 100 |
| | Or, rate as iritis, diagnostic Code 6003. | |
| 6013 | Glaucoma, simple, primary, noncongestive: Rate on impairment of visual acuity or field loss. Minimum rating | 10 |
| 6014 | New growths, malignant (eyeball only): Pending completion of operation or other indicated treatment | 100 |
| | Healed; rate on residuals. | |
| 6015 | New growths, benign (eyeball and adnexa, other than superficial) Rate on impaired vision, minimum | 10 |
| | Healed; rate on residuals. | |
| 6016 | Nystagmus, central | 10 |
| 6017 | Conjunctivitis, trachomatous, chronic: Active; rate for impairment of visual acuity; minimum rating while there is active pathology | 30 |
| | Healed; rate on residuals, if no residuals | 0 |
| 6018 | Conjunctivitis, other, chronic: Active, with objective symptoms | 10 |
| | Healed; rate on residuals, if no residuals | 0 |
| 6019 | Ptosis, unilateral or bilateral: Pupil wholly obscured. Rate equivalent to 5/200 (1.5/60). Pupile one-half or more obscured. Rate equivalent to 20/100 (6/30). With less interference with vision. Rate as disfigurement. | |
| 6020 | Ectropion: Bilateral | 20 |
| | Unilateral | 10 |
| 6021 | Entropion: Bilateral | 20 |
| | Unilateral | 10 |
| 6022 | Lagophthalmos: Bilateral | 20 |
| | Unilateral | 10 |

| DISEASES OF THE EYE—Continued | | Rat- ing |
|-------------------------------|---|-------------|
| 6023 | Eyebrows, loss of, complete, unilateral or bilateral | 10 |
| 6024 | Eyelashes, loss of, complete, unilateral or bilateral | 10 |
| 6025 | Epiphora (lacrymal duct, interference with, from any cause): Bilateral | 20 |
| | Unilateral | 10 |
| 6026 | Neuritis, optic: Rate underlying disease, and combine impairment of visual acuity or field loss. | |
| 6027 | Cataract, traumatic: Preoperative. Rate on impairment of vision. Postoperative. Rate on impairment of vision and aphakia. | |
| 6028 | Cataract, senile, and others: Preoperative. Rate on impairment of vision. Postoperative. Rate on impairment of vision and aphakia. | |
| 6029 | Aphakia: Bilateral or unilateral | 30 |
| | NOTE: The 30 percent rating prescribed for aphakia is a minimum rating to be applied to the unilateral or bilateral condition and is not to be combined with any other rating for impaired vision. When only one eye is aphakic, the eye having poorer corrected visual acuity will be rated on the basis of its acuity without correction. When both eyes are aphakic, both will be rated on corrected vision. The corrected vision of one or both aphakic eyes will be taken one step worse than the ascertained value, however, not better than 20/70 (6/21). Combined ratings for disabilities of the same eye should not exceed the amount for total loss of vision of that eye unless there is an enucleation or a serious cosmetic defect added to the total loss of vision. | |
| 6030 | Accommodation, paralysis of | 20 |
| 6031 | Dacryocystitis Rate as epiphora. | |
| 6032 | Eyelids, loss of portion of: Rate as disfigurement. (See diseases of the skin.) | |
| 6033 | Lens, crystalline, dislocation of: Rate as aphakia. | |
| 6034 | Pterygium: Rate for loss of vision, if any. | |
| 6035 | Keratoconus: To be evaluated on impairment of corrected visual acuity using contact lenses. NOTE: When contact lenses are medically required for keratoconus, either unilateral or bilateral, the minimum rating will be 30 percent. | |

TABLE IV—TABLE FOR RATING BILATERAL BLINDNESS OR BLINDNESS COMBINED WITH HEARING LOSS WITH DICTATOR’S CODE AND 38 CFR CITATIONS

| Vision one eye | Vision other eye | | | Plus service-connected Hearing loss | | | | |
|---|--|---|--|--|---|---|---|-------------------------------------|
| | 5/200 (1.5/60) or less | Light perception only | No light perception or anatomical loss | Total deafness one ear | 10% or 20% at least one ear SC | 30% at least one ear SC | 40% at least one ear SC | 60% or more at least one ear SC |
| 5/200 (1.5/60) or less. | L ¹ Code LB-1 38 CFR 3.350(b)(2). | L+½ ¹ Code LB-2 38 CFR 3.350(f)(2)(i). | M Code MB-2 a or b 38 CFR 3.350(f)(2)(ii). | Add ½ step Code PB-1 38 CFR 3.350(f)(2)(iv). | No additional SMC. | Add a full step Code PB-3 38 CFR 3.350(f)(2)(vi). | Add a full step Code PB-3 38 CFR 3.350(f)(2)(vi). | O Code OB-1 38 CFR 3.350(e)(1)(iii) |
| Light perception only. | | M Code MB-1 a 38 CFR 3.350(c)(1)(iv). | M+½ Code MB-3 a or b 38 CFR 3.350(f)(iii). | O Code OB-2 38 CFR 3.350(e)(1)(iv). | Add ½ step Code PB-2 38 CFR 3.350(f)(2)(v). | Add a full step Code PB-3 38 CFR 3.350(f)(2)(iv). | O Code OB-2 38 CFR 3.350(e)(1)(iv). | O Code OB-1 38 CFR 3.350(e)(1)(iii) |
| No light perception or anatomical loss. | | | N Code NB-1 a-b or c 38 CFR 3.350(d)(4). | O Code OB-2 38 CFR 3.350(e)(1)(iv). | Add ½ step Code PB-2 38 CFR 3.350(f)(2)(v). | Add full step Code PB-3 38 CFR 3.350(f)(2)(vi). | O Code OB-2 38 CFR 3.350(e)(1)(iv). | O Code OB-1 38 CFR 3.350(e)(1)(iii) |

¹ With need for aid and attendance qualifies for Subpar. m. code MB-1, b; 38 CFR 3.350(c)(1)(v).

NOTE.—(1) Any of the additional SMC payable under Dictator’s Codes PB-1, PB-2, or PB-3 is not to exceed the rate payable under Subpar. O. (2) If in addition to any of the above the veteran has the service-connected loss or loss of use of an extremity, additional SMC is payable, not to exceed the rate payable under Subpar. O. See Dictator’s Codes PB-4, PB-5, PB-6, and 38 CFR 3.350(f)(2)(vii) (A), (B), (C).

(Authority: 38 U.S.C. 1115)

§ 4.84a

38 CFR Ch. I (7-1-01 Edition)

IMPAIRMENT OF CENTRAL VISUAL ACUITY

IMPAIRMENT OF CENTRAL VISUAL ACUITY—
Continued

| | Rat- ing |
|--|-------------|
| 6061 Anatomical loss both eyes | 5 100 |
| 6062 Blindness in both eyes having only light perception | 5 100 |
| Anatomical loss of 1 eye: | |
| 6063 In the other eye 5/200 (1.5/60) | 5 100 |
| 6064 In the other eye 10/200 (3/60) | 6 90 |
| 6064 In the other eye 15/200 (4.5/60) | 6 80 |
| 6064 In the other eye 20/200 (6/60) | 6 70 |
| 6065 In the other eye 20/100 (6/30) | 6 60 |
| 6065 In the other eye 20/70 (6/21) | 6 60 |
| 6065 In the other eye 20/50 (6/15) | 6 50 |
| 6066 In the other eye 20/40 (6/12) | 6 40 |
| Blindness in 1 eye, having only light perception: | |
| 6067 In the other eye 5/200 (1.5/60) | 5 100 |
| 6068 In the other eye 10/200 (3/60) | 5 90 |
| 6068 In the other eye 15/200 (4.5/60) | 5 80 |
| 6068 In the other eye 20/200 (6/60) | 5 70 |
| 6069 In the other eye 20/100 (6/30) | 5 60 |
| 6069 In the other eye 20/70 (6/21) | 5 50 |
| 6069 In the other eye 20/50 (6/15) | 5 40 |
| 6070 In the other eye 20/40 (6/12) | 5 30 |
| Vision in 1 eye 5/200 (1.5/60): | |
| 6071 In the other eye 5/200 (1.5/60) | 5 100 |
| 6072 In the other eye 10/200 (3/60) | 90 |
| 6072 In the other eye 15/200 (4.5/60) | 80 |
| 6072 In the other eye 20/200 (6/60) | 70 |
| 6073 In the other eye 20/100 (6/30) | 60 |
| 6073 In the other eye 20/70 (6/21) | 50 |
| 6073 In the other eye 20/50 (6/15) | 40 |
| 6074 In the other eye 20/40 (6/12) | 30 |
| Vision in 1 eye 10/200 (3/60): | |
| 6075 In the other eye 10/200 (3/60) | 90 |
| 6075 In the other eye 15/200 (4.5/60) | 80 |
| 6075 In the other eye 20/200 (6/60) | 70 |
| 6076 In the other eye 20/100 (6/30) | 60 |

| | Rat- ing |
|---|-------------|
| 6076 In the other eye 20/70 (6/21) | 50 |
| 6076 In the other eye 20/50 (6/15) | 40 |
| 6077 In the other eye 20/40 (6/12) | 30 |
| Vision in 1 eye 15/200 (4.5/60): | |
| 6075 In the other eye 15/200 (4.5/60) | 80 |
| 6075 In the other eye 20/200 (6/60) | 70 |
| 6076 In the other eye 20/100 (6/30) | 60 |
| 6076 In the other eye 20/70 (6/21) | 40 |
| 6076 In the other eye 20/50 (6/15) | 30 |
| 6077 In the other eye 20/40 (6/12) | 20 |
| Vision in 1 eye 20/200 (6/60): | |
| 6075 In the other eye 20/200 (6/60) | 70 |
| 6076 In the other eye 20/100 (6/30) | 60 |
| 6076 In the other eye 20/70 (6/21) | 40 |
| 6076 In the other eye 20/50 (6/15) | 30 |
| 6077 In the other eye 20/40 (6/12) | 20 |
| Vision in 1 eye 20/100 (6/30): | |
| 6078 In the other eye 20/100 (6/30) | 50 |
| 6078 In the other eye 20/70 (6/21) | 30 |
| 6078 In the other eye 20/50 (6/15) | 20 |
| 6079 In the other eye 20/40 (6/12) | 10 |
| Vision in 1 eye 20/50 (6/15): | |
| 6078 In the other eye 20/50 (6/15) | 10 |
| 6079 In the other eye 20/40 (6/12) | 10 |
| Vision in 1 eye 20/40 (6/12): | |
| In the other eye 20/40 (6/12) | 0 |

⁵ Also entitled to special monthly compensation.
⁶ Add 10% if artificial eye cannot be worn; also entitled to special monthly compensation.

TABLE V—RATINGS FOR CENTRAL VISUAL ACUITY IMPAIRMENT
[With Diagnostic Code]

| Vision in one eye | Vision in other eye | | | | | | | | Light perception only/anatomical loss |
|-----------------------|---------------------|--------------|--------------|---------------|---------------|-----------------|---------------|----------------|---------------------------------------|
| | 20/40 (6/12) | 20/50 (6/15) | 20/70 (6/21) | 20/100 (6/30) | 20/200 (6/60) | 15/200 (4.5/60) | 10/200 (3/60) | 5/200 (1.5/60) | |
| 20/40 (6/12) | 0 | | | | | | | | |
| 20/50 (6/15) | 10 (6079) | 10 (6078) | | | | | | | |
| 20/70 (6/21) | 10 (6079) | 20 (6078) | 30 (6078) | | | | | | |
| 20/100 (6/30) | 10 (6079) | 20 (6078) | 30 (6078) | 50 (6078) | | | | | |
| 20/200 (6/60) | 20 (6077) | 30 (6076) | 40 (6076) | 60 (6076) | 70 (6075) | | | | |
| 15/200 (4.5/60) | 20 (6077) | 30 (6076) | 40 (6076) | 60 (6076) | 70 (6075) | 80 (6075) | | | |
| 10/200 (3/60) | 30 (6077) | 40 (6076) | 50 (6076) | 60 (6076) | 70 (6075) | 80 (6075) | 90 (6075) | | |
| 5/200 (1.5/60) | 30 (6074) | 40 (6073) | 50 (6073) | 60 (6073) | 70 (6072) | 80 (6072) | 90 (6072) | 5 100 (6071) | |
| Light perception only | 5 30 | 5 40 | 5 50 | 5 60 | 5 70 | 5 80 | 5 90 | 5 100 | 5 100 |

TABLE V—RATINGS FOR CENTRAL VISUAL ACUITY IMPAIRMENT—Continued
[With Diagnostic Code]

| Vision in one eye | Vision in other eye | | | | | | | | Light perception only/anatomical loss |
|----------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|-------------------------|---------------------------------------|
| | 20/40 (6/12) | 20/50 (6/15) | 20/70 (6/21) | 20/100 (6/30) | 20/200 (6/60) | 15/200 (4.5/60) | 10/200 (3/60) | 5/200 (1.5/60) | |
| | (6070) | (6069) | (6069) | (6069) | (6068) | (6068) | (6068) | (6067) | (6062) |
| Anatomical loss of one eye | ⁶ 40 (6066) | ⁶ 50 (6065) | ⁶ 60 (6065) | ⁶ 60 (6065) | ⁶ 70 (6064) | ⁶ 80 (6064) | ⁶ 90 (6064) | ⁵ 100 (6063) | ⁵ 100 (6061) |

⁵Also entitled to special monthly compensation.

⁶Add 10 percent if artificial eye cannot be worn; also entitled to special monthly compensation.

RATINGS FOR IMPAIRMENT OF FIELD VISION

| | Rating |
|---|--------|
| 6080 Field vision, impairment of: | |
| Homonymous hemianopsia | 30 |
| Field, visual, loss of temporal half: | |
| Bilateral | 30 |
| Unilateral | 10 |
| Or rate as 20/70 (6/21). | |
| Field, visual, loss of nasal half: | |
| Bilateral | 20 |
| Unilateral | 10 |
| Or rate as 20/50 (6/15). | |
| Field, visual, concentric contraction of: | |
| To 5°: | |
| Bilateral | 100 |
| Unilateral | 30 |
| Or rate as 5/200 (1.5/60). | |
| To 15° but not to 5°: | |
| Bilateral | 70 |
| Unilateral | 20 |
| Or rate as 20/200 (6/60). | |
| To 30° but not to 15°: | |
| Bilateral | 50 |
| Unilateral | 10 |
| Or rate as 20/100 (6/30). | |
| To 45° but not to 30°: | |
| Bilateral | 30 |
| Unilateral | 10 |
| Or rate as 20/70 (6/21): | |
| To 60° but not to 45°: | |
| Bilateral | 20 |
| Unilateral | 10 |
| Or rate as 20/50 (6/15). | |
| Note (1): Correct diagnosis reflecting disease or injury should be cited.. | |
| Note (2): Demonstrable pathology commensurate with the functional loss will be required. The concentric contraction ratings require contraction within the stated degrees, temporally; the nasal contraction may be less. The alternative ratings are to be employed when there is ratable defect of visual acuity, or a different impairment of the visual field in the other eye. Concentric contraction resulting from demonstrable pathology to 5 degrees or less will be considered on a parity with reduction of central visual acuity to 5/200 (1.5/60) or less for all purposes including entitlement under § 3.350(b)(2) of this chapter; not however, for the purpose of § 3.350(a) of this chapter. Entitlement on account of blindness requiring regular aid and attendance, § 3.350(c) of this chapter, will continue to be determined on the facts in the individual case.. | |
| 6081 Scotoma, pathological, unilateral: | |

RATINGS FOR IMPAIRMENT OF FIELD VISION—Continued

| | Rating |
|--|--------|
| Large or centrally located, minimum | 10 |
| NOTE: Rate on loss of central visual acuity or impairment of field vision. Do not combine with any other rating for visual impairment. | |

RATINGS FOR IMPAIRMENT OF MUSCLE FUNCTION [6090 Diplopia (double vision)]

| Degree of diplopia | Equivalent visual acuity |
|--|--------------------------|
| (a) Central 20° | 5/200 |
| (b) 21° to 30°: | |
| (1) Down | 15/200 |
| (2) Lateral | 20/100 |
| (3) Up | 20/70 |
| (c) 31° to 40°: | |
| (1) Down | 20/200 |
| (2) Lateral | 20/70 |
| (3) Up | 20/40 |
| Note: (1) Correct diagnosis reflecting disease or injury should be cited.. | |
| Note: (2) The above ratings will be applied to only one eye. Ratings will not be applied for both diplopia and decreased visual acuity or field of vision in the same eye. When diplopia is present and there is also ratable impairment of visual acuity or field of vision of both eyes the above diplopia ratings will be applied to the poorer eye while the better eye is rated according to the best corrected visual acuity or visual field.. | |
| Note: (3) When the diplopia field extends beyond more than one quadrant or more than one range of degrees, the evaluation for diplopia will be based on the quadrant and degree range that provide the highest evaluation.. | |
| Note: (4) When diplopia exists in two individual and separate areas of the same eye, the equivalent visual acuity will be taken one step worse, but no worse than 5/200.. | |
| 6091 Symblepharon.. | |
| Rate as limited muscle function, diagnostic code 6090.. | |
| 6092 Diplopia, due to limited muscle function.. | |
| Rate as diagnostic code 6090.. | |

§ 4.85

38 CFR Ch. I (7-1-01 Edition)

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 40 FR 42537, Sept. 15, 1975; 41 FR 11297, Mar. 18, 1976; 43 FR 45354, Oct. 2, 1978; 51 FR 6411, Feb. 24, 1986; 53 FR 30264, Aug. 11, 1988; 53 FR 50955, Dec. 19, 1988; 57 FR 24364, June 9, 1992]

IMPAIRMENT OF AUDITORY ACUITY

§ 4.85 Evaluation of hearing impairment.

(a) An examination for hearing impairment for VA purposes must be conducted by a state-licensed audiologist and must include a controlled speech discrimination test (Maryland CNC) and a puretone audiometry test. Examinations will be conducted without the use of hearing aids.

(b) Table VI, "Numeric Designation of Hearing Impairment Based on Puretone Threshold Average and Speech Discrimination," is used to determine a Roman numeral designation (I through XI) for hearing impairment based on a combination of the percent of speech discrimination (horizontal rows) and the puretone threshold average (vertical columns). The Roman numeral designation is located at the point where the percentage of speech discrimination and puretone threshold average intersect.

(c) Table VIa, "Numeric Designation of Hearing Impairment Based Only on Puretone Threshold Average," is used to determine a Roman numeral designation (I through XI) for hearing impairment based only on the puretone threshold average. Table VIa will be used when the examiner certifies that

use of the speech discrimination test is not appropriate because of language difficulties, inconsistent speech discrimination scores, etc., or when indicated under the provisions of § 4.86.

(d) "Puretone threshold average," as used in Tables VI and VIa, is the sum of the puretone thresholds at 1000, 2000, 3000 and 4000 Hertz, divided by four. This average is used in all cases (including those in § 4.86) to determine the Roman numeral designation for hearing impairment from Table VI or VIa.

(e) Table VII, "Percentage Evaluations for Hearing Impairment," is used to determine the percentage evaluation by combining the Roman numeral designations for hearing impairment of each ear. The horizontal rows represent the ear having the better hearing and the vertical columns the ear having the poorer hearing. The percentage evaluation is located at the point where the row and column intersect.

(f) If impaired hearing is service-connected in only one ear, in order to determine the percentage evaluation from Table VII, the non-service-connected ear will be assigned a Roman Numeral designation for hearing impairment of I, subject to the provisions of § 3.383 of this chapter.

(g) When evaluating any claim for impaired hearing, refer to § 3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation due either to deafness, or to deafness in combination with other specified disabilities.

(h) *Numeric tables VI, VIa*, and VII.*

TABLE VI
NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ON
PURETONE THRESHOLD AVERAGE AND SPEECH DISCRIMINATION

Puretone Threshold Average

| % of discrimination | 0-41 | 42-49 | 50-57 | 58-65 | 66-73 | 74-81 | 82-89 | 90-97 | 98+ |
|----------------------------|-------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|------------|
| 92-100 | I | I | I | II | II | II | III | III | IV |
| 84-90 | II | II | II | III | III | III | IV | IV | IV |
| 76-82 | III | III | IV | IV | IV | V | V | V | V |
| 68-74 | IV | IV | V | V | VI | VI | VII | VII | VII |
| 60-66 | V | V | VI | VI | VII | VII | VIII | VIII | VIII |
| 52-58 | VI | VI | VII | VII | VIII | VIII | VIII | VIII | IX |
| 44-50 | VII | VII | VIII | VIII | VIII | IX | IX | IX | X |
| 36-42 | VIII | VIII | VIII | IX | IX | IX | X | X | X |
| 0-34 | IX | X | XI | XI | XI | XI | XI | XI | XI |

TABLE VIA*
NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ONLY ON
PURETONE THRESHOLD AVERAGE

Puretone Threshold Average

| 0-41 | 42-48 | 49-55 | 56-62 | 63-69 | 70-76 | 77-83 | 84-90 | 91-97 | 98-104 | 105+ |
|-------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|---------------|-------------|
| I | II | III | IV | V | VI | VII | VIII | IX | X | XI |

* This table is for use only as specified in §§ 4.85 and 4.86.

TABLE VII
PERCENTAGE EVALUATION FOR HEARING IMPAIRMENT
(DIAGNOSTIC CODE 6100)

| | | | | | | | | | | | | |
|-------------------|-------------|-------------------|----------|-----------|-------------|------------|-----------|----------|-----------|------------|-----------|----------|
| | | Poorer Ear | | | | | | | | | | |
| Better Ear | XI | 100* | | | | | | | | | | |
| | X | 90 | 80 | | | | | | | | | |
| | IX | 80 | 70 | 60 | | | | | | | | |
| | VIII | 70 | 60 | 50 | 50 | | | | | | | |
| | VII | 60 | 60 | 50 | 40 | 40 | | | | | | |
| | VI | 50 | 50 | 40 | 40 | 30 | 30 | | | | | |
| | V | 40 | 40 | 40 | 30 | 30 | 20 | 20 | | | | |
| | IV | 30 | 30 | 30 | 20 | 20 | 20 | 10 | 10 | | | |
| | III | 20 | 20 | 20 | 20 | 20 | 10 | 10 | 10 | 0 | | |
| | II | 10 | 10 | 10 | 10 | 10 | 10 | 10 | 0 | 0 | 0 | |
| | I | 10 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | XI | X | IX | VIII | VII | VI | V | IV | III | II | I |

* Review for entitlement to special monthly compensation under §3.350 of this chapter.

[64 FR 25206, May 11, 1999]

§ 4.86 Exceptional patterns of hearing impairment.

(a) When the puretone threshold at each of the four specified frequencies (1000, 2000, 3000, and 4000 Hertz) is 55 decibels or more, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. Each ear will be evaluated separately.

(b) When the puretone threshold is 30 decibels or less at 1000 Hertz, and 70 decibels or more at 2000 Hertz, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. That numeral will then be elevated to the next higher

Department of Veterans Affairs

§ 4.88a

Roman numeral. Each ear will be evaluated separately.

(Authority: 38 U.S.C. 1155)

[64 FR 25209, May 11, 1999]

§ 4.87 Schedule of ratings—ear.

| DISEASES OF THE EAR | | Rating |
|--|-----|--------|
| 6200 Chronic suppurative otitis media, mastoiditis, or cholesteatoma (or any combination): During suppuration, or with aural polyps | | 10 |
| NOTE: Evaluate hearing impairment, and complications such as labyrinthitis, tinnitus, facial nerve paralysis, or bone loss of skull, separately. | | |
| 6201 Chronic nonsuppurative otitis media with effusion (serous otitis media): Rate hearing impairment | | |
| 6202 Otosclerosis: Rate hearing impairment | | |
| 6204 Peripheral vestibular disorders: Dizziness and occasional staggering | 30 | |
| Occasional dizziness | 10 | |
| NOTE: Objective findings supporting the diagnosis of vestibular disequilibrium are required before a compensable evaluation can be assigned under this code. Hearing impairment or suppuration shall be separately rated and combined. | | |
| 6205 Meniere's syndrome (endolymphatic hydrops): Hearing impairment with attacks of vertigo and cerebellar gait occurring more than once weekly, with or without tinnitus | 100 | |
| Hearing impairment with attacks of vertigo and cerebellar gait occurring from one to four times a month, with or without tinnitus | 60 | |
| Hearing impairment with vertigo less than once a month, with or without tinnitus | 30 | |
| NOTE: Evaluate Meniere's syndrome either under these criteria or by separately evaluating vertigo (as a peripheral vestibular disorder), hearing impairment, and tinnitus, whichever method results in a higher overall evaluation. But do not combine an evaluation for hearing impairment, tinnitus, or vertigo with an evaluation under diagnostic code 6205. | | |
| 6207 Loss of auricle: Complete loss of both | 50 | |
| Complete loss of one | 30 | |
| Deformity of one, with loss of one-third or more of the substance | 10 | |
| 6208 Malignant neoplasm of the ear (other than skin only) | 100 | |
| NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, radiation treatment, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based on that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | | |
| 6209 Benign neoplasms of the ear (other than skin only): Rate on impairment of function. | | |
| 6210 Chronic otitis externa: | | |

DISEASES OF THE EAR—Continued

| | Rating |
|---|--------|
| Swelling, dry and scaly or serous discharge, and itching requiring frequent and prolonged treatment | 10 |
| 6211 Tympanic membrane, perforation of | 0 |
| 6260 Tinnitus, recurrent | 10 |
| NOTE: A separate evaluation for tinnitus may be combined with an evaluation under diagnostic codes 6100, 6200, 6204, or other diagnostic code, except when tinnitus supports an evaluation under one of those diagnostic codes. | |

(Authority: 38 U.S.C. 1155)

[64 FR 25210, May 11, 1999]

§ 4.87a Schedule of ratings—other sense organs.

| | Rating |
|---|--------|
| 6275 Sense of smell, complete loss | 10 |
| 6276 Sense of taste, complete loss | 10 |
| NOTE: Evaluation will be assigned under diagnostic codes 6275 or 6276 only if there is an anatomical or pathological basis for the condition. | |

(Authority: 38 U.S.C. 1155)

[64 FR 25210, May 11, 1999]

INFECTIOUS DISEASES, IMMUNE DISORDERS AND NUTRITIONAL DEFICIENCIES

§ 4.88 [Reserved]

§ 4.88a Chronic fatigue syndrome.

(a) For VA purposes, the diagnosis of chronic fatigue syndrome requires:

- (1) new onset of debilitating fatigue severe enough to reduce daily activity to less than 50 percent of the usual level for at least six months; and
- (2) the exclusion, by history, physical examination, and laboratory tests, of all other clinical conditions that may produce similar symptoms; and
- (3) six or more of the following:
 - (i) acute onset of the condition,
 - (ii) low grade fever,
 - (iii) nonexudative pharyngitis,
 - (iv) palpable or tender cervical or axillary lymph nodes,
 - (v) generalized muscle aches or weakness,
 - (vi) fatigue lasting 24 hours or longer after exercise,

§ 4.88b

38 CFR Ch. I (7-1-01 Edition)

- (vii) headaches (of a type, severity, or pattern that is different from headaches in the pre-morbid state),
- (viii) migratory joint pains,
- (ix) neuropsychologic symptoms,
- (x) sleep disturbance.
- (b) [Reserved]

[59 FR 60902, Nov. 29, 1994]

§ 4.88b Schedule of ratings—infectious diseases, immune disorders and nutritional deficiencies.

| | Rating |
|---|--------|
| 6300 Cholera, Asiatic: As active disease, and for 3 months convalescence | 100 |
| Thereafter rate residuals such as renal necrosis under the appropriate system | |
| 6301 Visceral Leishmaniasis: During treatment for active disease | 100 |
| NOTE: A 100 percent evaluation shall continue beyond the cessation of treatment for active disease. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. Rate residuals such as liver damage or lymphadenopathy under the appropriate system. | |
| 6302 Leprosy (Hansen's Disease): As active disease | 100 |
| NOTE: A 100 percent evaluation shall continue beyond the date that an examining physician has determined that this has become inactive. Six months after the date of inactivity, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. Rate residuals such as skin lesions or peripheral neuropathy under the appropriate system. | |
| 6304 Malaria: As active disease | 100 |
| NOTE: The diagnosis of malaria depends on the identification of the malarial parasites in blood smears. If the veteran served in an endemic area and presents signs and symptoms compatible with malaria, the diagnosis may be based on clinical grounds alone. Relapses must be confirmed by the presence of malarial parasites in blood smears. Thereafter rate residuals such as liver or spleen damage under the appropriate system | |
| 6305 Lymphatic Filariasis: As active disease | 100 |
| Thereafter rate residuals such as epididymitis or lymphangitis under the appropriate system | |
| 6306 Bartonellosis: As active disease, and for 3 months convalescence | 100 |
| Thereafter rate residuals such as skin lesions under the appropriate system | |
| 6307 Plague: As active disease | 100 |
| Thereafter rate residuals such as lymphadenopathy under the appropriate system | |
| 6308 Relapsing Fever: As active disease | 100 |
| Thereafter rate residuals such as liver or spleen damage or central nervous system involvement under the appropriate system | |
| 6309 Rheumatic fever: As active disease | 100 |
| Thereafter rate residuals such as heart damage under the appropriate system | |
| 6310 Syphilis, and other treponemal infections: Rate the complications of nervous system, vascular system, eyes or ears. (See DC 7004, syphilitic heart disease, DC 8013, cerebrospinal syphilis, DC 8014, meningovascular syphilis, DC 8015, tabes dorsalis, and DC 9301, dementia associated with central nervous system syphilis) | |
| 6311 Tuberculosis, miliary: As active disease | 100 |
| Inactive: See §§ 4.88c and 4.89. | |
| 6313 Avitaminosis: Marked mental changes, moist dermatitis, inability to retain adequate nourishment, exhaustion, and cachexia | 100 |
| With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor | 60 |
| With stomatitis, diarrhea, and symmetrical dermatitis | 40 |
| With stomatitis, or achlorhydria, or diarrhea | 20 |
| Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discomfort, weakness, inability to concentrate and irritability | 10 |
| 6314 Beriberi: As active disease: With congestive heart failure, anasarca, or Wernicke-Korsakoff syndrome | 100 |
| With cardiomegaly, or; with peripheral neuropathy with footdrop or atrophy of thigh or calf muscles | 60 |
| With peripheral neuropathy with absent knee or ankle jerks and loss of sensation, or; with symptoms such as weakness, fatigue, anorexia, dizziness, heaviness and stiffness of legs, headache or sleep disturbance | 30 |
| Thereafter rate residuals under the appropriate body system. | |
| 6315 Pellagra: Marked mental changes, moist dermatitis, inability to retain adequate nourishment, exhaustion, and cachexia | 100 |

Department of Veterans Affairs

§ 4.88b

| | Rating |
|---|--------|
| With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor | 60 |
| With stomatitis, diarrhea, and symmetrical dermatitis | 40 |
| With stomatitis, or achlorhydria, or diarrhea | 20 |
| Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discomfort, weakness, inability to concentrate and irritability | 10 |
| 6316 Brucellosis: | |
| As active disease | 100 |
| Thereafter rate residuals such as liver or spleen damage or meningitis under the appropriate system | |
| 6317 Typhus, scrub: | |
| As active disease, and for 3 months convalescence | 100 |
| Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system | |
| 6318 Melioidosis: | |
| As active disease | 100 |
| Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system | |
| 6319 Lyme Disease: | |
| As active disease | 100 |
| Thereafter rate residuals such as arthritis under the appropriate system | |
| 6320 Parasitic diseases otherwise not specified: | |
| As active disease | 100 |
| Thereafter rate residuals such as spleen or liver damage under the appropriate system | |
| 6350 Lupus erythematosus, systemic (disseminated): | |
| Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health | 100 |
| Exacerbations lasting a week or more, 2 or 3 times per year | 60 |
| Exacerbations once or twice a year or symptomatic during the past 2 years | 10 |
| NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation. | |
| 6351 HIV-Related Illness: | |
| AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions | 100 |
| Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neoplasm | 60 |
| Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis | 30 |
| Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations | 10 |
| Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count | 0 |
| NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredited medical institution. | |
| NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated separately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above. | |
| 6354 Chronic Fatigue Syndrome (CFS): | |
| Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms: | |
| Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care | 100 |
| Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year | 60 |
| Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year | 40 |
| Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year | 20 |
| Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication | 10 |
| NOTE: For the purpose of evaluating this disability, the condition will be considered incapacitating only while it requires bed rest and treatment by a physician. | |

[61 FR 39875, July 31, 1996]

§ 4.88c

§ 4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19, 1968.

| | Rating |
|---|--------|
| For 1 year after date of inactivity, following active tuberculosis | 100 |
| Thereafter: Rate residuals under the specific body system or systems affected. | |
| Following the total rating for the 1 year period after date of inactivity, the schedular evaluation for residuals of nonpulmonary tuberculosis, i.e., ankylosis, surgical removal of a part, etc., will be assigned under the appropriate diagnostic code for the residual preceded by the diagnostic code for tuberculosis of the body part affected. For example, tuberculosis of the hip joint with residual ankylosis would be coded 5001-5250. Where there are existing residuals of pulmonary and nonpulmonary conditions, the evaluations for residual separate functional impairment may be combined. | |
| Where there are existing pulmonary and nonpulmonary conditions, the total rating for the 1 year, after attainment of inactivity, may not be applied to both conditions during the same period. However, the total rating during the 1-year period for the pulmonary or for the nonpulmonary condition will be utilized, combined with evaluation for residuals of the condition not covered by the 1-year total evaluation, so as to allow any additional benefit provided during such period. | |

[34 FR 5062, Mar. 11, 1969. Redesignated at 59 FR 60902, Nov. 29, 1994]

§ 4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

Public Law 90-493 repealed section 356 of title 38, United States Code which provided graduated ratings for inactive tuberculosis. The repealed section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90-493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For use in rating cases in which the protective provisions of Pub. L. 90-493 apply, the former evaluations are retained in this section.

| | Rating |
|---|--------|
| For 2 years after date of inactivity, following active tuberculosis, which was clinically identified during service or subsequently | 100 |
| Thereafter, for 4 years, or in any event, to 6 years after date of inactivity | 50 |
| Thereafter, for 5 years, or to 11 years after date of inactivity | 30 |
| Thereafter, in the absence of a schedular compensable permanent residual | 0 |

| | Rating |
|---|--------|
| Following the total rating for the 2-year period after date of inactivity, the schedular evaluation for residuals of nonpulmonary tuberculosis, i.e., ankylosis, surgical removal of a part, etc., if in excess of 50 percent or 30 percent will be assigned under the appropriate diagnostic code for the specific residual preceded by the diagnostic code for tuberculosis of the body part affected. For example, tuberculosis of the hipjoint with residual ankylosis would be coded 5001-5250. | |
| The graduated ratings for nonpulmonary tuberculosis will not be combined with residuals of nonpulmonary tuberculosis unless the graduated rating and the rating for residual disability cover separate functional losses, e.g., graduated ratings for tuberculosis of the kidney and residuals of tuberculosis of the spine. Where there are existing pulmonary and nonpulmonary conditions, the graduated evaluation for the pulmonary, or for the nonpulmonary, condition will be utilized, combined with evaluations for residuals of the condition not covered by the graduated evaluation utilized, so as to provide the higher evaluation over such period. | |
| The ending dates of all graduated ratings of nonpulmonary tuberculosis will be controlled by the date of attainment of inactivity. | |
| These ratings are applicable only to veterans with nonpulmonary tuberculosis active on or after October 10, 1949. | |

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 43 FR 45361, Oct. 2, 1978]

THE RESPIRATORY SYSTEM

§ 4.96 Special provisions regarding evaluation of respiratory conditions.

(a) *Rating coexisting respiratory conditions.* Ratings under diagnostic codes 6600 through 6817 and 6822 through 6847 will not be combined with each other. Where there is lung or pleural involvement, ratings under diagnostic codes 6819 and 6820 will not be combined with each other or with diagnostic codes 6600 through 6817 or 6822 through 6847. A single rating will be assigned under the diagnostic code which reflects the predominant disability with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation. However, in cases protected by the provisions of Pub. L. 90-493, the graduated ratings of 50 and 30 percent for inactive tuberculosis will not be elevated.

(b) *Rating "protected" tuberculosis cases.* Public Law 90-493 repealed section 356 of title 38, United States Code which had provided graduated ratings for inactive tuberculosis. The repealed

Department of Veterans Affairs

§ 4.97

section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90-493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For application in rating cases in which the protective provisions of Pub. L. 90-493 apply the former evaluations pertaining to pulmonary tuberculosis are retained in § 4.97.

(c) *Special monthly compensation.* When evaluating any claim involving complete organic aphonia, refer to § 3.350 of this chapter to determine

whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, there are other conditions in this section which under certain circumstances also establish entitlement to special monthly compensation.

(Authority: 38 U.S.C. 1155)

[34 FR 5062, Mar. 11, 1969, as amended at 61 FR 46727, Sept. 5, 1996]

§ 4.97 Schedule of ratings—respiratory system.

| | Rating |
|--|--------|
| DISEASES OF THE NOSE AND THROAT | |
| 6502 Septum, nasal, deviation of: Traumatic only, With 50-percent obstruction of the nasal passage on both sides or complete obstruction on one side | 10 |
| 6504 Nose, loss of part of, or scars: Exposing both nasal passages | 30 |
| Loss of part of one ala, or other obvious disfigurement | 10 |
| Note: Or evaluate as DC 7800, scars, disfiguring, head, face, or neck. | |
| 6510 Sinusitis, pansinusitis, chronic. | |
| 6511 Sinusitis, ethmoid, chronic. | |
| 6512 Sinusitis, frontal, chronic. | |
| 6513 Sinusitis, maxillary, chronic. | |
| 6514 Sinusitis, sphenoid, chronic. | |
| General Rating Formula for Sinusitis (DC's 6510 through 6514): | |
| Following radical surgery with chronic osteomyelitis, or; near constant sinusitis characterized by headaches, pain and tenderness of affected sinus, and purulent discharge or crusting after repeated surgeries | 50 |
| Three or more incapacitating episodes per year of sinusitis requiring prolonged (lasting four to six weeks) antibiotic treatment, or; more than six non-incapacitating episodes per year of sinusitis characterized by headaches, pain, and purulent discharge or crusting | 30 |
| One or two incapacitating episodes per year of sinusitis requiring prolonged (lasting four to six weeks) antibiotic treatment, or; three to six non-incapacitating episodes per year of sinusitis characterized by headaches, pain, and purulent discharge or crusting | 10 |
| Detected by X-ray only | 0 |
| Note: An incapacitating episode of sinusitis means one that requires bed rest and treatment by a physician. | |
| 6515 Laryngitis, tuberculous, active or inactive. Rate under §§ 4.88c or 4.89, whichever is appropriate. | |
| 6516 Laryngitis, chronic: Hoarseness, with thickening or nodules of cords, polyps, submucous infiltration, or pre-malignant changes on biopsy | 30 |
| Hoarseness, with inflammation of cords or mucous membrane | 10 |
| 6518 Laryngectomy, total. | 100 |
| Rate the residuals of partial laryngectomy as laryngitis (DC 6516), aphonia (DC 6519), or stenosis of larynx (DC 6520). | |
| 6519 Aphonia, complete organic: Constant inability to communicate by speech | 100 |
| Constant inability to speak above a whisper | 60 |
| Note: Evaluate incomplete aphonia as laryngitis, chronic (DC 6516). | |
| 6520 Larynx, stenosis of, including residuals of laryngeal trauma (unilateral or bilateral): Forced expiratory volume in one second (FEV-1) less than 40 percent of predicted value, with Flow-Volume Loop compatible with upper airway obstruction, or; permanent tracheostomy | 100 |
| FEV-1 of 40- to 55-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction | 60 |
| FEV-1 of 56- to 70-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction | 30 |
| FEV-1 of 71- to 80-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction | 10 |
| Note: Or evaluate as aphonia (DC 6519). | |
| 6521 Pharynx, injuries to: Stricture or obstruction of pharynx or nasopharynx, or; absence of soft palate secondary to trauma, chemical burn, or granulomatous disease, or; paralysis of soft palate with swallowing difficulty (nasal regurgitation) and speech impairment | 50 |

| | Rating |
|--|--------|
| 6522 Allergic or vasomotor rhinitis: | |
| With polyps | 30 |
| Without polyps, but with greater than 50-percent obstruction of nasal passage on both sides or complete obstruction on one side | 10 |
| 6523 Bacterial rhinitis: | |
| Rhinoscleroma | 50 |
| With permanent hypertrophy of turbinates and with greater than 50-percent obstruction of nasal passage on both sides or complete obstruction on one side | 10 |
| 6524 Granulomatous rhinitis: | |
| Wegener's granulomatosis, lethal midline granuloma | 100 |
| Other types of granulomatous infection | 20 |

DISEASES OF THE TRACHEA AND BRONCHI

| | |
|---|-----|
| 6600 Bronchitis, chronic: | |
| FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy | 100 |
| FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) | 60 |
| FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent predicted | 30 |
| FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted | 10 |
| 6601 Bronchiectasis: | |
| With incapacitating episodes of infection of at least six weeks total duration per year | 100 |
| With incapacitating episodes of infection of four to six weeks total duration per year, or; near constant findings of cough with purulent sputum associated with anorexia, weight loss, and frank hemoptysis and requiring antibiotic usage almost continuously | 60 |
| With incapacitating episodes of infection of two to four weeks total duration per year, or; daily productive cough with sputum that is at times purulent or blood-tinged and that requires prolonged (lasting four to six weeks) antibiotic usage more than twice a year | 30 |
| Intermittent productive cough with acute infection requiring a course of antibiotics at least twice a year | 10 |
| Or rate according to pulmonary impairment as for chronic bronchitis (DC 6600). | |
| Note: An incapacitating episode is one that requires bedrest and treatment by a physician. | |
| 6602 Asthma, bronchial: | |
| FEV-1 less than 40-percent predicted, or; FEV-1/FVC less than 40 percent, or; more than one attack per week with episodes of respiratory failure, or; requires daily use of systemic (oral or parenteral) high dose corticosteroids or immuno-suppressive medications | 100 |
| FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; at least monthly visits to a physician for required care of exacerbations, or; intermittent (at least three per year) courses of systemic (oral or parenteral) corticosteroids | 60 |
| FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; daily inhalational or oral bronchodilator therapy, or; inhalational anti-inflammatory medication | 30 |
| FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; intermittent inhalational or oral bronchodilator therapy | 10 |
| Note: In the absence of clinical findings of asthma at time of examination, a verified history of asthmatic attacks must be of record. | |
| 6603 Emphysema, pulmonary: | |
| FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy | 100 |
| FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) | 60 |
| FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent predicted | 30 |
| FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted | 10 |
| 6604 Chronic obstructive pulmonary disease: | |
| FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy | 100 |
| FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) | 60 |

Department of Veterans Affairs

§ 4.97

| | Rating |
|--|--------|
| FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent predicted | 30 |
| FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted | 10 |

DISEASES OF THE LUNGS AND PLEURA—TUBERCULOSIS
Ratings for Pulmonary Tuberculosis Entitled on August 19, 1968

| | |
|--|-----|
| 6701 Tuberculosis, pulmonary, chronic, far advanced, active | 100 |
| 6702 Tuberculosis, pulmonary, chronic, moderately advanced, active | 100 |
| 6703 Tuberculosis, pulmonary, chronic, minimal, active | 100 |
| 6704 Tuberculosis, pulmonary, chronic, active, advancement unspecified | 100 |
| 6721 Tuberculosis, pulmonary, chronic, far advanced, inactive. | |
| 6722 Tuberculosis, pulmonary, chronic, moderately advanced, inactive. | |
| 6723 Tuberculosis, pulmonary, chronic, minimal, inactive. | |
| 6724 Tuberculosis, pulmonary, chronic, inactive, advancement unspecified. | |
| General Rating Formula for Inactive Pulmonary Tuberculosis: For two years after date of inactivity, following active tuberculosis, which was clinically identified during service or subsequently | 100 |
| Thereafter for four years, or in any event, to six years after date of inactivity | 50 |
| Thereafter, for five years, or to eleven years after date of inactivity | 30 |
| Following far advanced lesions diagnosed at any time while the disease process was active, minimum | 30 |
| Following moderately advanced lesions, provided there is continued disability, emphysema, dyspnea on exertion, impairment of health, etc | 20 |
| Otherwise | 0 |
| Note (1): The 100-percent rating under codes 6701 through 6724 is not subject to a requirement of precedent hospital treatment. It will be reduced to 50 percent for failure to submit to examination or to follow prescribed treatment upon report to that effect from the medical authorities. When a veteran is placed on the 100-percent rating for inactive tuberculosis, the medical authorities will be appropriately notified of the fact, and of the necessity, as given in footnote 1 to 38 U.S.C. 1156 (and formerly in 38 U.S.C. 356, which has been repealed by Public Law 90-493), to notify the Adjudication Division in the event of failure to submit to examination or to follow treatment. | |
| Note (2): The graduated 50-percent and 30-percent ratings and the permanent 30 percent and 20 percent ratings for inactive pulmonary tuberculosis are not to be combined with ratings for other respiratory disabilities. Following thoracoplasty the rating will be for removal of ribs combined with the rating for collapsed lung. Resection of the ribs incident to thoracoplasty will be rated as removal. | |

Ratings for Pulmonary Tuberculosis Initially Evaluated After August 19, 1968

| | |
|---|-----|
| 6730 Tuberculosis, pulmonary, chronic, active | 100 |
| Note: Active pulmonary tuberculosis will be considered permanently and totally disabling for non-service-connected pension purposes in the following circumstances: (a) Associated with active tuberculosis involving other than the respiratory system. (b) With severe associated symptoms or with extensive cavity formation. (c) Reactivated cases, generally. (d) With advancement of lesions on successive examinations or while under treatment. (e) Without retrogression of lesions or other evidence of material improvement at the end of six months hospitalization or without change of diagnosis from "active" at the end of 12 months hospitalization. Material improvement means lessening or absence of clinical symptoms, and X-ray findings of a stationary or retrogressive lesion. | |
| 6731 Tuberculosis, pulmonary, chronic, inactive: Depending on the specific findings, rate residuals as interstitial lung disease, restrictive lung disease, or, when obstructive lung disease is the major residual, as chronic bronchitis (DC 6600). Rate thoracoplasty as removal of ribs under DC 5297. | |
| Note: A mandatory examination will be requested immediately following notification that active tuberculosis evaluated under DC 6730 has become inactive. Any change in evaluation will be carried out under the provisions of § 3.105(e). | |
| 6732 Pleurisy, tuberculous, active or inactive: Rate under §§ 4.88c or 4.89, whichever is appropriate. | |

NONTUBERCULOUS DISEASES

| | |
|--|-----|
| 6817 Pulmonary Vascular Disease: Primary pulmonary hypertension, or; chronic pulmonary thromboembolism with evidence of pulmonary hypertension, right ventricular hypertrophy, or cor pulmonale, or; pulmonary hypertension secondary to other obstructive disease of pulmonary arteries or veins with evidence of right ventricular hypertrophy or cor pulmonale | 100 |
| Chronic pulmonary thromboembolism requiring anticoagulant therapy, or; following inferior vena cava surgery without evidence of pulmonary hypertension or right ventricular dysfunction | 60 |
| Symptomatic, following resolution of acute pulmonary embolism | 30 |
| Asymptomatic, following resolution of pulmonary thromboembolism | 0 |
| Note: Evaluate other residuals following pulmonary embolism under the most appropriate diagnostic code, such as chronic bronchitis (DC 6600) or chronic pleural effusion or fibrosis (DC 6844), but do not combine that evaluation with any of the above evaluations. | |
| 6819 Neoplasms, malignant, any specified part of respiratory system exclusive of skin growths | 100 |

| | Rating |
|--|-----------------------|
| <p>Note: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.</p> | |
| 6820 Neoplasms, benign, any specified part of respiratory system. Evaluate using an appropriate respiratory analogy. | |
| Bacterial Infections of the Lung | |
| 6822 Actinomycosis. 6823 Nocardiosis. 6824 Chronic lung abscess. General Rating Formula for Bacterial Infections of the Lung (diagnostic codes 6822 through 6824): Active infection with systemic symptoms such as fever, night sweats, weight loss, or hemoptysis Depending on the specific findings, rate residuals as interstitial lung disease, restrictive lung disease, or, when obstructive lung disease is the major residual, as chronic bronchitis (DC 6600). | 100 |
| Interstitial Lung Disease | |
| 6825 Diffuse interstitial fibrosis (interstitial pneumonitis, fibrosing alveolitis). 6826 Desquamative interstitial pneumonitis. 6827 Pulmonary alveolar proteinosis. 6828 Eosinophilic granuloma of lung. 6829 Drug-induced pulmonary pneumonitis and fibrosis. 6830 Radiation-induced pulmonary pneumonitis and fibrosis. 6831 Hypersensitivity pneumonitis (extrinsic allergic alveolitis). 6832 Pneumoconiosis (silicosis, anthracosis, etc.). 6833 Asbestosis. General Rating Formula for Interstitial Lung Disease (diagnostic codes 6825 through 6833): Forced Vital Capacity (FVC) less than 50-percent predicted, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption with cardiorespiratory limitation, or; cor pulmonale or pulmonary hypertension, or; requires outpatient oxygen therapy FVC of 50- to 64-percent predicted, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum exercise capacity of 15 to 20 ml/kg/min oxygen consumption with cardiorespiratory limitation FVC of 65- to 74-percent predicted, or; DLCO (SB) of 56- to 65-percent predicted FVC of 75- to 80-percent predicted, or; DLCO (SB) of 66- to 80-percent predicted | 100 60 30 10 |
| Mycotic Lung Disease | |
| 6834 Histoplasmosis of lung. 6835 Coccidioidomycosis. 6836 Blastomycosis. 6837 Cryptococcosis. 6838 Aspergillosis. 6839 Mucormycosis. General Rating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive cough Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough Healed and inactive mycotic lesions, asymptomatic | 100 50 30 0 |
| <p>Note: Coccidioidomycosis has an incubation period up to 21 days, and the disseminated phase is ordinarily manifest within six months of the primary phase. However, there are instances of dissemination delayed up to many years after the initial infection which may have been unrecognized. Accordingly, when service connection is under consideration in the absence of record or other evidence of the disease in service, service in southwestern United States where the disease is endemic and absence of prolonged residence in this locality before or after service will be the deciding factor.</p> | |
| Restrictive Lung Disease | |
| 6840 Diaphragm paralysis or paresis. 6841 Spinal cord injury with respiratory insufficiency. 6842 Kyphoscoliosis, pectus excavatum, pectus carinatum. 6843 Traumatic chest wall defect, pneumothorax, hernia, etc. 6844 Post-surgical residual (lobectomy, pneumonectomy, etc.). 6845 Chronic pleural effusion or fibrosis. | |

| | Rating |
|---|--------|
| General Rating Formula for Restrictive Lung Disease (diagnostic codes 6840 through 6845): | |
| FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy | 100 |
| FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) | 60 |
| FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent predicted | 30 |
| FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted | 10 |
| Or rate primary disorder. | |
| Note (1): A 100-percent rating shall be assigned for pleurisy with empyema, with or without pleurocutaneous fistula, until resolved. | |
| Note (2): Following episodes of total spontaneous pneumothorax, a rating of 100 percent shall be assigned as of the date of hospital admission and shall continue for three months from the first day of the month after hospital discharge. | |
| Note (3): Gunshot wounds of the pleural cavity with bullet or missile retained in lung, pain or discomfort on exertion, or with scattered rales or some limitation of excursion of diaphragm or of lower chest expansion shall be rated at least 20-percent disabling. Disabling injuries of shoulder girdle muscles (Groups I to IV) shall be separately rated and combined with ratings for respiratory involvement. Involvement of Muscle Group XXI (DC 5321), however, will not be separately rated. | |
| 6846 Sarcoidosis: | |
| Cor pulmonale, or; cardiac involvement with congestive heart failure, or; progressive pulmonary disease with fever, night sweats, and weight loss despite treatment | 100 |
| Pulmonary involvement requiring systemic high dose (therapeutic) corticosteroids for control | 60 |
| Pulmonary involvement with persistent symptoms requiring chronic low dose (maintenance) or intermittent corticosteroids | 30 |
| Chronic hilar adenopathy or stable lung infiltrates without symptoms or physiologic impairment | 0 |
| Or rate active disease or residuals as chronic bronchitis (DC 6600) and extra-pulmonary involvement under specific body system involved. | |
| 6847 Sleep Apnea Syndromes (Obstructive, Central, Mixed): | |
| Chronic respiratory failure with carbon dioxide retention or cor pulmonale, or; requires tracheostomy | 100 |
| Requires use of breathing assistance device such as continuous airway pressure (CPAP) machine | 50 |
| Persistent day-time hypersomnolence | 30 |
| Asymptomatic but with documented sleep disorder breathing | 0 |

¹ Review for entitlement to special monthly compensation under § 3.350 of this chapter.

[61 FR 46728, Sept. 5, 1996]

THE CARDIOVASCULAR SYSTEM

DISEASES OF THE HEART—Continued

§§ 4.100–4.103 [Reserved]

§ 4.104 Schedule of ratings—cardiovascular system.

DISEASES OF THE HEART

| | Rating |
|--|--------|
|--|--------|

NOTE (1): Evaluate cor pulmonale, which is a form of secondary heart disease, as part of the pulmonary condition that causes it.

| | Rating |
|--|--------|
|--|--------|

NOTE (2): One MET (metabolic equivalent) is the energy cost of standing quietly at rest and represents an oxygen uptake of 3.5 milliliters per kilogram of body weight per minute. When the level of METs at which dyspnea, fatigue, angina, dizziness, or syncope develops is required for evaluation, and a laboratory determination of METs by exercise testing cannot be done for medical reasons, an estimation by a medical examiner of the level of activity (expressed in METs and supported by specific examples, such as slow stair climbing or shoveling snow) that results in dyspnea, fatigue, angina, dizziness, or syncope may be used.

7000 Valvular heart disease (including rheumatic heart disease):
 During active infection with valvular heart damage and for three months following cessation of therapy for the active infection

100

DISEASES OF THE HEART—Continued

DISEASES OF THE HEART—Continued

| | Rating |
|--|--------|
| Thereafter, with valvular heart disease (documented by findings on physical examination and either echocardiogram, Doppler echocardiogram, or cardiac catheterization) resulting in: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electro-cardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| 7001 Endocarditis: | |
| For three months following cessation of therapy for active infection with cardiac involvement | 100 |
| Thereafter, with endocarditis (documented by findings on physical examination and either echocardiogram, Doppler echocardiogram, or cardiac catheterization) resulting in: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| 7002 Pericarditis: | |
| For three months following cessation of therapy for active infection with cardiac involvement | 100 |
| Thereafter, with documented pericarditis resulting in: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |

| | Rating |
|--|--------|
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electro-cardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| 7003 Pericardial adhesions: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electro-cardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| 7004 Syphilitic heart disease: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |

NOTE: Evaluate syphilitic aortic aneurysms under DC 7110 (aortic aneurysm).

DISEASES OF THE HEART—Continued

DISEASES OF THE HEART—Continued

| | Rating |
|--|--------|
| 7005 Arteriosclerotic heart disease (Coronary artery disease): | |
| With documented coronary artery disease resulting in: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| NOTE: If nonservice-connected arteriosclerotic heart disease is superimposed on service-connected valvular or other non-arteriosclerotic heart disease, request a medical opinion as to which condition is causing the current signs and symptoms. | |
| 7006 Myocardial infarction: | |
| During and for three months following myocardial infarction, documented by laboratory tests | 100 |
| Thereafter: | |
| With history of documented myocardial infarction, resulting in: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| 7007 Hypertensive heart disease: | |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |

| | Rating |
|---|--------|
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| 7008 Hyperthyroid heart disease: | |
| Include as part of the overall evaluation for hyperthyroidism under DC 7900. However, when atrial fibrillation is present, hyperthyroidism may be evaluated either under DC 7900 or under DC 7010 (supraventricular arrhythmia), whichever results in a higher evaluation. | |
| 7010 Supraventricular arrhythmias: | |
| Paroxysmal atrial fibrillation or other supraventricular tachycardia, with more than four episodes per year documented by ECG or Holter monitor | 30 |
| Permanent atrial fibrillation (one atrial fibrillation), or; one to four episodes per year of paroxysmal atrial fibrillation or other supraventricular tachycardia documented by ECG or Holter monitor | 10 |
| 7011 Ventricular arrhythmias (sustained): | |
| For indefinite period from date of hospital admission for initial evaluation and medical therapy for a sustained ventricular arrhythmia, or; for indefinite period from date of hospital admission for ventricular aneurysmectomy, or; with an automatic implantable Cardioverter-Defibrillator (AICD) in place | 100 |
| Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |

DISEASES OF THE HEART—Continued

DISEASES OF THE HEART—Continued

| | Rating |
|--|--------|
| NOTE: A rating of 100 percent shall be assigned from the date of hospital admission for initial evaluation and medical therapy for a sustained ventricular arrhythmia or for ventricular aneurysmectomy. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. | |
| 7015 Atrioventricular block: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication or a pacemaker required | 10 |
| NOTE: Unusual cases of arrhythmia such as atrioventricular block associated with a supraventricular arrhythmia or pathological bradycardia should be submitted to the Director, Compensation and Pension Service. Simple delayed P-R conduction time, in the absence of other evidence of cardiac disease, is not a disability. | |
| 7016 Heart valve replacement (prosthesis): For indefinite period following date of hospital admission for valve replacement | 100 |
| Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |

| | Rating |
|---|--------|
| NOTE: A rating of 100 percent shall be assigned as of the date of hospital admission for valve replacement. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. | |
| 7017 Coronary bypass surgery: For three months following hospital admission for surgery | 100 |
| Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| 7018 Implantable cardiac pacemakers: For two months following hospital admission for implantation or reimplantation | 100 |
| Thereafter: Evaluate as supraventricular arrhythmias (DC 7010), ventricular arrhythmias (DC 7011), or atrioventricular block (DC 7015). Minimum | 10 |
| NOTE: Evaluate implantable Cardioverter-Defibrillators (AICD's) under DC 7011. | |
| 7019 Cardiac transplantation: For an indefinite period from date of hospital admission for cardiac transplantation | 100 |
| Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Minimum | 30 |
| NOTE: A rating of 100 percent shall be assigned as of the date of hospital admission for cardiac transplantation. One year following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. | |

DISEASES OF THE HEART—Continued

DISEASES OF THE HEART—Continued

| | Rating |
|---|--------|
| 7020 Cardiomyopathy: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent | 100 |
| More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent | 60 |
| Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray | 30 |
| Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required | 10 |
| Diseases of the Arteries and Veins | |
| 7101 Hypertensive vascular disease (hypertension and isolated systolic hypertension): Diastolic pressure predominantly 130 or more | 60 |
| Diastolic pressure predominantly 120 or more | 40 |
| Diastolic pressure predominantly 110 or more, or; systolic pressure predominantly 200 or more | 20 |
| Diastolic pressure predominantly 100 or more, or; systolic pressure predominantly 160 or more, or; minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control | 10 |
| NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm. | |
| NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation. | |
| 7110 Aortic aneurysm: If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion) | 100 |
| Precluding exertion | 60 |
| Evaluate residuals of surgical correction according to organ systems affected. | |
| NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. | |
| 7111 Aneurysm, any large artery: | |

| | Rating |
|---|--------|
| If symptomatic, or; for indefinite period from date of hospital admission for surgical correction | 100 |
| Following surgery: Ischemic limb pain at rest, and; either deep ischemic ulcers or ankle/brachial index of 0.4 or less | 100 |
| Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; persistent coldness of the extremity, one or more deep ischemic ulcers, or ankle/brachial index of 0.5 or less | 60 |
| Claudication on walking between 25 and 100 yards on a level grade at 2 miles per hour, and; trophic changes (thin skin, absence of hair, dystrophic nails) or ankle/brachial index of 0.7 or less | 40 |
| Claudication on walking more than 100 yards, and; diminished peripheral pulses or ankle/brachial index of 0.9 or less | 20 |
| NOTE (1): The ankle/brachial index is the ratio of the systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater. | |
| NOTE (2): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor, if applicable. | |
| NOTE (3): A rating of 100 percent shall be assigned as of the date of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. | |
| 7112 Aneurysm, any small artery: Asymptomatic | 0 |
| NOTE: If symptomatic, evaluate according to body system affected. Following surgery, evaluate residuals under the body system affected. | |
| 7113 Arteriovenous fistula, traumatic: With high output heart failure | 100 |
| Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia | 60 |
| Without cardiac involvement but with edema, stasis dermatitis, and either ulceration or cellulitis: Lower extremity | 50 |
| Upper extremity | 40 |
| With edema or stasis dermatitis: Lower extremity | 30 |
| Upper extremity | 20 |
| 7114 Arteriosclerosis obliterans: Ischemic limb pain at rest, and; either deep ischemic ulcers or ankle/brachial index of 0.4 or less | 100 |
| Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; either persistent coldness of the extremity or ankle/brachial index of 0.5 or less | 60 |
| Claudication on walking between 25 and 100 yards on a level grade at 2 miles per hour, and; trophic changes (thin skin, absence of hair, dystrophic nails) or ankle/brachial index of 0.7 or less | 40 |

DISEASES OF THE HEART—Continued

| | Rating |
|--|--------|
| Claudication on walking more than 100 yards, and; diminished peripheral pulses or ankle/brachial index of 0.9 or less | 20 |
| NOTE (1): The ankle/brachial index is the ratio of the systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater. | |
| NOTE (2): Evaluate residuals of aortic and large arterial bypass surgery or arterial graft as arteriosclerosis obliterans. | |
| NOTE (3): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor (§ 4.26), if applicable. | |
| 7115 Thrombo-angiitis obliterans (Buerger's Disease): | |
| Ischemic limb pain at rest, and; either deep ischemic ulcers or ankle/brachial index of 0.4 or less | 100 |
| Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; either persistent coldness of the extremity or ankle/brachial index of 0.5 or less | 60 |
| Claudication on walking between 25 and 100 yards on a level grade at 2 miles per hour, and; trophic changes (thin skin, absence of hair, dystrophic nails) or ankle/brachial index of 0.7 or less | 40 |
| Claudication on walking more than 100 yards, and; diminished peripheral pulses or ankle/brachial index of 0.9 or less | 20 |
| NOTE (1): The ankle/brachial index is the ratio of the systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater. | |
| NOTE (2): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor (§ 4.26), if applicable. | |
| 7117 Raynaud's syndrome: | |
| With two or more digital ulcers plus autoamputation of one or more digits and history of characteristic attacks | 100 |
| With two or more digital ulcers and history of characteristic attacks | 60 |
| Characteristic attacks occurring at least daily | 40 |
| Characteristic attacks occurring four to six times a week | 20 |
| Characteristic attacks occurring one to three times a week | 10 |
| NOTE: For purposes of this section, characteristic attacks consist of sequential color changes of the digits of one or more extremities lasting minutes to hours, sometimes with pain and paresthesias, and precipitated by exposure to cold or by emotional upsets. These evaluations are for the disease as a whole, regardless of the number of extremities involved or whether the nose and ears are involved. | |
| 7118 Angioneurotic edema: | |
| Attacks without laryngeal involvement lasting one to seven days or longer and occurring more than eight times a year, or; attacks with laryngeal involvement of any duration occurring more than twice a year | 40 |

DISEASES OF THE HEART—Continued

| | Rating |
|--|--------|
| Attacks without laryngeal involvement lasting one to seven days and occurring five to eight times a year, or; attacks with laryngeal involvement of any duration occurring once or twice a year | 20 |
| Attacks without laryngeal involvement lasting one to seven days and occurring two to four times a year | 10 |
| 7119 Erythromelalgia: | |
| Characteristic attacks that occur more than once a day, last an average of more than two hours each, respond poorly to treatment, and that restrict most routine daily activities | 100 |
| Characteristic attacks that occur more than once a day, last an average of more than two hours each, and respond poorly to treatment, but that do not restrict most routine daily activities | 60 |
| Characteristic attacks that occur daily or more often but that respond to treatment | 30 |
| Characteristic attacks that occur less than daily but at least three times a week and that respond to treatment | 10 |
| NOTE: For purposes of this section, a characteristic attack of erythromelalgia consists of burning pain in the hands, feet, or both, usually bilateral and symmetrical, with increased skin temperature and redness, occurring at warm ambient temperatures. These evaluations are for the disease as a whole, regardless of the number of extremities involved. | |
| 7120 Varicose veins: | |
| With the following findings attributed to the effects of varicose veins: Massive board-like edema with constant pain at rest | 100 |
| Persistent edema or subcutaneous induration, stasis pigmentation or eczema, and persistent ulceration | 60 |
| Persistent edema and stasis pigmentation or eczema, with or without intermittent ulceration | 40 |
| Persistent edema, incompletely relieved by elevation of extremity, with or without beginning stasis pigmentation or eczema | 20 |
| Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by elevation of extremity or compression hosiery | 10 |
| Asymptomatic palpable or visible varicose veins | 0 |
| NOTE: These evaluations are for involvement of a single extremity. If more than one extremity is involved, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor (§ 4.26), if applicable. | |
| 7121 Post-phlebotic syndrome of any etiology: | |
| With the following findings attributed to venous disease: | |
| Massive board-like edema with constant pain at rest | 100 |
| Persistent edema or subcutaneous induration, stasis pigmentation or eczema, and persistent ulceration | 60 |
| Persistent edema and stasis pigmentation or eczema, with or without intermittent ulceration | 40 |

DISEASES OF THE HEART—Continued

THE DIGESTIVE SYSTEM

| | Rat- ing |
|---|-------------|
| Persistent edema, incompletely relieved by elevation of extremity, with or without beginning stasis pigmentation or eczema | 20 |
| Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by elevation of extremity or compression hosiery | 10 |
| Asymptomatic palpable or visible varicose veins | 0 |
| NOTE: These evaluations are for involvement of a single extremity. If more than one extremity is involved, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable. | |
| 7122 Cold injury residuals: With the following in affected parts: | |
| Arthralgia or other pain, numbness, or cold sensitivity plus two or more of the following: tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteoarthritis) | 30 |
| Arthralgia or other pain, numbness, or cold sensitivity plus tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, or X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteoarthritis) | 20 |
| Arthralgia or other pain, numbness, or cold sensitivity | 10 |
| NOTE (1): Separately evaluate amputations of fingers or toes, and complications such as squamous cell carcinoma at the site of a cold injury scar or peripheral neuropathy, under other diagnostic codes. Separately evaluate other disabilities that have been diagnosed as the residual effects of cold injury, such as Raynaud's phenomenon, muscle atrophy, etc., unless they are used to support an evaluation under diagnostic code 7122. | |
| NOTE (2): Evaluate each affected part (e.g., hand, foot, ear, nose) separately and combine the ratings in accordance with §§4.25 and 4.26. | |
| 7123 Soft tissue sarcoma (of vascular origin) | 100 |
| NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | |

§4.110 Ulcers.

Experience has shown that the term "peptic ulcer" is not sufficiently specific for rating purposes. Manifest differences in ulcers of the stomach or duodenum in comparison with those at an anastomotic stoma are sufficiently recognized as to warrant two separate graduated descriptions. In evaluating the ulcer, care should be taken that the findings adequately identify the particular location.

§4.111 Postgastrectomy syndromes.

There are various postgastrectomy symptoms which may occur following anastomotic operations of the stomach. When present, those occurring during or immediately after eating and known as the "dumping syndrome" are characterized by gastrointestinal complaints and generalized symptoms simulating hypoglycemia; those occurring from 1 to 3 hours after eating usually present definite manifestations of hypoglycemia.

§4.112 Weight loss.

Minor weight loss or greater losses of weight for periods of brief duration are not considered of importance in rating. Rather, weight loss becomes of importance where there is appreciable loss which is sustained over a period of time. In evaluating weight loss generally, consideration will be given not only to standard age, height, and weight tables, but also to the particular individual's predominant weight pattern as reflected by the records. The use of the term "inability to gain weight" indicates that there has been a significant weight loss with inability to regain it despite appropriate therapy.

EFFECTIVE DATE NOTE: At 66 FR 29488, May 31, 2001, §4.112 was revised, effective July 2, 2001. For the convenience of the user, the revised text is set forth as follows:

§4.112 Weight loss.

For purposes of evaluating conditions in §4.114, the term "substantial weight loss" means a loss of greater than 20 percent of the individual's baseline weight, sustained for three months or longer; and the term "minor weight loss" means a weight loss of 10 to 20 percent of the individual's baseline weight.

(Authority: 38 U.S.C. 1155)

[62 FR 65219, Dec. 11, 1997, as amended at 63 FR 37779, July 14, 1998]

§4.113

sustained for three months or longer. The term “inability to gain weight” means that there has been substantial weight loss with inability to regain it despite appropriate therapy. “Baseline weight” means the average weight for the two-year-period preceding onset of the disease.

(Authority: 38 U.S.C. 1155)

§4.113 Coexisting abdominal conditions.

There are diseases of the digestive system, particularly within the abdomen, which, while differing in the site of pathology, produce a common disability picture characterized in the main by varying degrees of abdominal distress or pain, anemia and disturbances in nutrition. Consequently, certain coexisting diseases in this area, as indicated in the instruction under the title “Diseases of the Digestive System,” do not lend themselves to distinct and separate disability evaluations without violating the fundamental principle relating to pyramiding as outlined in §4.14.

§4.114 Schedule of ratings—digestive system.

Ratings under diagnostic codes 7301 to 7329, inclusive, 7331, 7342, and 7345 to 7348 inclusive will not be combined with each other. A single evaluation will be assigned under the diagnostic code which reflects the predominant disability picture, with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation.

| | Rating |
|--|-----------------|
| 7200 Mouth, injuries of. Rate as for disfigurement and impairment of function of mastication. | |
| 7201 Lips, injuries of. Rate as for disfigurement of face. | |
| 7202 Tongue, loss of whole or part: With inability to communicate by speech One-half or more With marked speech impairment | 100 60 30 |
| 7203 Esophagus, stricture of: Permitting passage of liquids only, with marked impairment of general health Severe, permitting liquids only Moderate | 80 50 30 |
| 7204 Esophagus, spasm of (cardiospasm). If not amenable to dilation, rate as for the degree of obstruction (stricture). | |
| 7205 Esophagus, diverticulum of, acquired. Rate as for obstruction (stricture). | |
| 7301 Peritoneum, adhesions of: | |

38 CFR Ch. I (7-1-01 Edition)

| | Rating |
|---|--------|
| Severe; definite partial obstruction shown by X-ray, with frequent and prolonged episodes of severe colic distension, nausea or vomiting, following severe peritonitis, ruptured appendix, perforated ulcer, or operation with drainage | 50 |
| Moderately severe; partial obstruction manifested by delayed motility of barium meal and less frequent and less prolonged episodes of pain | 30 |
| Moderate; pulling pain on attempting work or aggravated by movements of the body, or occasional episodes of colic pain, nausea, constipation (perhaps alternating with diarrhea) or abdominal distension | 10 |
| Mild | 0 |
| NOTE: Ratings for adhesions will be considered when there is history of operative or other traumatic or infectious (intraabdominal) process, and at least two of the following: disturbance of motility, actual partial obstruction, reflex disturbances, presence of pain. | |
| 7304 Ulcer, gastric. | |
| 7305 Ulcer, duodenal: Severe; pain only partially relieved by standard ulcer therapy, periodic vomiting, recurrent hematemesis or melena, with manifestations of anemia and weight loss productive of definite impairment of health | 60 |
| Moderately severe; less than severe but with impairment of health manifested by anemia and weight loss; or recurrent incapacitating episodes averaging 10 days or more in duration at least four or more times a year | 40 |
| Moderate; recurring episodes of severe symptoms two or three times a year averaging 10 days in duration; or with continuous moderate manifestations | 20 |
| Mild; with recurring symptoms once or twice yearly | 10 |
| 7306 Ulcer, marginal (gastrojejunal): Pronounced; periodic or continuous pain unrelieved by standard ulcer therapy with periodic vomiting, recurring melena or hematemesis, and weight loss. Totally incapacitating | 100 |
| Severe; same as pronounced with less pronounced and less continuous symptoms with definite impairment of health | 60 |
| Moderately severe; intercurrent episodes of abdominal pain at least once a month partially or completely relieved by ulcer therapy, mild and transient episodes of vomiting or melena | 40 |
| Moderate; with episodes of recurring symptoms several times a year | 20 |
| Mild; with brief episodes of recurring symptoms once or twice yearly | 10 |
| 7307 Gastritis, hypertrophic (identified by gastro-scope): Chronic; with severe hemorrhages, or large ulcerated or eroded areas | 60 |
| Chronic; with multiple small eroded or ulcerated areas, and symptoms | 30 |
| Chronic; with small nodular lesions, and symptoms | 10 |
| Gastritis, atrophic. A complication of a number of diseases, including pernicious anemia. Rate the underlying condition. | |
| 7308 Postgastrectomy syndromes: Severe; associated with nausea, sweating, circulatory disturbance after meals, diarrhea, hypoglycemic symptoms, and weight loss with malnutrition and anemia | 60 |

Department of Veterans Affairs

§4.114

| | Rating | | Rating |
|---|--------|--|--------|
| Moderate; less frequent episodes of epigastric disorders with characteristic mild circulatory symptoms after meals but with diarrhea and weight loss | 40 | 7323 Colitis, ulcerative: Pronounced; resulting in marked malnutrition, anemia, and general debility, or with serious complication as liver abscess | 100 |
| Mild; infrequent episodes of epigastric distress with characteristic mild circulatory symptoms or continuous mild manifestations | 20 | Severe; with numerous attacks a year and malnutrition, the health only fair during remissions | 60 |
| 7309 Stomach, stenosis of. Rate as for gastric ulcer. | | Moderately severe; with frequent exacerbations | 30 |
| 7310 Stomach, injury of, residuals. Rate as peritoneal adhesions. | | Moderate; with infrequent exacerbations | 10 |
| 7311 Liver, injury of. With residual disability, rate as peritoneal adhesions. | | 7324 Distomiasis, intestinal or hepatic: Severe symptoms | 30 |
| Healed, no residuals | 0 | Moderate symptoms | 10 |
| 7312 Liver, cirrhosis of: Pronounced; aggravation of the symptoms for moderate and severe, necessitating frequent tapping | 100 | Mild or no symptoms | 0 |
| Severe; ascites requiring infrequent tapping, or recurrent hemorrhage from esophageal varices, aggravated symptoms and impaired health | 70 | 7325 Enteritis, chronic. Rate as for irritable colon syndrome. | |
| Moderately severe; liver definitely enlarged with abdominal distention due to early ascites and with muscle wasting and loss of strength | 50 | 7326 Enterocolitis, chronic. Rate as for irritable colon syndrome. | |
| Moderate; with dilation of superficial abdominal veins, chronic dyspepsia, slight loss of weight or impairment of health | 30 | 7327 Diverticulitis. Rate as for irritable colon syndrome, peritoneal adhesions, or colitis, ulcerative, depending upon the predominant disability picture. | |
| 7313 Liver, abscess of, residuals: With severe symptoms | 30 | 7328 Intestine, small, resection of: With marked interference with absorption and nutrition, manifested by severe impairment of health objectively supported by examination findings including material weight loss | 60 |
| With moderate symptoms | 20 | With definite interference with absorption and nutrition, manifested by impairment of health objectively supported by examination findings including definite weight loss | 40 |
| 7314 Cholecystitis, chronic: Severe; frequent attacks of gall bladder colic | 30 | Symptomatic with diarrhea, anemia and inability to gain weight | 20 |
| Moderate; gall bladder dyspepsia, confirmed by X-ray technique, and with infrequent attacks (not over two or three a year) of gall bladder colic, with or without jaundice | 10 | NOTE: Where residual adhesions constitute the predominant disability, rate under diagnostic code 7301. | |
| Mild | 0 | 7329 Intestine, large, resection of: With severe symptoms, objectively supported by examination findings | 40 |
| 7315 Cholelithiasis, chronic. Rate as for chronic cholecystitis. | | With moderate symptoms | 20 |
| 7316 Cholangitis, chronic. Rate as for chronic cholecystitis. | | With slight symptoms | 10 |
| 7317 Gall bladder, injury of. Rate as for peritoneal adhesions. | | NOTE: Where residual adhesions constitute the predominant disability, rate under diagnostic code 7301. | |
| 7318 Gall bladder, removal of: With severe symptoms | 30 | 7330 Intestine, fistula of, persistent, or after attempt at operative closure: Copious and frequent, fecal discharge | 100 |
| With mild symptoms | 10 | Constant or frequent, fecal discharge | 60 |
| Nonsymptomatic | 0 | Slight infrequent, fecal discharge | 30 |
| Spleen, disease or injury of. See Hemic and Lymphatic Systems. | | Healed; rate for peritoneal adhesions. | |
| 7319 Irritable colon syndrome (spastic colitis, mucous colitis, etc.): Severe; diarrhea, or alternating diarrhea and constipation, with more or less constant abdominal distress | 30 | 7331 Peritonitis, tuberculous, active or inactive: Active | 100 |
| Moderate; frequent episodes of bowel disturbance with abdominal distress | 10 | Inactive: See §§ 4.88b and 4.89. | |
| Mild; disturbances of bowel function with occasional episodes of abdominal distress | 0 | 7332 Rectum and anus, impairment of sphincter control: Complete loss of sphincter control | 100 |
| 7321 Amebiasis: Mild gastrointestinal disturbances, lower abdominal cramps, nausea, gaseous distention, chronic constipation interrupted by diarrhea | 10 | Extensive leakage and fairly frequent involuntary bowel movements | 60 |
| Asymptomatic | 0 | Occasional involuntary bowel movements, necessitating wearing of pad | 30 |
| NOTE: Amebiasis with or without liver abscess is parallel in symptomatology with ulcerative colitis and should be rated on the scale provided for the latter. Similarly, lung abscess due to amebiasis will be rated under the respiratory system schedule, diagnostic code 6809. | | Constant slight, or occasional moderate leakage | 10 |
| 7322 Dysentery, bacillary. Rate as for ulcerative colitis.. | | Healed or slight, without leakage | 0 |
| | | 7333 Rectum and anus, stricture of: Requiring colostomy | 100 |
| | | Great reduction of lumen, or extensive leakage .. | 50 |
| | | Moderate reduction of lumen, or moderate constant leakage | 30 |
| | | 7334 Rectum, prolapse of: Severe (or complete), persistent | 50 |
| | | Moderate, persistent or frequently recurring | 30 |
| | | Mild with constant slight or occasional moderate leakage | 10 |
| | | 7335 Ano, fistula in. Rate as for impairment of sphincter control. | |
| | | 7336 Hemorrhoids, external or internal: | |

§4.114

38 CFR Ch. I (7-1-01 Edition)

| | Rat- ing |
|--|-------------|
| With persistent bleeding and with secondary anemia, or with fissures | 20 |
| Large or thrombotic, irreducible, with excessive redundant tissue, evidencing frequent recurrences | 10 |
| Mild or moderate | 0 |
| 7337 Pruritus ani. Rate for the underlying condition. | |
| 7338 Hernia, inguinal: Large, postoperative, recurrent, not well supported under ordinary conditions and not readily reducible, when considered inoperable | 60 |
| Small, postoperative recurrent, or unoperated irremediable, not well supported by truss, or not readily reducible | 30 |
| Postoperative recurrent, readily reducible and well supported by truss or belt | 10 |
| Not operated, but remediable | 0 |
| Small, reducible, or without true hernia protrusion | 0 |
| NOTE: Add 10 percent for bilateral involvement, provided the second hernia is compensable. This means that the more severely disabling hernia is to be evaluated, and 10 percent, only, added for the second hernia, if the latter is of compensable degree. | |
| 7339 Hernia, ventral, postoperative: Massive, persistent, severe diastasis of recti muscles or extensive diffuse destruction or weakening of muscular and fascial support of abdominal wall so as to be inoperable | 100 |
| Large, not well supported by belt under ordinary conditions | 40 |
| Small, not well supported by belt under ordinary conditions, or healed ventral hernia or postoperative wounds with weakening of abdominal wall and indication for a supporting belt | 20 |
| Wounds, postoperative, healed, no disability, belt not indicated | 0 |
| 7340 Hernia, femoral. Rate as for inguinal hernia. | |
| 7342 Visceroptosis, symptomatic, marked | 10 |
| 7343 New growths, malignant, exclusive of skin growths | 100 |
| NOTE: The rating under diagnostic code 7343 will be continued for 1 year following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. At this point, if there has been no local recurrence or metastases, the rating will be made on residuals. | |
| 7344 New growths, benign, any specified part of digestive system, exclusive of skin growths. The rating will be based on interference with digestion, using any applicable digestive analogy. | |
| 7345 Hepatitis, infectious: With marked liver damage manifest by liver function test and marked gastrointestinal symptoms, or with episodes of several weeks duration aggregating three or more a year and accompanied by disabling symptoms requiring rest therapy | 100 |
| With moderate liver damage and disabling recurrent episodes of gastrointestinal disturbance, fatigue, and mental depression | 60 |
| Minimal liver damage with associated fatigue, anxiety, and gastrointestinal disturbance of lesser degree and frequency but necessitating dietary restriction or other therapeutic measures | 30 |
| Demonstrable liver damage with mild gastrointestinal disturbance | 10 |
| Healed, nonsymptomatic | 0 |

| | Rat- ing |
|---|-------------|
| 7346 Hernia hiatal: Symptoms of pain, vomiting, material weight loss and hematemesis or melena with moderate anemia; or other symptom combinations productive of severe impairment of health | 60 |
| Persistently recurrent epigastric distress with dysphagia, pyrosis, and regurgitation, accompanied by substernal or arm or shoulder pain, productive of considerable impairment of health | 30 |
| With two or more of the symptoms for the 30 percent evaluation of less severity | 10 |
| 7347 Pancreatitis: With frequently recurrent disabling attacks of abdominal pain with few pain free intermissions and with steatorrhea, malabsorption, diarrhea and severe malnutrition | 100 |
| With frequent attacks of abdominal pain, loss of normal body weight and other findings showing continuing pancreatic insufficiency between acute attacks | 60 |
| Moderately severe; with at least 4-7 typical attacks of abdominal pain per year with good remission between attacks | 30 |
| With at least one recurring attack of typical severe abdominal pain in the past year | 10 |
| NOTE 1: Abdominal pain in this condition must be confirmed as resulting from pancreatitis by appropriate laboratory and clinical studies. | |
| NOTE 2: Following total or partial pancreatectomy, rate under above, symptoms, minimum rating 30 percent. | |
| 7348 Vagotomy with pyloroplasty or gastroenterostomy: Followed by demonstrably confirmative postoperative complications of stricture or continuing gastric retention | 40 |
| With symptoms and confirmed diagnosis of alkaline gastritis, or of confirmed persisting diarrhea | 30 |
| Recurrent ulcer with incomplete vagotomy | 20 |
| NOTE: Rate recurrent ulcer following complete vagotomy under diagnostic code 7305, minimum rating 20 percent; and rate dumping syndrome under diagnostic code 7308. | |

[29 FR 6718, May 22, 1964, as amended at 34 FR 5063, Mar. 11, 1969; 40 FR 42540, Sept. 15, 1975; 41 FR 11301, Mar. 18, 1976]

EFFECTIVE DATE NOTE: At 66 FR 29488, May 31, 2001, §4.114 was amended by revising diagnostic codes 7311, 7312, 7343, 7344, and 7345; removing diagnostic code 7313; adding diagnostic codes 7351 and 7354; and adding a new authority citation at the end of the section, effective July 2, 2001. For the convenience of the user, the added and revised text is set forth as follows:

§ 4.114 Schedule of ratings-digestive system.

| | Rat- ing |
|--|-------------|
| * * * * * | |
| 7311 Residuals of injury of the liver: | |

| | Rating | | Rating |
|--|--------|--|--------|
| Depending on the specific residuals, separately evaluate as adhesions of peritoneum (diagnostic code 7301), cirrhosis of liver (diagnostic code 7312), and chronic liver disease without cirrhosis (diagnostic code 7345). | | Daily fatigue, malaise, and anorexia (without weight loss or hepatomegaly), requiring dietary restriction or continuous medication, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least two weeks, but less than four weeks, during the past 12-month period | 20 |
| 7312 Cirrhosis of the liver, primary biliary cirrhosis, or cirrhotic phase of sclerosing cholangitis: Generalized weakness, substantial weight loss, and persistent jaundice, or; with one of the following refractory to treatment: ascites, hepatic encephalopathy, hemorrhage from varices or portal gastropathy (erosive gastritis) | 100 | Intermittent fatigue, malaise, and anorexia, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least one week, but less than two weeks, during the past 12-month period | 10 |
| History of two or more episodes of ascites, hepatic encephalopathy, or hemorrhage from varices or portal gastropathy (erosive gastritis), but with periods of remission between attacks | 70 | Nonsymptomatic | 0 |
| History of one episode of ascites, hepatic encephalopathy, or hemorrhage from varices or portal gastropathy (erosive gastritis) | 50 | NOTE (1): Evaluate sequelae, such as cirrhosis or malignancy of the liver, under an appropriate diagnostic code, but do not use the same signs and symptoms as the basis for evaluation under DC 7354 and under a diagnostic code for sequelae. (See §4.14.) | |
| Portal hypertension and splenomegaly, with weakness, anorexia, abdominal pain, malaise, and at least minor weight loss | 30 | NOTE (2): For purposes of evaluating conditions under diagnostic code 7345, "incapacitating episode" means a period of acute signs and symptoms severe enough to require bed rest and treatment by a physician. | |
| Symptoms such as weakness, anorexia, abdominal pain, and malaise | 10 | NOTE (3): Hepatitis B infection must be confirmed by serologic testing in order to evaluate it under diagnostic code 7345. | |
| NOTE: For evaluation under diagnostic code 7312, documentation of cirrhosis (by biopsy or imaging) and abnormal liver function tests must be present. * * * * * | | * * * * * | |
| 7343 Malignant neoplasms of the digestive system, exclusive of skin growths NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | 100 | 7351 Liver transplant: For an indefinite period from the date of hospital admission for transplant surgery | 100 |
| | | Minimum | 30 |
| | | NOTE: A rating of 100 percent shall be assigned as of the date of hospital admission for transplant surgery and shall continue. One year following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. | |
| 7344 Benign neoplasms, exclusive of skin growths: Evaluate under an appropriate diagnostic code, depending on the predominant disability or the specific residuals after treatment. | | 7354 Hepatitis C (or non-A, non-B hepatitis): With serologic evidence of hepatitis C infection and the following signs and symptoms due to hepatitis C infection: Near-constant debilitating symptoms (such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) | 100 |
| 7345 Chronic liver disease without cirrhosis (including hepatitis B, chronic active hepatitis, autoimmune hepatitis, hemochromatosis, drug-induced hepatitis, etc., but excluding bile duct disorders and hepatitis C): Near-constant debilitating symptoms (such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) | 100 | Daily fatigue, malaise, and anorexia, with substantial weight loss (or other indication of malnutrition), and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least six weeks during the past 12-month period, but not occurring constantly | 60 |
| Daily fatigue, malaise, and anorexia, with substantial weight loss (or other indication of malnutrition), and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least six weeks during the past 12-month period, but not occurring constantly | 60 | Daily fatigue, malaise, and anorexia, with minor weight loss and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least four weeks, but less than six weeks, during the past 12-month period | 40 |
| Daily fatigue, malaise, and anorexia, with minor weight loss and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least four weeks, but less than six weeks, during the past 12-month period | 40 | | |

§4.115

38 CFR Ch. I (7-1-01 Edition)

| | Rating |
|--|--------|
| Daily fatigue, malaise, and anorexia (without weight loss or hepatomegaly), requiring dietary restriction or continuous medication, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least two weeks, but less than four weeks, during the past 12-month period | 20 |
| Intermittent fatigue, malaise, and anorexia, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least one week, but less than two weeks, during the past 12-month period | 10 |
| Nonsymptomatic | 0 |
| NOTE (1): Evaluate sequelae, such as cirrhosis or malignancy of the liver, under an appropriate diagnostic code, but do not use the same signs and symptoms as the basis for evaluation under DC 7354 and under a diagnostic code for sequelae. (See §4.14.) | |
| NOTE (2): For purposes of evaluating conditions under diagnostic code 7354, "incapacitating episode" means a period of acute signs and symptoms severe enough to require bed rest and treatment by a physician. | |

pertension or heart disease will be separately rated.

[41 FR 34258, Aug. 13, 1976, as amended at 59 FR 2527, Jan. 18, 1994]

§4.115a Ratings of the genitourinary system—dysfunctions.

Diseases of the genitourinary system generally result in disabilities related to renal or voiding dysfunctions, infections, or a combination of these. The following section provides descriptions of various levels of disability in each of these symptom areas. Where diagnostic codes refer the decisionmaker to these specific areas dysfunction, only the predominant area of dysfunction shall be considered for rating purposes. Since the areas of dysfunction described below do not cover all symptoms resulting from genitourinary diseases, specific diagnoses may include a description of symptoms assigned to that diagnosis.

(Authority: 38 U.S.C. 1155)

THE GENITOURINARY SYSTEM

§4.115 Nephritis.

Albuminuria alone is not nephritis, nor will the presence of transient albumin and casts following acute febrile illness be taken as nephritis. The glomerular type of nephritis is usually preceded by or associated with severe infectious disease; the onset is sudden, and the course marked by red blood cells, salt retention, and edema; it may clear up entirely or progress to a chronic condition. The nephrosclerotic type, originating in hypertension or arteriosclerosis, develops slowly, with minimum laboratory findings, and is associated with natural progress. Separate ratings are not to be assigned for disability from disease of the heart and any form of nephritis, on account of the close interrelationships of cardiovascular disabilities. If, however, absence of a kidney is the sole renal disability, even if removal was required because of nephritis, the absent kidney and any hypertension or heart disease will be separately rated. Also, in the event that chronic renal disease has progressed to the point where regular dialysis is required, any coexisting hy-

| | Rating |
|---|--------|
| Renal dysfunction: Requiring regular dialysis, or precluding more than sedentary activity from one of the following: persistent edema and albuminuria; or, BUN more than 80mg%; or, creatinine more than 8mg%; or, markedly decreased function of kidney or other organ systems, especially cardiovascular | 100 |
| Persistent edema and albuminuria with BUN 40 to 80mg%; or, creatinine 4 to 8mg%; or, generalized poor health characterized by lethargy, weakness, anorexia, weight loss, or limitation of exertion | 80 |
| Constant albuminuria with some edema; or, definite decrease in kidney function; or, hypertension at least 40 percent disabling under diagnostic code 7101 | 60 |
| Albumin constant or recurring with hyaline and granular casts or red blood cells; or, transient or slight edema or hypertension at least 10 percent disabling under diagnostic code 7101 | 30 |
| Albumin and casts with history of acute nephritis; or, hypertension non-compensable under diagnostic code 7101 | 0 |
| Voiding dysfunction: Rate particular condition as urine leakage, frequency, or obstructed voiding | |
| Continual Urine Leakage, Post Surgical Urinary Diversion, Urinary Incontinence, or Stress Incontinence: Requiring the use of an appliance or the wearing of absorbent materials which must be changed more than 4 times per day | 60 |
| Requiring the wearing of absorbent materials which must be changed 2 to 4 times per day | 40 |
| Requiring the wearing of absorbent materials which must be changed less than 2 times per day | 20 |
| Urinary frequency: Daytime voiding interval less than one hour, or; awakening to void five or more times per night ... | 40 |

Department of Veterans Affairs

§ 4.115b

| | Rating | | Rating |
|---|--------|---|--------|
| Daytime voiding interval between one and two hours, or; awakening to void three to four times per night | 20 | 7502 Nephritis, chronic: Rate as renal dysfunction. | |
| Daytime voiding interval between two and three hours, or; awakening to void two times per night | 10 | 7504 Pyelonephritis, chronic: Rate as renal dysfunction or urinary tract infection, whichever is predominant. | |
| Obstructed voiding: | | | |
| Urinary retention requiring intermittent or continuous catheterization | 30 | 7505 Kidney, tuberculosis of: Rate in accordance with §§ 4.88b or 4.89, whichever is appropriate. | |
| Marked obstructive symptomatology (hesitancy, slow or weak stream, decreased force of stream) with any one or combination of the following: | | 7507 Nephrosclerosis, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation. | |
| 1. Post void residuals greater than 150 cc. | | | |
| 2. Uroflowmetry; markedly diminished peak flow rate (less than 10 cc/sec). | | | |
| 3. Recurrent urinary tract infections secondary to obstruction. | | | |
| 4. Stricture disease requiring periodic dilatation every 2 to 3 months | 10 | | |
| Obstructive symptomatology with or without stricture disease requiring dilatation 1 to 2 times per year | 0 | 7508 Nephrolithiasis: Rate as hydronephrosis, <i>except for</i> recurrent stone formation requiring one or more of the following: | |
| Urinary tract infection: | | 1. diet therapy | |
| Poor renal function: Rate as renal dysfunction. | | 2. drug therapy | |
| Recurrent symptomatic infection requiring drainage/frequent hospitalization (greater than two times/year), and/or requiring continuous intensive management | 30 | 3. invasive or non-invasive procedures more than two times/year | 30 |
| Long-term drug therapy, 1-2 hospitalizations per year and/or requiring intermittent intensive management | 10 | 7509 Hydronephrosis: Severe; Rate as renal dysfunction. | |
| | | Frequent attacks of colic with infection (pyonephrosis), kidney function impaired | 30 |
| | | Frequent attacks of colic, requiring catheter drainage | 20 |
| | | Only an occasional attack of colic, not infected and not requiring catheter drainage | 10 |
| | | 7510 Ureterolithiasis: Rate as hydronephrosis, <i>except for</i> recurrent stone formation requiring one or more of the following: | |
| | | 1. diet therapy | |
| | | 2. drug therapy | |
| | | 3. invasive or non-invasive procedures more than two times/year | 30 |
| | | 7511 Ureter, stricture of: Rate as hydronephrosis, <i>except for</i> recurrent stone formation requiring one or more of the following: | |
| | | 1. diet therapy | |
| | | 2. drug therapy | |
| | | 3. invasive or non-invasive procedures more than two times/year | 30 |
| | | 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. | |
| | | 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction | |
| | | 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant. | |
| | | Postoperative, suprapubic cystostomy | 100 |

[59 FR 2527, Jan. 18, 1994; 59 FR 10676, Mar. 7, 1994]

§ 4.115b Ratings of the genitourinary system—diagnoses.

Note: When evaluating any claim involving loss or loss of use of one or more creative organs, refer to § 3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, there are other conditions in this section which under certain circumstances also establish entitlement to special monthly compensation.

| | | | |
|---|----|--|--|
| 7500 Kidney, removal of one: Minimum evaluation | 30 | | |
| Or rate as renal dysfunction if there is nephritis, infection, or pathology of the other. | | | |
| 7501 Kidney, abscess of: Rate as urinary tract infection | | | |

§4.115b

38 CFR Ch. I (7-1-01 Edition)

| | Rating | | Rating |
|--|--------|---|--------|
| 7517 Bladder, injury of: Rate as voiding dysfunction. | | <p>Note—Following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure, the rating of 100 percent shall continue with a mandatory VA examination at the expiration of six months. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local reoccurrence or metastasis, rate on residuals as voiding dysfunction or renal dysfunction, whichever is predominant.</p> <p>7529 Benign neoplasms of the genitourinary system: Rate as voiding dysfunction or renal dysfunction, whichever is predominant.</p> <p>7530 Chronic renal disease requiring regular dialysis: Rate as renal dysfunction.</p> <p>7531 Kidney transplant: Following transplant surgery 100 Thereafter: Rate on residuals as renal dysfunction, minimum rating 30</p> <p>Note—The 100 percent evaluation shall be assigned as of the date of hospital admission for transplant surgery and shall continue with a mandatory VA examination one year following hospital discharge. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.</p> <p>7532 Renal tubular disorders (such as renal glycosurias, aminoacidurias, renal tubular acidosis, Fanconi's syndrome, Bartter's syndrome, related disorders of Henle's loop and proximal or distal nephron function, etc.): Minimum rating for symptomatic condition 20</p> | |
| 7518 Urethra, stricture of: Rate as voiding dysfunction. | | | |
| 7519 Urethra, fistula of: Rate as voiding dysfunction. | | | |
| Multiple urethroperineal fistulae | 100 | | |
| 7520 Penis, removal of half or more | 30 | | |
| Or rate as voiding dysfunction. | | | |
| 7521 Penis removal of glans | 20 | | |
| Or rate as voiding dysfunction. | | | |
| 7522 Penis, deformity, with loss of erectile power—20 ¹ . | | | |
| 7523 Testis, atrophy complete: Both—20 ¹ One—0 ¹ | | | |
| 7524 Testis, removal: Both—30 ¹ One—0 ¹ | | | |
| Note: In cases of the removal of one testis as the result of a service-incurred injury or disease, other than an undescended or congenitally undeveloped testis, with the absence or nonfunctioning of the other testis unrelated to service, an evaluation of 30 percent will be assigned for the service-connected testicular loss. Testis, undescended, or congenitally undeveloped is not a ratable disability. | | | |
| 7525 Epididymo-orchitis, chronic only: Rate as urinary tract infection. For tubercular infections: Rate in accordance with §§4.88b or 4.89, whichever is appropriate. | | | |
| 7527 Prostate gland injuries, infections, hypertrophy, postoperative residuals: Rate as voiding dysfunction or urinary tract infection, whichever is predominant. | | | |
| 7528 Malignant neoplasms of the genitourinary system | 100 | | |

Department of Veterans Affairs

§4.116

| | Rating |
|---|--------|
| Or rate as renal dysfunction. | |
| 7533 Cystic diseases of the kidneys (polycystic disease, uremic medullary cystic disease, Medullary sponge kidney, and similar conditions): | |
| Rate as renal dysfunction. | |
| 7534 Atherosclerotic renal disease (renal artery stenosis or atheroembolic renal disease): | |
| Rate as renal dysfunction. | |
| 7535 Toxic nephropathy (antibiotics, radiocontrast agents, nonsteroidal anti-inflammatory agents, heavy metals, and similar agents): | |
| Rate as renal dysfunction. | |
| 7536 Glomerulonephritis: | |
| Rate as renal dysfunction. | |
| 7537 Interstitial nephritis: | |
| Rate as renal dysfunction. | |
| 7538 Papillary necrosis: | |
| Rate as renal dysfunction. | |
| 7539 Renal amyloid disease: | |
| Rate as renal dysfunction. | |
| 7540 Disseminated intravascular coagulation with renal cortical necrosis: | |
| Rate as renal dysfunction. | |
| 7541 Renal involvement in diabetes mellitus, sickle cell anemia, systemic lupus erythematosus, vasculitis, or other systemic disease processes. | |
| Rate as renal dysfunction. | |
| 7542 Neurogenic bladder: | |
| Rate as voiding dysfunction. | |

¹ Review for entitlement to special monthly compensation under § 3.350 of this chapter.

[59 FR 2527, Jan. 18, 1994; 59 FR 14567, Mar. 29, 1994, as amended at 59 FR 46339, Sept. 8, 1994]

GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

§ 4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

| | Rating |
|---|--------|
| Note 1: Natural menopause, primary amenorrhea, and pregnancy and childbirth are not disabilities for rating purposes. Chronic residuals of medical or surgical complications of pregnancy may be disabilities for rating purposes. | |
| Note 2: When evaluating any claim involving loss or loss of use of one or more creative organs, refer to § 3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, almost any condition in this section might, under certain circumstances, establish entitlement to special monthly compensation. | |
| 7610 Vulva, disease or injury of (including vulvovaginitis). | |
| 7611 Vagina, disease or injury of. | |
| 7612 Cervix, disease or injury of. | |
| 7613 Uterus, disease, injury, or adhesions of. | |
| 7614 Fallopian tube, disease, injury, or adhesions of (including pelvic inflammatory disease (PID)). | |
| 7615 Ovary, disease, injury, or adhesions of. | |
| General Rating Formula for Disease, Injury, or Adhesions of Female Reproductive Organs (diagnostic codes 7610 through 7615): | |
| Symptoms not controlled by continuous treatment | 30 |
| Symptoms that require continuous treatment | 10 |
| Symptoms that do not require continuous treatment | 0 |
| 7617 Uterus and both ovaries, removal of, complete: | |
| For three months after removal | 100 |
| Thereafter | 50 |
| 7618 Uterus, removal of, including corpus: | |
| For three months after removal | 100 |
| Thereafter | 30 |
| 7619 Ovary, removal of: | |
| For three months after removal | 100 |
| Thereafter: | |
| Complete removal of both ovaries | 30 |
| Removal of one with or without partial removal of the other | 10 |
| 7620 Ovaries, atrophy of both, complete | 120 |
| 7621 Uterus, prolapse: | |
| Complete, through vagina and introitus | 50 |
| Incomplete | 30 |
| 7622 Uterus, displacement of: | |
| With marked displacement and frequent or continuous menstrual disturbances | 30 |
| With adhesions and irregular menstruation | 10 |
| 7623 Pregnancy, surgical complications of: | |
| With rectocele or cystocele | 50 |
| With relaxation of perineum | 10 |
| 7624 Fistula, rectovaginal: | |
| Vaginal fecal leakage at least once a day requiring wearing of pad | 100 |
| Vaginal fecal leakage four or more times per week, but less than daily, requiring wearing of pad | 60 |
| Vaginal fecal leakage one to three times per week requiring wearing of pad | 30 |
| Vaginal fecal leakage less than once a week ... | 10 |

| | Rating |
|--|--------|
| Without leakage | 0 |
| 7625 Fistula, urethrovaginal: | |
| Multiple urethrovaginal fistulae | 100 |
| Requiring the use of an appliance or the wearing of absorbent materials which must be changed more than four times per day | 60 |
| Requiring the wearing of absorbent materials which must be changed two to four times per day | 40 |
| Requiring the wearing of absorbent materials which must be changed less than two times per day | 20 |
| 7626 Breast, surgery of: | |
| Following radical mastectomy: | |
| Both | 80 |
| One | 50 |
| Following modified radical mastectomy: | |
| Both | 60 |
| One | 40 |
| Following simple mastectomy or wide local excision with significant alteration of size or form: | |
| Both | 50 |
| One | 30 |
| Following wide local excision without significant alteration of size or form: | |
| Both or one | 0 |
| Note: For VA purposes: | |
| (1) <i>Radical mastectomy</i> means removal of the entire breast, underlying pectoral muscles, and regional lymph nodes up to the coracoclavicular ligament. | |
| (2) <i>Modified radical mastectomy</i> means removal of the entire breast and axillary lymph nodes (in continuity with the breast). Pectoral muscles are left intact. | |
| (3) <i>Simple (or total) mastectomy</i> means removal of all of the breast tissue, nipple, and a small portion of the overlying skin, but lymph nodes and muscles are left intact. | |
| (4) <i>Wide local excision</i> (including partial mastectomy, lumpectomy, tylectomy, segmentectomy, and quadrantectomy) means removal of a portion of the breast tissue. | |
| 7627 Malignant neoplasms of gynecological system or breast | 100 |
| Note: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | |
| 7628 Benign neoplasms of the gynecological system or breast. Rate according to impairment in function of the urinary or gynecological systems, or skin. | |
| 7629 Endometriosis: | |
| Lesions involving bowel or bladder confirmed by laparoscopy, pelvic pain or heavy or irregular bleeding not controlled by treatment, and bowel or bladder symptoms | 50 |
| Pelvic pain or heavy or irregular bleeding not controlled by treatment | 30 |
| Pelvic pain or heavy or irregular bleeding requiring continuous treatment for control | 10 |

| | Rating |
|---|--------|
| Note: Diagnosis of endometriosis must be substantiated by laparoscopy. | |
| ¹ Review for entitlement to special monthly compensation under § 3.350 of this chapter. | |
| [60 FR 19855, Apr. 21, 1995] | |
| THE HEMIC AND LYMPHATIC SYSTEMS | |
| § 4.117 Schedule of ratings—hemic and lymphatic systems. | |
| | Rating |
| 7700 Anemia, hypochromic-microcytic and megaloblastic, such as iron-deficiency and pernicious anemia: | |
| Hemoglobin 5gm/100ml or less, with findings such as high output congestive heart failure or dyspnea at rest | 100 |
| Hemoglobin 7gm/100ml or less, with findings such as dyspnea on mild exertion, cardiomegaly, tachycardia (100 to 120 beats per minute) or syncope (three episodes in the last six months) | 70 |
| Hemoglobin 8gm/100ml or less, with findings such as weakness, easy fatigability, headaches, lightheadedness, or shortness of breath | 30 |
| Hemoglobin 10gm/100ml or less with findings such as weakness, easy fatigability or headaches | 10 |
| Hemoglobin 10gm/100ml or less, asymptomatic | 0 |
| NOTE: Evaluate complications of pernicious anemia, such as dementia or peripheral neuropathy, separately. | |
| 7702 Agranulocytosis, acute: | |
| Requiring bone marrow transplant, or; requiring transfusion of platelets or red cells at least once every six weeks, or; infections recurring at least once every six weeks | 100 |
| Requiring transfusion of platelets or red cells at least once every three months, or; infections recurring at least once every three months ... | 60 |
| Requiring transfusion of platelets or red cells at least once per year but less than once every three months, or; infections recurring at least once per year but less than once every three months | 30 |
| Requiring continuous medication for control | 10 |
| NOTE: The 100 percent rating for bone marrow transplant shall be assigned as of the date of hospital admission and shall continue with a mandatory VA examination six months following hospital discharge. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. | |
| 7703 Leukemia: | |
| With active disease or during a treatment phase | 100 |
| Otherwise rate as anemia (code 7700) or aplastic anemia (code 7716), whichever would result in the greater benefit. | |
| NOTE: The 100 percent rating shall continue beyond the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no recurrence, rate on residuals. | |

Department of Veterans Affairs

§4.118

| | Rating |
|---|--------|
| 7704 Polycythemia vera: During periods of treatment with myelosuppressants and for three months following cessation of myelosuppressant therapy | 100 |
| Requiring phlebotomy | 40 |
| Stable, with or without continuous medication ... | 10 |
| NOTE: Rate complications such as hypertension, gout, stroke or thrombotic disease separately. | |
| 7705 Thrombocytopenia, primary, idiopathic or immune: Platelet count of less than 20,000, with active bleeding, requiring treatment with medication and transfusions | 100 |
| Platelet count between 20,000 and 70,000, not requiring treatment, without bleeding | 70 |
| Stable platelet count between 70,000 and 100,000, without bleeding | 30 |
| Stable platelet count of 100,000 or more, without bleeding | 0 |
| 7706 Splenectomy | 20 |
| NOTE: Rate complications such as systemic infections with encapsulated bacteria separately. | |
| 7707 Spleen, injury of, healed. Rate for any residuals. | |
| 7709 Hodgkin's disease: With active disease or during a treatment phase | 100 |
| NOTE: The 100 percent rating shall continue beyond the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | |
| 7710 Adenitis, tuberculous, active or inactive. Rate under §§ 4.88c or 4.89 of this part, whichever is appropriate. | |
| 7714 Sickle cell anemia: With repeated painful crises, occurring in skin, joints, bones or any major organs caused by hemolysis and sickling of red blood cells, with anemia, thrombosis and infarction, with symptoms precluding even light manual labor | 100 |
| With painful crises several times a year or with symptoms precluding other than light manual labor | 60 |
| Following repeated hemolytic sickling crises with continuing impairment of health | 30 |
| Asymptomatic, established case in remission, but with identifiable organ impairment | 10 |
| NOTE: Sickle cell trait alone, without a history of directly attributable pathological findings, is not a ratable disability. Cases of symptomatic sickle cell trait will be forwarded to the Director, Compensation and Pension Service, for consideration under §3.321(b)(1) of this chapter. | |
| 7715 Non-Hodgkin's lymphoma: With active disease or during a treatment phase | 100 |
| NOTE: The 100 percent rating shall continue beyond the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | |

| | Rating |
|--|--------|
| 7716 Aplastic anemia: Requiring bone marrow transplant, or; requiring transfusion of platelets or red cells at least once every six weeks, or; infections recurring at least once every six weeks | 100 |
| Requiring transfusion of platelets or red cells at least once every three months, or; infections recurring at least once every three months ... | 60 |
| Requiring transfusion of platelets or red cells at least once per year but less than once every three months, or; infections recurring at least once per year but less than once every three months | 30 |
| Requiring continuous medication for control | 10 |
| NOTE: The 100 percent rating for bone marrow transplant shall be assigned as of the date of hospital admission and shall continue with a mandatory VA examination six months following hospital discharge. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. | |

[60 FR 49227, Sept. 22, 1995]

THE SKIN

§4.118 Schedule of ratings—skin.

| | Rating |
|---|--------|
| 7800 Scars, disfiguring, head, face or neck: Complete or exceptionally repugnant deformity of one side of face or marked or repugnant bilateral disfigurement | 50 |
| Severe, especially if producing a marked and unsightly deformity of eyelids, lips, or auricles | 30 |
| Moderate; disfiguring | 10 |
| Slight | 0 |
| NOTE: When in addition to tissue loss and cicatrization there is marked discoloration, color contrast, or the like, the 50 percent rating under Code 7800 may be increased to 80 percent, the 30 percent to 50 percent, and the 10 percent to 30 percent. The most repugnant, disfiguring conditions, including scars and diseases of the skin, may be submitted for central office rating, with several unretouched photographs. | |
| 7801 Scars, burns, third degree: Area or areas exceeding 1 square foot (0.1 m. ²) | 40 |
| Area or areas exceeding one-half square foot (0.05 m. ²) | 30 |
| Area or areas exceeding 12 square inches (77.4 cm. ²) | 20 |
| Area or areas exceeding 6 square inches (38.7 cm. ²) | 10 |
| NOTE (1): Actual third degree residual involvement required to the extent shown under 7801. | |
| NOTE (2): Ratings for widely separated areas, as on two or more extremities or on anterior and posterior surfaces of extremities or trunk, will be separately rated and combined. | |
| 7802 Scars, burns, second degree: Area or areas approximating 1 square foot (0.1 m. ²) | 10 |
| NOTE: See NOTE (2) under diagnostic code 7801. | |
| 7803 Scars, superficial, poorly nourished, with repeated ulceration | 10 |

§4.119

38 CFR Ch. I (7-1-01 Edition)

| | Rat- ing |
|--|-------------|
| 7804 Scars, superficial, tender and painful on objective demonstration | 10 |
| NOTE: The 10 percent rating will be assigned, when the requirements are met, even though the location may be on tip of finger or toe, and the rating may exceed the amputation value for the limited involvement. | |
| 7805 Scars, other. Rate on limitation of function of part affected. | |
| 7806 Eczema: With ulceration or extensive exfoliation or crusting, and systemic or nervous manifestations, or exceptionally repugnant | 50 |
| With exudation or itching constant, extensive lesions, or marked disfigurement | 30 |
| With exfoliation, exudation or itching, if involving an exposed surface or extensive area | 10 |
| With slight, if any, exfoliation, exudation or itching, if on a nonexposed surface or small area | 0 |
| 7807 Leishmaniasis, americana (mucocutaneous, espundia). | |
| 7808 Leishmaniasis, old world (cutaneous, oriental sore). | |
| 7809 Lupus erythematosus, discoid. (Not to be combined with ratings under diagnostic code 6350.) | |
| 7810 Pinta. | |
| 7811 Tuberculosis luposa (lupus vulgaris), active or inactive: Active | 100 |
| Inactive: See §§ 4.88b and 4.89. | |
| 7812 Verruga peruana. | |
| 7813 Dermatophytosis. | |
| 7814 Tinea barbae. | |
| 7815 Pemphigus. | |
| 7816 Psoriasis. | |
| 7817 Dermatitis exfoliativa. | |
| 7818 New growths, malignant, skin. Rate scars, disfigurement, etc., on the extent of constitutional symptoms, physical impairment. | |
| 7819 New growths, benign, skin. Rate as scars, disfigurement, etc. Unless otherwise provided, rate codes 7807 through 7819 as for eczema, dependent upon location, extent, and repugnant or otherwise disabling character of manifestations. NOTE: The most repugnant conditions may be submitted for central office rating with several unretouched photographs. Total disability ratings may be assigned without reference to Central Office in the most severe cases of pemphigus and dermatitis exfoliativa with constitutional symptoms. | |

| | Rat- ing |
|---|-------------|
| Thyroid enlargement, tachycardia (more than 100 beats per minute), eye involvement, muscular weakness, loss of weight, and sympathetic nervous system, cardiovascular, or astrointestinal symptoms | 100 |
| Emotional instability, tachycardia, fatigability, and increased pulse pressure or blood pressure | 60 |
| Tachycardia, tremor, and increased pulse pressure or blood pressure | 30 |
| Tachycardia, which may be intermittent, and tremor, or; continuous medication required for control | 10 |
| NOTE (1): If disease of the heart is the predominant finding, evaluate as hyperthyroid heart disease (DC 7008) if doing so would result in a higher evaluation than using the criteria above. | |
| NOTE (2): If ophthalmopathy is the sole finding, evaluate as field vision, impairment of (DC 6080); diplopia (DC 6090); or impairment of central visual acuity (DC 6061-6079). | |
| 7901 Thyroid gland, toxic adenoma of Thyroid enlargement, tachycardia (more than 100 beats per minute), eye involvement, muscular weakness, loss of weight, and sympathetic nervous system, cardiovascular, or gastrointestinal symptoms | 100 |
| Emotional instability, tachycardia, fatigability, and increased pulse pressure or blood pressure | 60 |
| Tachycardia, tremor, and increased pulse pressure or blood pressure | 30 |
| Tachycardia, which may be intermittent, and tremor, or; continuous medication required for control | 10 |
| NOTE (1): If disease of the heart is the predominant finding, evaluate as hyperthyroid heart disease (DC 7008) if doing so would result in a higher evaluation than using the criteria above. | |
| NOTE (2): If ophthalmopathy is the sole finding, evaluate as field vision, impairment of (DC 6080); diplopia (DC 6090); or impairment of central visual acuity (DC 6061-6079). | |
| 7902 Thyroid gland, nontoxic adenoma of With disfigurement of the head or neck | 20 |
| Without disfigurement of the head or neck | 0 |
| NOTE: If there are symptoms due to pressure on adjacent organs such as the trachea, larynx, or esophagus, evaluate under the diagnostic code for disability of that organ, if doing so would result in a higher evaluation than using this diagnostic code. | |
| 7903 Hypothyroidism Cold intolerance, muscular weakness, cardiovascular involvement, mental disturbance (dementia, slowing of thought, depression), bradycardia (less than 60 beats per minute), and sleepiness ... | 100 |
| Muscular weakness, mental disturbance, and weight gain | 60 |
| Fatigability, constipation, and mental sluggishness .. | 30 |
| Fatigability, or; continuous medication required for control | 10 |
| 7904 Hyperparathyroidism Generalized decalcification of bones, kidney stones, gastrointestinal symptoms (nausea, vomiting, anorexia, constipation, weight loss, or peptic ulcer), and weakness | 100 |
| Gastrointestinal symptoms and weakness | 60 |
| Continuous medication required for control | 10 |
| NOTE: Following surgery or treatment, evaluate as digestive, skeletal, renal, or cardiovascular residuals or as endocrine dysfunction. | |
| 7905 Hypoparathyroidism | |

[29 FR 6718, May 22, 1964, as amended at 34 FR 5063, Mar. 11, 1969; 40 FR 42540, Sept. 15, 1975; 43 FR 45262, Oct. 2, 1978]

THE ENDOCRINE SYSTEM

§4.119 Schedule of ratings—endocrine system.

| | Rat- ing |
|----------------------|-------------|
| 7900 Hyperthyroidism | |

Department of Veterans Affairs

§4.119

| | Rat- ing | | Rat- ing |
|---|-------------|---|-------------|
| Marked neuromuscular excitability (such as convulsions, muscular spasms (tetany), or laryngeal stridor) plus either cataract or evidence of increased intracranial pressure (such as papilledema) | 100 | NOTE (1): An Addisonian "crisis" consists of the rapid onset of peripheral vascular collapse (with acute hypotension and shock), with findings that may include: anorexia; nausea; vomiting; dehydration; profound weakness; pain in abdomen, legs, and back; fever; apathy, and depressed mentation with possible progression to coma, renal shutdown, and death. | |
| Marked neuromuscular excitability, or; paresthesias (of arms, legs, or circumoral area) plus either cataract or evidence of increased intracranial pressure | 60 | NOTE (2): An Addisonian "episode," for VA purposes, is a less acute and less severe event than an Addisonian crisis and may consist of anorexia, nausea, vomiting, diarrhea, dehydration, weakness, malaise, orthostatic hypotension, or hypoglycemia, but no peripheral vascular collapse. | |
| Continuous medication required for control | 10 | NOTE (3): Tuberculous Addison's disease will be evaluated as active or inactive tuberculosis. If inactive, these evaluations are not to be combined with the graduated ratings of 50 percent or 30 percent for non-pulmonary tuberculosis specified under §4.88b. Assign the higher rating. | |
| 7907 Cushing's syndrome | | 7912 Pluriglandular syndrome | |
| As active, progressive disease including loss of muscle strength, areas of osteoporosis, hypertension, weakness, and enlargement of pituitary or adrenal gland | 100 | Evaluate according to major manifestations. | |
| Loss of muscle strength and enlargement of pituitary or adrenal gland | 60 | 7913 Diabetes mellitus | |
| With striae, obesity, moon face, glucose intolerance, and vascular fragility | 30 | Requiring more than one daily injection of insulin, restricted diet, and regulation of activities (avoidance of strenuous occupational and recreational activities) with episodes of ketoacidosis or hypoglycemic reactions requiring at least three hospitalizations per year or weekly visits to a diabetic care provider, plus either progressive loss of weight and strength or complications that would be compensable if separately evaluated | 100 |
| NOTE: With recovery or control, evaluate as residuals of adrenal insufficiency or cardiovascular, psychiatric, skin, or skeletal complications under appropriate diagnostic code. | | Requiring insulin, restricted diet, and regulation of activities with episodes of ketoacidosis or hypoglycemic reactions requiring one or two hospitalizations per year or twice a month visits to a diabetic care provider, plus complications that would not be compensable if separately evaluated | 60 |
| 7908 Acromegaly | | Requiring insulin, restricted diet, and regulation of activities | 40 |
| Evidence of increased intracranial pressure (such as visual field defect), arthropathy, glucose intolerance, and either hypertension or cardiomegaly | 100 | Requiring insulin and restricted diet, or; oral hypoglycemic agent and restricted diet | 20 |
| Arthropathy, glucose intolerance, and hypertension | 60 | Manageable by restricted diet only | 10 |
| Enlargement of acral parts or overgrowth of long bones, and enlarged sella turcica | 30 | NOTE (1): Evaluate compensable complications of diabetes separately unless they are part of the criteria used to support a 100 percent evaluation. Noncompensable complications are considered part of the diabetic process under diagnostic code 7913. | |
| 7909 Diabetes insipidus | | NOTE (2): When diabetes mellitus has been conclusively diagnosed, do not request a glucose tolerance test solely for rating purposes. | |
| Polyuria with near-continuous thirst, and more than two documented episodes of dehydration requiring parenteral hydration in the past year | 100 | 7914 Neoplasm, malignant, any specified part of the endocrine system | 100 |
| Polyuria with near-continuous thirst, and one or two documented episodes of dehydration requiring parenteral hydration in the past year | 60 | NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | |
| Polyuria with near-continuous thirst, and one or more episodes of dehydration in the past year not requiring parenteral hydration | 40 | 7915 Neoplasm, benign, any specified part of the endocrine system rate as residuals of endocrine dysfunction. | |
| Polyuria with near-continuous thirst | 20 | 7916 Hyperpituitarism (prolactin secreting pituitary dysfunction) | |
| 7911 Addison's disease (Adrenal Cortical Hypofunction) | | 7917 Hyperaldosteronism (benign or malignant) | |
| Four or more crises during the past year | 60 | | |
| Three crises during the past year, or; five or more episodes during the past year | 40 | | |
| One or two crises during the past year, or; two to four episodes during the past year, or; weakness and fatigability, or; corticosteroid therapy required for control | 20 | | |

§ 4.120

| | Rat- ing |
|--|-------------|
| 7918 Pheochromocytoma (benign or malignant) NOTE: Evaluate diagnostic codes 7916, 7917, and 7918 as malignant or benign neoplasm as appropriate. | 100 |
| 7919 C-cell hyperplasia of the thyroid NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals. | |

[61 FR 20446, May 7, 1996]

NEUROLOGICAL CONDITIONS AND
CONVULSIVE DISORDERS

§ 4.120 Evaluations by comparison.

Disability in this field is ordinarily to be rated in proportion to the impairment of motor, sensory or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, injury to the skull, etc. In rating disability from the conditions in the preceding sentence refer to the appropriate schedule. In rating peripheral nerve injuries and their residuals, attention should be given to the site and character of the injury, the relative impairment in motor function, trophic changes, or sensory disturbances.

§ 4.121 Identification of epilepsy.

When there is doubt as to the true nature of epileptiform attacks, neurological observation in a hospital adequate to make such a study is necessary. To warrant a rating for epilepsy, the seizures must be witnessed or verified at some time by a physician. As to frequency, competent, consistent lay testimony emphasizing convulsive and immediate post-convulsive characteristics may be accepted. The frequency of seizures should be ascertained under the ordinary conditions of life (while not hospitalized).

§ 4.122 Psychomotor epilepsy.

The term psychomotor epilepsy refers to a condition that is character-

ized by seizures and not uncommonly by a chronic psychiatric disturbance as well.

(a) Psychomotor seizures consist of episodic alterations in conscious control that may be associated with automatic states, generalized convulsions, random motor movements (chewing, lip smacking, fumbling), hallucinatory phenomena (involving taste, smell, sound, vision), perceptual illusions (deja vu, feelings of loneliness, strangeness, macropsia, micropsia, dreamy states), alterations in thinking (not open to reason), alterations in memory, abnormalities of mood or affect (fear, alarm, terror, anger, dread, well-being), and autonomic disturbances (sweating, pallor, flushing of the face, visceral phenomena such as nausea, vomiting, defecation, a rising feeling of warmth in the abdomen). Automatic states or automatisms are characterized by episodes of irrational, irrelevant, disjointed, unconventional, asocial, purposeless though seemingly coordinated and purposeful, confused or inappropriate activity of one to several minutes (or, infrequently, hours) duration with subsequent amnesia for the seizure. Examples: A person of high social standing remained seated, muttered angrily, and rubbed the arms of his chair while the National Anthem was being played; an apparently normal person suddenly disrobed in public; a man traded an expensive automobile for an antiquated automobile in poor mechanical condition and after regaining conscious control, discovered that he had signed an agreement to pay an additional sum of money in the trade. The seizure manifestations of psychomotor epilepsy vary from patient to patient and in the same patient from seizure to seizure.

(b) A chronic mental disorder is not uncommon as an interseizure manifestation of psychomotor epilepsy and may include psychiatric disturbances extending from minimal anxiety to severe personality disorder (as distinguished from developmental) or almost complete personality disintegration (psychosis). The manifestations of a chronic mental disorder associated with psychomotor epilepsy, like those of the seizures, are protean in character.

Department of Veterans Affairs

§ 4.124a

§ 4.123 Neuritis, cranial or peripheral.

Neuritis, cranial or peripheral, characterized by loss of reflexes, muscle atrophy, sensory disturbances, and constant pain, at times excruciating, is to be rated on the scale provided for injury of the nerve involved, with a maximum equal to severe, incomplete, paralysis. See nerve involved for diagnostic code number and rating. The maximum rating which may be assigned for neuritis not characterized by organic changes referred to in this section will be that for moderate, or with sciatic nerve involvement, for moderately severe, incomplete paralysis.

§ 4.124 Neuralgia, cranial or peripheral.

Neuralgia, cranial or peripheral, characterized usually by a dull and intermittent pain, of typical distribution so as to identify the nerve, is to be rated on the same scale, with a maximum equal to moderate incomplete paralysis. See nerve involved for diagnostic code number and rating. Tic douloureux, or trifacial neuralgia, may be rated up to complete paralysis of the affected nerve.

§ 4.124a Schedule of ratings—neurological conditions and convulsive disorders.

[With the exceptions noted, disability from the following diseases and their residuals may be rated from 10 percent to 100 percent in proportion to the impairment of motor, sensory, or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, etc., referring to the appropriate bodily system of the schedule. With partial loss of use of one or more extremities from neurological lesions, rate by comparison with the mild, moderate, severe, or complete paralysis of peripheral nerves]

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM

| | Rating |
|---------------------------------------|--------|
| 8000 Encephalitis, epidemic, chronic: | |
| As active febrile disease | 100 |
| Rate residuals, minimum | 10 |
| Brain, new growth of: | |
| 8002 Malignant | 100 |

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

| | Rating |
|--|--------|
| NOTE: The rating in code 8002 will be continued for 2 years following cessation of surgical, chemotherapeutic or other treatment modality. At this point, if the residuals have stabilized, the rating will be made on neurological residuals according to symptomatology. | |
| Minimum rating | 30 |
| 8003 Benign, minimum | 60 |
| Rate residuals, minimum | 10 |
| 8004 Paralysis agitans: | |
| Minimum rating | 30 |
| 8005 Bulbar palsy | 100 |
| 8007 Brain, vessels, embolism of. | |
| 8008 Brain, vessels, thrombosis of. | |
| 8009 Brain, vessels, hemorrhage from: | |
| Rate the vascular conditions under Codes 8007 through 8009, for 6 months | 100 |
| Rate residuals, thereafter, minimum | 10 |
| 8010 Myelitis: | |
| Minimum rating | 10 |
| 8011 Poliomyelitis, anterior: | |
| As active febrile disease | 100 |
| Rate residuals, minimum | 10 |
| 8012 Hematomyelia: | |
| For 6 months | 100 |
| Rate residuals, minimum | 10 |
| 8013 Syphilis, cerebrospinal. | |
| 8014 Syphilis, meningovascular. | |
| 8015 Tabes dorsalis. | |
| NOTE: Rate upon the severity of convulsions, paralysis, visual impairment or psychotic involvement, etc. | |
| 8017 Amyotrophic lateral sclerosis: | |
| Minimum rating | 30 |
| 8018 Multiple sclerosis: | |
| Minimum rating | 30 |
| 8019 Meningitis, cerebrospinal, epidemic: | |
| As active febrile disease | 100 |
| Rate residuals, minimum | 10 |
| 8020 Brain, abscess of: | |
| As active disease | 100 |
| Rate residuals, minimum | 10 |
| Spinal cord, new growths of: | |
| 8021 Malignant | 100 |
| NOTE: The rating in code 8021 will be continued for 2 years following cessation of surgical, chemotherapeutic or other treatment modality. At this point, if the residuals have stabilized, the rating will be made on neurological residuals according to symptomatology. | |
| Minimum rating | 30 |
| 8022 Benign, minimum rating | 60 |
| Rate residuals, minimum | 10 |
| 8023 Progressive muscular atrophy: | |
| Minimum rating | 30 |
| 8024 Syringomyelia: | |
| Minimum rating | 30 |
| 8025 Myasthenia gravis: | |
| Minimum rating | 30 |

§ 4.124a

38 CFR Ch. I (7-1-01 Edition)

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

| | Rat- ing |
|---|-------------|
| NOTE: It is required for the minimum ratings for residuals under diagnostic codes 8000-8025, that there be ascertainable residuals. Determinations as to the presence of residuals not capable of objective verification, i.e., headaches, dizziness, fatigability, must be approached on the basis of the diagnosis recorded; subjective residuals will be accepted when consistent with the disease and not more likely attributable to other disease or no disease. It is of exceptional importance that when ratings in excess of the prescribed minimum ratings are assigned, the diagnostic codes utilized as bases of evaluation be cited, in addition to the codes identifying the diagnoses. | |
| 8045 Brain disease due to trauma: Purely neurological disabilities, such as hemiplegia, epileptiform seizures, facial nerve paralysis, etc., following trauma to the brain, will be rated under the diagnostic codes specifically dealing with such disabilities, with citation of a hyphenated diagnostic code (e.g., 8045-8207). Purely subjective complaints such as headache, dizziness, insomnia, etc., recognized as symptomatic of brain trauma, will be rated 10 percent and no more under diagnostic code 9304. This 10 percent rating will not be combined with any other rating for a disability due to brain trauma. Ratings in excess of 10 percent for brain disease due to trauma under diagnostic code 9304 are not assignable in the absence of a diagnosis of multi-infarct dementia associated with brain trauma. | |
| 8046 Cerebral arteriosclerosis: Purely neurological disabilities, such as hemiplegia, cranial nerve paralysis, etc., due to cerebral arteriosclerosis will be rated under the diagnostic codes dealing with such specific disabilities, with citation of a hyphenated diagnostic code (e.g., 8046-8207). Purely subjective complaints such as headache, dizziness, tinnitus, insomnia and irritability, recognized as symptomatic of a properly diagnosed cerebral arteriosclerosis, will be rated 10 percent and no more under diagnostic code 9305. This 10 percent rating will not be combined with any other rating for a disability due to cerebral or generalized arteriosclerosis. Ratings in excess of 10 percent for cerebral arteriosclerosis under diagnostic code 9305 are not assignable in the absence of a diagnosis of multi-infarct dementia with cerebral arteriosclerosis. NOTE: The ratings under code 8046 apply only when the diagnosis of cerebral arteriosclerosis is substantiated by the entire clinical picture and not solely on findings of retinal arteriosclerosis. | |
| MISCELLANEOUS DISEASES | |
| | Rat- ing |
| 8100 Migraine: | |

MISCELLANEOUS DISEASES—Continued

| | Rat- ing |
|--|-------------|
| With very frequent completely prostrating and prolonged attacks productive of severe economic inadaptability | 50 |
| With characteristic prostrating attacks occurring on an average once a month over last several months | 30 |
| With characteristic prostrating attacks averaging one in 2 months over last several months | 10 |
| With less frequent attacks | 0 |
| 8103 Tic, convulsive: | |
| Severe | 30 |
| Moderate | 10 |
| Mild | 0 |
| NOTE: Depending upon frequency, severity, muscle groups involved. | |
| 8104 Paramyoclonus multiplex (convulsive state, myoclonic type): | |
| Rate as tic; convulsive; severe cases | 60 |
| 8105 Chorea, Sydenham's: | |
| Pronounced, progressive grave types | 100 |
| Severe | 80 |
| Moderately severe | 50 |
| Moderate | 30 |
| Mild | 10 |
| NOTE: Consider rheumatic etiology and complications. | |
| 8106 Chorea, Huntington's. | |
| Rate as Sydenham's chorea. This, though a familial disease, has its onset in late adult life, and is considered a ratable disability. | |
| 8107 Athetosis, acquired. | |
| Rate as chorea. | |
| 8108 Narcolepsy. | |
| Rate as for epilepsy, petit mal. | |
| DISEASES OF THE CRANIAL NERVES | |
| | Rat- ing |
| Disability from lesions of peripheral portions of first, second, third, fourth, sixth, and eighth nerves will be rated under the Organs of Special Sense. The ratings for the cranial nerves are for unilateral involvement; when bilateral, combine but without the bilateral factor. | |
| Fifth (trigeminal) cranial nerve | |
| 8205 Paralysis of: | |
| Complete | 50 |
| Incomplete, severe | 30 |
| Incomplete, moderate | 10 |
| NOTE: Dependent upon relative degree of sensory manifestation or motor loss. | |
| 8305 Neuritis. | |
| 8405 Neuralgia. | |
| NOTE: Tic douloureux may be rated in accordance with severity, up to complete paralysis. | |
| Seventh (facial) cranial nerve | |
| 8207 Paralysis of: | |
| Complete | 30 |
| Incomplete, severe | 20 |
| Incomplete, moderate | 10 |
| NOTE: Dependent upon relative loss of innervation of facial muscles. | |
| 8307 Neuritis. | |
| 8407 Neuralgia. | |
| Ninth (glossopharyngeal) cranial nerve. | |
| 8209 Paralysis of: | |
| Complete | 30 |
| Incomplete, severe | 20 |
| Incomplete, moderate | 10 |

Department of Veterans Affairs

§ 4.124a

DISEASES OF THE CRANIAL NERVES—Continued

| | Rating |
|--|--------|
| NOTE: Dependent upon relative loss of ordinary sensation in mucous membrane of the pharynx, fauces, and tonsils. | |
| 8309 Neuritis. | |
| 8409 Neuralgia. | |
| Tenth (pneumogastric, vagus) cranial nerve. | |
| 8210 Paralysis of: | |
| Complete | 50 |
| Incomplete, severe | 30 |
| Incomplete, moderate | 10 |
| NOTE: Dependent upon extent of sensory and motor loss to organs of voice, respiration, pharynx, stomach and heart. | |
| 8310 Neuritis. | |
| 8410 Neuralgia. | |
| Eleventh (spinal accessory, external branch) cranial nerve. | |
| 8211 Paralysis of: | |
| Complete | 30 |
| Incomplete, severe | 20 |
| Incomplete, moderate | 10 |
| NOTE: Dependent upon loss of motor function of sternomastoid and trapezius muscles. | |
| 8311 Neuritis. | |
| 8411 Neuralgia. | |
| Twelfth (hypoglossal) cranial nerve. | |
| 8212 Paralysis of: | |
| Complete | 50 |
| Incomplete, severe | 30 |
| Incomplete, moderate | 10 |
| NOTE: Dependent upon loss of motor function of tongue. | |
| 8312 Neuritis. | |
| 8412 Neuralgia. | |

DISEASES OF THE PERIPHERAL NERVES

| Schedule of ratings | Rating | |
|--|--------|-------|
| | Major | Minor |
| The term "incomplete paralysis," with this and other peripheral nerve injuries, indicates a degree of lost or impaired function substantially less than the type picture for complete paralysis given with each nerve, whether due to varied level of the nerve lesion or to partial regeneration. When the involvement is wholly sensory, the rating should be for the mild, or at most, the moderate degree. The ratings for the peripheral nerves are for unilateral involvement; when bilateral, combine with application of the bilateral factor. | | |
| Upper radicular group (fifth and sixth cervicals) | | |
| 8510 Paralysis of: | | |
| Complete; all shoulder and elbow movements lost or severely affected, hand and wrist movements not affected | 70 | 60 |
| Incomplete: | | |
| Severe | 50 | 40 |
| Moderate | 40 | 30 |
| Mild | 20 | 20 |

DISEASES OF THE PERIPHERAL NERVES—Continued

| Schedule of ratings | Rating | |
|---|--------|-------|
| | Major | Minor |
| 8610 Neuritis. | | |
| 8710 Neuralgia. | | |
| Middle radicular group | | |
| 8511 Paralysis of: | | |
| Complete; adduction, abduction and rotation of arm, flexion of elbow, and extension of wrist lost or severely affected | 70 | 60 |
| Incomplete: | | |
| Severe | 50 | 40 |
| Moderate | 40 | 30 |
| Mild | 20 | 20 |
| 8611 Neuritis. | | |
| 8711 Neuralgia. | | |
| Lower radicular group | | |
| 8512 Paralysis of: | | |
| Complete; all intrinsic muscles of hand, and some or all of flexors of wrist and fingers, paralyzed (substantial loss of use of hand) | 70 | 60 |
| Incomplete: | | |
| Severe | 50 | 40 |
| Moderate | 40 | 30 |
| Mild | 20 | 20 |
| 8612 Neuritis. | | |
| 8712 Neuralgia. | | |
| All radicular groups | | |
| 8513 Paralysis of: | | |
| Complete | 90 | 80 |
| Incomplete: | | |
| Severe | 70 | 60 |
| Moderate | 40 | 30 |
| Mild | 20 | 20 |
| 8613 Neuritis. | | |
| 8713 Neuralgia. | | |
| The musculospiral nerve (radial nerve) | | |
| 8514 Paralysis of: | | |
| Complete; drop of hand and fingers, wrist and fingers perpetually flexed, the thumb adducted falling within the line of the outer border of the index finger; can not extend hand at wrist, extend proximal phalanges of fingers, extend thumb, or make lateral movement of wrist; supination of hand, extension and flexion of elbow weakened, the loss of synergic motion of extensors impairs the hand grip seriously; total paralysis of the triceps occurs only as the greatest rarity | 70 | 60 |
| Incomplete: | | |
| Severe | 50 | 40 |
| Moderate | 30 | 20 |
| Mild | 20 | 20 |

§ 4.124a

38 CFR Ch. I (7-1-01 Edition)

DISEASES OF THE PERIPHERAL NERVES—
Continued

| Schedule of ratings | Rating | |
|--|--------|-------|
| | Major | Minor |
| 8614 Neuritis. | | |
| 8714 Neuralgia. | | |
| NOTE: Lesions involving only "dissociation of extensor communis digitorum" and "paralysis below the extensor communis digitorum," will not exceed the moderate rating under code 8514. | | |
| The median nerve | | |
| 8515 Paralysis of: | | |
| Complete; the hand inclined to the ulnar side, the index and middle fingers more extended than normally, considerable atrophy of the muscles of the thenar eminence, the thumb in the plane of the hand (ape hand); pronation incomplete and defective, absence of flexion of index finger and feeble flexion of middle finger, cannot make a fist, index and middle fingers remain extended; cannot flex distal phalanx of thumb, defective opposition and abduction of the thumb, at right angles to palm; flexion of wrist weakened; pain with trophic disturbances ... | 70 | 60 |
| Incomplete: | | |
| Severe | 50 | 40 |
| Moderate | 30 | 20 |
| Mild | 10 | 10 |
| 8615 Neuritis. | | |
| 8715 Neuralgia. | | |
| The ulnar nerve | | |
| 8516 Paralysis of: | | |
| Complete; the "griffin claw" deformity, due to flexor contraction of ring and little fingers, atrophy very marked in dorsal interspace and thenar and hypothenar eminences; loss of extension of ring and little fingers cannot spread the fingers (or reverse), cannot adduct the thumb; flexion of wrist weakened | 60 | 50 |
| Incomplete: | | |
| Severe | 40 | 30 |
| Moderate | 30 | 20 |
| Mild | 10 | 10 |
| 8616 Neuritis. | | |
| 8716 Neuralgia. | | |
| Musculocutaneous nerve | | |
| 8517 Paralysis of: | | |
| Complete; weakness but not loss of flexion of elbow and supination of forearm | 30 | 20 |
| Incomplete: | | |
| Severe | 20 | 20 |
| Moderate | 10 | 10 |
| Mild | 0 | 0 |
| 8617 Neuritis. | | |
| 8717 Neuralgia. | | |
| Circumflex nerve | | |
| 8518 Paralysis of: | | |
| Complete; abduction of arm is impossible, outward rotation is weakened; muscles supplied are deltoid and teres minor | 50 | 40 |
| Incomplete: | | |
| Severe | 30 | 20 |
| Moderate | 10 | 10 |

DISEASES OF THE PERIPHERAL NERVES—
Continued

| Schedule of ratings | Rating | |
|---|--------|--------|
| | Major | Minor |
| Mild | 0 | 0 |
| 8618 Neuritis. | | |
| 8718 Neuralgia. | | |
| Long thoracic nerve | | |
| 8519 Paralysis of: | | |
| Complete; inability to raise arm above shoulder level, winged scapula deformity | 30 | 20 |
| Incomplete: | | |
| Severe | 20 | 20 |
| Moderate | 10 | 10 |
| Mild | 0 | 0 |
| NOTE: Not to be combined with lost motion above shoulder level. | | |
| 8619 Neuritis. | | |
| 8719 Neuralgia. | | |
| NOTE: Combined nerve injuries should be rated by reference to the major involvement, or if sufficient in extent, consider radicular group ratings. | | |
| | | Rating |
| Sciatic nerve | | |
| 8520 Paralysis of: | | |
| Complete; the foot dangles and drops, no active movement possible of muscles below the knee, flexion of knee weakened or (very rarely) lost | | 80 |
| Incomplete: | | |
| Severe, with marked muscular atrophy | | 60 |
| Moderately severe | | 40 |
| Moderate | | 20 |
| Mild | | 10 |
| 8620 Neuritis. | | |
| 8720 Neuralgia. | | |
| External popliteal nerve (common peroneal) | | |
| 8521 Paralysis of: | | |
| Complete; foot drop and slight droop of first phalanges of all toes, cannot dorsiflex the foot, extension (dorsal flexion) of proximal phalanges of toes lost; abduction of foot lost, adduction weakened; anesthesia covers entire dorsum of foot and toes | | 40 |
| Incomplete: | | |
| Severe | | 30 |
| Moderate | | 20 |
| Mild | | 10 |
| 8621 Neuritis. | | |
| 8721 Neuralgia. | | |
| Musculocutaneous nerve (superficial peroneal) | | |
| 8522 Paralysis of: | | |
| Complete; eversion of foot weakened | | 30 |
| Incomplete: | | |
| Severe | | 20 |
| Moderate | | 10 |
| Mild | | 0 |

Department of Veterans Affairs

§ 4.124a

| | Rating |
|--|--------|
| 8622 Neuritis. | |
| 8722 Neuralgia. | |
| Anterior tibial nerve (deep peroneal) | |
| 8523 Paralysis of: | |
| Complete; dorsal flexion of foot lost | 30 |
| Incomplete: | |
| Severe | 20 |
| Moderate | 10 |
| Mild | 0 |
| 8623 Neuritis. | |
| 8723 Neuralgia. | |
| Internal popliteal nerve (tibial) | |
| 8524 Paralysis of: | |
| Complete; plantar flexion lost, frank adduction of foot impossible, flexion and separation of toes abolished; no muscle in sole can move; in lesions of the nerve high in popliteal fossa, plantar flexion of foot is lost | 40 |
| Incomplete: | |
| Severe | 30 |
| Moderate | 20 |
| Mild | 10 |
| 8624 Neuritis. | |
| 8724 Neuralgia. | |
| Posterior tibial nerve | |
| 8525 Paralysis of: | |
| Complete; paralysis of all muscles of sole of foot, frequently with painful paralysis of a causalgic nature; toes cannot be flexed; adduction is weakened; plantar flexion is impaired | 30 |
| Incomplete: | |
| Severe | 20 |
| Moderate | 10 |
| Mild | 10 |
| 8625 Neuritis. | |
| 8725 Neuralgia. | |
| Anterior crural nerve (femoral) | |
| 8526 Paralysis of: | |
| Complete; paralysis of quadriceps extensor muscles | 40 |
| Incomplete: | |
| Severe | 30 |
| Moderate | 20 |
| Mild | 10 |
| 8626 Neuritis. | |
| 8726 Neuralgia. | |
| Internal saphenous nerve | |
| 8527 Paralysis of: | |
| Severe to complete | 10 |
| Mild to moderate | 0 |
| 8627 Neuritis. | |
| 8727 Neuralgia. | |
| Obturator nerve | |
| 8528 Paralysis of: | |
| Severe to complete | 10 |
| Mild or moderate | 0 |

| | Rating |
|---|--------|
| 8628 Neuritis. | |
| 8728 Neuralgia. | |
| External cutaneous nerve of thigh | |
| 8529 Paralysis of: | |
| Severe to complete | 10 |
| Mild or moderate | 0 |
| 8629 Neuritis. | |
| 8729 Neuralgia. | |
| Ilio-inguinal nerve | |
| 8530 Paralysis of: | |
| Severe to complete | 10 |
| Mild or moderate | 0 |
| 8630 Neuritis. | |
| 8730 Neuralgia. | |
| 8540 Soft-tissue sarcoma (of neurogenic origin) | 100 |
| NOTE: The 100 percent rating will be continued for 6 months following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. At this point, if there has been no local recurrence or metastases, the rating will be made on residuals. | |

THE EPILEPSIES

| | Rating |
|---|--------|
| A thorough study of all material in §§ 4.121 and 4.122 of the preface and under the ratings for epilepsy is necessary prior to any rating action. | |
| 8910 Epilepsy, grand mal. Rate under the general rating formula for major seizures. | 30 |
| 8911 Epilepsy, petit mal. Rate under the general rating formula for minor seizures. | 10 |
| NOTE (1): A major seizure is characterized by the generalized tonic-clonic convulsion with unconsciousness. | |
| NOTE (2): A minor seizure consists of a brief interruption in consciousness or conscious control associated with staring or rhythmic blinking of the eyes or nodding of the head ("pure" petit mal), or sudden jerking movements of the arms, trunk, or head (myoclonic type) or sudden loss of postural control (akinetic type). | |
| General Rating Formula for Major and Minor Epileptic Seizures: | |
| Averaging at least 1 major seizure per month over the last year | 100 |
| Averaging at least 1 major seizure in 3 months over the last year; or more than 10 minor seizures weekly | 80 |
| Averaging at least 1 major seizure in 4 months over the last year; or 9-10 minor seizures per week | 60 |
| At least 1 major seizure in the last 6 months or 2 in the last year; or averaging at least 5 to 8 minor seizures weekly | 40 |
| At least 1 major seizure in the last 2 years; or at least 2 minor seizures in the last 6 months | 20 |
| A confirmed diagnosis of epilepsy with a history of seizures | 10 |

§ 4.125

38 CFR Ch. I (7-1-01 Edition)

THE EPILEPSIES—Continued

MENTAL DISORDERS

| | Rating |
|---|--------|
| NOTE (1): When continuous medication is shown necessary for the control of epilepsy, the minimum evaluation will be 10 percent. This rating will not be combined with any other rating for epilepsy. | |
| NOTE (2): In the presence of major and minor seizures, rate the predominating type. | |
| NOTE (3): There will be no distinction between diurnal and nocturnal major seizures. | |
| 8912 Epilepsy, Jacksonian and focal motor or sensory. | |
| 8913 Epilepsy, diencephalic. Rate as minor seizures, except in the presence of major and minor seizures, rate the predominating type. | |
| 8914 Epilepsy, psychomotor. Major seizures: Psychomotor seizures will be rated as major seizures under the general rating formula when characterized by automatic states and/or generalized convulsions with unconsciousness. Minor seizures: Psychomotor seizures will be rated as minor seizures under the general rating formula when characterized by brief transient episodes of random motor movements, hallucinations, perceptual illusions, abnormalities of thinking, memory or mood, or autonomic disturbances. | |

Mental Disorders in Epilepsies: A nonpsychotic organic brain syndrome will be rated separately under the appropriate diagnostic code (e.g., 9304 or 9307). In the absence of a diagnosis of non-psychotic organic psychiatric disturbance (psychotic, psychoneurotic or personality disorder) if diagnosed and shown to be secondary to or directly associated with epilepsy will be rated separately. The psychotic or psychoneurotic disorder will be rated under the appropriate diagnostic code. The personality disorder will be rated as a dementia (e.g., diagnostic code 9304 or 9307).

Epilepsy and Unemployability: (1) Rating specialists must bear in mind that the epileptic, although his or her seizures are controlled, may find employment and rehabilitation difficult of attainment due to employer reluctance to the hiring of the epileptic.

(2) Where a case is encountered with a definite history of unemployment, full and complete development should be undertaken to ascertain whether the epilepsy is the determining factor in his or her inability to obtain employment.

(3) The assent of the claimant should first be obtained for permission to conduct this economic and social survey. The purpose of this survey is to secure all the relevant facts and data necessary to permit a true judgment as to the reason for his or her unemployment and should include information as to:

- (a) Education;
- (b) Occupations prior and subsequent to service;
- (c) Places of employment and reasons for termination;
- (d) Wages received;
- (e) Number of seizures.

(4) Upon completion of this survey and current examination, the case should have rating board consideration. Where in the judgment of the rating board the veteran's unemployability is due to epilepsy and jurisdiction is not vested in that body by reason of schedular evaluations, the case should be submitted to the Director, Compensation and Pension Service.

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 40 FR 42540, Sept. 15, 1975; 41 FR 11302, Mar. 18, 1976; 43 FR 45362, Oct. 2, 1978; 54 FR 4282, Jan. 30, 1989; 54 FR 49755, Dec. 1, 1989; 55 FR 154, Jan. 3, 1990; 56 FR 51653, Oct. 15, 1991; 57 FR 24364, June 9, 1992]

§ 4.125 Diagnosis of mental disorders.

(a) If the diagnosis of a mental disorder does not conform to DSM-IV or is not supported by the findings on the examination report, the rating agency shall return the report to the examiner to substantiate the diagnosis.

(b) If the diagnosis of a mental disorder is changed, the rating agency shall determine whether the new diagnosis represents progression of the prior diagnosis, correction of an error in the prior diagnosis, or development of a new and separate condition. If it is not clear from the available records what the change of diagnosis represents, the rating agency shall return the report to the examiner for a determination.

(Authority: 38 U.S.C. 1155)

[61 FR 52700, Oct. 8, 1996]

§ 4.126 Evaluation of disability from mental disorders.

(a) When evaluating a mental disorder, the rating agency shall consider the frequency, severity, and duration of psychiatric symptoms, the length of remissions, and the veteran's capacity for adjustment during periods of remission. The rating agency shall assign an evaluation based on all the evidence of record that bears on occupational and social impairment rather than solely on the examiner's assessment of the level of disability at the moment of the examination.

(b) When evaluating the level of disability from a mental disorder, the rating agency will consider the extent of social impairment, but shall not assign an evaluation solely on the basis of social impairment.

(c) Delirium, dementia, and amnesic and other cognitive disorders shall be evaluated under the general rating formula for mental disorders; neurologic deficits or other impairments stemming from the same etiology (e.g., a head injury) shall be evaluated separately and combined with the evaluation for delirium, dementia, or amnesic or other cognitive disorder (see § 4.25).

(d) When a single disability has been diagnosed both as a physical condition

Department of Veterans Affairs

§ 4.130

and as a mental disorder, the rating agency shall evaluate it using a diagnostic code which represents the dominant (more disabling) aspect of the condition (see § 4.14).

(Authority: 38 U.S.C. 1155)
[61 FR 52700, Oct. 8, 1996]

§ 4.127 Mental retardation and personality disorders.

Mental retardation and personality disorders are not diseases or injuries for compensation purposes, and, except as provided in § 3.310(a) of this chapter, disability resulting from them may not be service-connected. However, disability resulting from a mental disorder that is superimposed upon mental retardation or a personality disorder may be service-connected.

(Authority: 38 U.S.C. 1155)
[61 FR 52700, Oct. 8, 1996]

§ 4.128 Convalescence ratings following extended hospitalization.

If a mental disorder has been assigned a total evaluation due to a continuous period of hospitalization lasting six months or more, the rating agency shall continue the total evaluation indefinitely and schedule a mandatory examination six months after the veteran is discharged or released to nonbed care. A change in evaluation based on that or any subsequent exam-

ination shall be subject to the provisions of § 3.105(e) of this chapter.

(Authority: 38 U.S.C. 1155)
[61 FR 52700, Oct. 8, 1996]

§ 4.129 Mental disorders due to traumatic stress.

When a mental disorder that develops in service as a result of a highly stressful event is severe enough to bring about the veteran's release from active military service, the rating agency shall assign an evaluation of not less than 50 percent and schedule an examination within the six month period following the veteran's discharge to determine whether a change in evaluation is warranted.

(Authority: 38 U.S.C. 1155)
[61 FR 52700, Oct. 8, 1996]

§ 4.130 Schedule of ratings—mental disorders.

The nomenclature employed in this portion of the rating schedule is based upon the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, of the American Psychiatric Association (DSM-IV). Rating agencies must be thoroughly familiar with this manual to properly implement the directives in § 4.125 through § 4.129 and to apply the general rating formula for mental disorders in § 4.130. The schedule for rating for mental disorders is set forth as follows:

| | Rating |
|--|--------|
| Schizophrenia and Other Psychotic Disorders | |
| 9201 Schizophrenia, disorganized type | |
| 9202 Schizophrenia, catatonic type | |
| 9203 Schizophrenia, paranoid type | |
| 9204 Schizophrenia, undifferentiated type | |
| 9205 Schizophrenia, residual type; other and unspecified types | |
| 9208 Delusional disorder | |
| 9210 Psychotic disorder, not otherwise specified (atypical psychosis) | |
| 9211 Schizoaffective disorder | |
| Delirium, Dementia, and Amnestic and Other Cognitive Disorders | |
| 9300 Delirium | |
| 9301 Dementia due to infection (HIV infection, syphilis, or other systemic or intracranial infections) | |
| 9304 Dementia due to head trauma | |
| 9305 Vascular dementia | |
| 9310 Dementia of unknown etiology | |
| 9312 Dementia of the Alzheimer's type | |
| 9326 Dementia due to other neurologic or general medical conditions (endocrine disorders, metabolic disorders, Pick's disease, brain tumors, etc.) or that are substance-induced (drugs, alcohol, poisons) | |
| 9327 Organic mental disorder, other (including personality change due to a general medical condition) | |

| | Rating |
|--|--------|
| Anxiety Disorders | |
| 9400 Generalized anxiety disorder | |
| 9403 Specific (simple) phobia; social phobia | |
| 9404 Obsessive compulsive disorder | |
| 9410 Other and unspecified neurosis | |
| 9411 Post-traumatic stress disorder | |
| 9412 Panic disorder and/or agoraphobia | |
| 9413 Anxiety disorder, not otherwise specified | |
| Dissociative Disorders | |
| 9416 Dissociative amnesia; dissociative fugue; dissociative identity disorder (multiple personality disorder) | |
| 9417 Depersonalization disorder | |
| Somatoform Disorders | |
| 9421 Somatization disorder | |
| 9422 Pain disorder | |
| 9423 Undifferentiated somatoform disorder | |
| 9424 Conversion disorder | |
| 9425 Hypochondriasis | |
| Mood Disorders | |
| 9431 Cyclothymic disorder | |
| 9432 Bipolar disorder | |
| 9433 Dysthymic disorder | |
| 9434 Major depressive disorder | |
| 9435 Mood disorder, not otherwise specified | |
| Chronic Adjustment Disorder | |
| 9440 Chronic adjustment disorder | |
| General Rating Formula for Mental Disorders: | |
| Total occupational and social impairment, due to such symptoms as: gross impairment in thought processes or communication; persistent delusions or hallucinations; grossly inappropriate behavior; persistent danger of hurting self or others; intermittent inability to perform activities of daily living (including maintenance of minimal personal hygiene); disorientation to time or place; memory loss for names of close relatives, own occupation, or own name | 100 |
| Occupational and social impairment, with deficiencies in most areas, such as work, school, family relations, judgment, thinking, or mood, due to such symptoms as: suicidal ideation; obsessional rituals which interfere with routine activities; speech intermittently illogical, obscure, or irrelevant; near-continuous panic or depression affecting the ability to function independently, appropriately and effectively; impaired impulse control (such as unprovoked irritability with periods of violence); spatial disorientation; neglect of personal appearance and hygiene; difficulty in adapting to stressful circumstances (including work or a worklike setting); inability to establish and maintain effective relationships | 70 |
| Occupational and social impairment with reduced reliability and productivity due to such symptoms as: flattened affect; circumstantial, circumlocutory, or stereotyped speech; panic attacks more than once a week; difficulty in understanding complex commands; impairment of short- and long-term memory (e.g., retention of only highly learned material, forgetting to complete tasks); impaired judgment; impaired abstract thinking; disturbances of motivation and mood; difficulty in establishing and maintaining effective work and social relationships | 50 |
| Occupational and social impairment with occasional decrease in work efficiency and intermittent periods of inability to perform occupational tasks (although generally functioning satisfactorily, with routine behavior, self-care, and conversation normal), due to such symptoms as: depressed mood, anxiety, suspiciousness, panic attacks (weekly or less often), chronic sleep impairment, mild memory loss (such as forgetting names, directions, recent events) | 30 |
| Occupational and social impairment due to mild or transient symptoms which decrease work efficiency and ability to perform occupational tasks only during periods of significant stress, or; symptoms controlled by continuous medication | 10 |
| A mental condition has been formally diagnosed, but symptoms are not severe enough either to interfere with occupational and social functioning or to require continuous medication | 0 |
| Eating Disorders | |
| 9520 Anorexia nervosa | |
| 9521 Bulimia nervosa | |

| | Rating |
|--|--------|
| Rating Formula for Eating Disorders: | |
| Self-induced weight loss to less than 80 percent of expected minimum weight, with incapacitating episodes of at least six weeks total duration per year, and requiring hospitalization more than twice a year for parenteral nutrition or tube feeding | 100 |
| Self-induced weight loss to less than 85 percent of expected minimum weight with incapacitating episodes of six or more weeks total duration per year | 60 |
| Self-induced weight loss to less than 85 percent of expected minimum weight with incapacitating episodes of more than two but less than six weeks total duration per year | 30 |
| Binge eating followed by self-induced vomiting or other measures to prevent weight gain, or resistance to weight gain even when below expected minimum weight, with diagnosis of an eating disorder and incapacitating episodes of up to two weeks total duration per year | 10 |
| Binge eating followed by self-induced vomiting or other measures to prevent weight gain, or resistance to weight gain even when below expected minimum weight, with diagnosis of an eating disorder but without incapacitating episodes | 0 |

NOTE: An incapacitating episode is a period during which bed rest and treatment by a physician are required.

(Authority: 38 U.S.C. 1155)

[61 FR 52700, Oct. 8, 1996]

DENTAL AND ORAL CONDITIONS

§ 4.149 [Reserved]

§ 4.150 Schedule of ratings—dental and oral conditions.

| | Rating | | Rating |
|------|--------|--|--------|
| | | Not involving loss of temporomandibular articulation | |
| | | Bilateral | 30 |
| | | Unilateral | 20 |
| 9900 | | Ramus, loss of less than one-half the substance of, not involving loss of continuity: | |
| | | Bilateral | 20 |
| | | Unilateral | 10 |
| 9908 | | Condylod process, loss of, one or both sides | 30 |
| 9909 | | Coronoid process, loss of: | |
| | | Bilateral | 20 |
| | | Unilateral | 10 |
| 9911 | | Hard palate, loss of half or more: | |
| | | Not replaceable by prosthesis | 30 |
| | | Replaceable by prosthesis | 10 |
| 9912 | | Hard palate, loss of less than half of: | |
| | | Not replaceable by prosthesis | 20 |
| | | Replaceable by prosthesis | 0 |
| 9913 | | Teeth, loss of, due to loss of substance of body of maxilla or mandible without loss of continuity: | |
| | | Where the lost masticatory surface cannot be restored by suitable prosthesis: | |
| | | Loss of all teeth | 40 |
| | | Loss of all upper teeth | 30 |
| | | Loss of all lower teeth | 30 |
| | | All upper and lower posterior teeth missing | 20 |
| | | All upper and lower anterior teeth missing ... | 20 |
| | | All upper anterior teeth missing | 10 |
| | | All lower anterior teeth missing | 10 |
| | | All upper and lower teeth on one side missing | 10 |
| | | Where the loss of masticatory surface can be restored by suitable prosthesis | 0 |
| | | NOTE—These ratings apply only to bone loss through trauma or disease such as osteomyelitis, and not to the loss of the alveolar process as a result of periodontal disease, since such loss is not considered disabling. | |
| 9914 | | Maxilla, loss of more than half: | |
| | | Not replaceable by prosthesis | 100 |
| | | Replaceable by prosthesis | 50 |
| 9915 | | Maxilla, loss of half or less: | |
| 9900 | 100 | Maxilla or mandible, chronic osteomyelitis or osteoradionecrosis of: | |
| | | Rate as osteomyelitis, chronic under diagnostic code 5000. | |
| 9901 | 100 | Mandible, loss of, complete, between angles | |
| 9902 | | Mandible, loss of approximately one-half: | |
| | | Involving temporomandibular articulation | 50 |
| | | Not involving temporomandibular articulation | 30 |
| 9903 | | Mandible, nonunion of: | |
| | | Severe | 30 |
| | | Moderate | 10 |
| | | NOTE—Dependent upon degree of motion and relative loss of masticatory function. | |
| 9904 | | Mandible, malunion of: | |
| | | Severe displacement | 20 |
| | | Moderate displacement | 10 |
| | | Slight displacement | 0 |
| | | NOTE—Dependent upon degree of motion and relative loss of masticatory function. | |
| 9905 | | Temporomandibular articulation, limited motion of: | |
| | | Inter-incisal range: | |
| | | 0 to 10 mm | 40 |
| | | 11 to 20 mm | 30 |
| | | 21 to 30 mm | 20 |
| | | 31 to 40 mm | 10 |
| | | Range of lateral excursion: | |
| | | 0 to 4 mm | 10 |
| | | NOTE—Ratings for limited inter-incisal movement shall not be combined with ratings for limited lateral excursion. | |

Pt. 4, App.A

38 CFR Ch. I (7-1-01 Edition)

| | Rating |
|--|--------|
| Loss of 25 to 50 percent: | |
| Not replaceable by prosthesis | 40 |
| Replaceable by prosthesis | 30 |
| Loss of less than 25 percent: | |
| Not replaceable by prosthesis | 20 |
| Replaceable by prosthesis | 0 |
| 9916 Maxilla, malunion or nonunion of: | |
| Severe displacement | 30 |
| Moderate displacement | 10 |
| Slight displacement | 0 |

[59 FR 2530, Jan. 18, 1994]

APPENDIX A TO PART 4—TABLE OF AMENDMENTS AND EFFECTIVE DATES SINCE 1946

| Sec. | |
|-------|---|
| 4.16 | Last sentence; March 1, 1963. |
| 4.17 | October 7, 1948. |
| 4.17a | March 1, 1963. |
| 4.29 | Introductory portion preceding paragraph (a); March 1, 1963. Paragraph (a) "first day of continuous hospitalization"; April 8, 1959. Paragraph (a) "terminated last day of month"; December 1, 1962. Paragraph (a) penultimate sentence; November 13, 1970. Paragraph (b); April 8, 1959. Paragraph (c); August 16, 1948. Paragraph (d); August 16, 1948. Paragraph (e); March 1, 1963. Paragraph (f); August 9, 1976. NOTE: Application of this section to psychoneurotic and psychophysiologic disorders effective October 1, 1961. |
| 4.30 | Introductory portion of paragraph (a) preceding subparagraph (1); July 6, 1950. Paragraph (a)(1); June 9, 1952. Paragraph (a)(2); June 9, 1952. Paragraph (a)(3); June 9, 1952. Effective as to outpatient treatment March 10, 1976. Paragraph (b)(1); March 1, 1963. Paragraph (b)(2); August 9, 1976. |
| 4.55 | Paragraph (b) first sentence; March 1, 1963. |
| 4.63 | June 17, 1948. |
| 4.64 | October 1, 1956. |
| 4.71a | Diagnostic Code 5000—60 percent; February 1, 1962. Diagnostic Code 5000 NOTE (2): First three sentences; July 10, 1956. Last sentence; July 6, 1950. Diagnostic Code 5002—100 percent, 60 percent, 40 percent, 20 percent; March 1, 1963. Diagnostic Code 5003; July 6, 1950. Diagnostic Code 5012—NOTE; March 10, 1976. In sentence following DC 5024: "except gout which will be rated under 5002"; March 1, 1963. Diagnostic Code 5051; Diagnostic Code 5052; Diagnostic Code 5053; Diagnostic Code 5054; September 9, 1975. Diagnostic Code 5055; September 9, 1975. Diagnostic Code 5056; Diagnostic Code 5164—60 percent; June 9, 1952. Diagnostic Code 5172; July 6, 1950. Diagnostic Code 5173; June 9, 1952. Diagnostic Code 5255 "or hip"; July 6, 1950. |

| Sec. | |
|-------|--|
| | Diagnostic Code 5257—Evaluation; July 6, 1950. Diagnostic Code 5297—(Removal of one rib) "or resection of 2 or more"; August 23, 1948. Diagnostic Code 5297—NOTE (2): Reference to lobectomy; pneumonectomy and graduated ratings; February 1, 1962. Diagnostic Code 5298; August 23, 1948. Diagnostic Code 5324; February 1, 1962. Diagnostic Code 5327; March 10, 1976. Diagnostic Code 5328; March 10, 1976. |
| 4.73 | Last sentence; December 1, 1963. Diagnostic Code 6029—Note; August 23, 1948. Diagnostic Code 6035; September 9, 1975. Diagnostic Code 6076—60%: Vision 1 eye 15/200 and other eye 20/100; August 23, 1948. Diagnostic Code 6080—Note—"as to 38 U.S.C. 1114(L)"; July 6, 1950. Diagnostic Code 6081—Words "unilateral", "minimal" and all of Note; March 10, 1976. |
| 4.84a | Removed-December 18, 1987 (text redesignated § 4.871, December 18, 1987) March 23, 1956. December 18, 1987. March 23, 1956. December 18, 1987. March 23, 1956. December 18, 1987. Tables VI and VII replaced by new Tables VI and VII December 18, 1987. Diagnostic Codes 6277 through 6297; March 23, 1956; removed December 18, 1987. (Text from § 4.84b redesignated § 4.87a, December 18, 1987). |
| 4.84b | Diagnostic Code 6304—Notes (1) and (2); August 23, 1948. Diagnostic Code 6309; March 1, 1963. Diagnostic Code 6350; 80% Evaluation and Criterion for 60% and 30% Evaluations; March 10, 1976. Other Evaluations and Note; March 1, 1963. |
| 4.85 | Ratings for nonpulmonary TB; December 1, 1949. |
| 4.86 | Diagnostic Code 6600—100% Evaluations and Criteria for 60%; September 9, 1975. Diagnostic Code 6602—Criteria for all Evaluations and Note; September 9, 1975. Diagnostic Code 6603; September 9, 1975. Second note following Diagnostic Code 6724; December 1, 1949. Diagnostic Code 6802—Criteria for all Evaluations; September 9, 1975. Diagnostic Code 6819—Note; March 10, 1976. Diagnostic Code 6821—Evaluations and Note; August 23, 1948. |
| 4.86a | Diagnostic Code 7000—30 percent; July 6, 1950. Diagnostic Code 7000—100 percent inactive "with signs of congestive failure upon any exertion beyond rest in bed" revoked; Diagnostic Code 7005—80 percent revoked; Diagnostic Code 7007—80 percent revoked; Diagnostic Code 7015—100 percent Evaluation. Criteria for All Evaluations and NOTES (1) and (2); September 9, 1975. Diagnostic Code 7016; September 9, 1975. Diagnostic Code 7017; Diagnostic Code 7100—20 percent; July 6, 1950. Diagnostic Code 7101 "or more"; September 1, 1960. Diagnostic Code 7101—NOTE (2); September 9, 1975. Diagnostic Code 7110—Criteria for 100 percent, NOTE and 60 percent and 20 percent Evaluations; September 9, 1975. |
| 4.87 | |
| 4.87a | |
| 4.88a | |
| 4.89 | |
| 4.97 | |
| 4.104 | |

Department of Veterans Affairs

Pt. 4, App. B

| Sec. | |
|--|--|
| 4.114 | Diagnostic Code 7111—NOTE; September 9, 1975. |
| | Diagnostic Codes 7114, 7115, 7116, and NOTE; June 9, 1952. |
| | Diagnostic Code 7117 and NOTE; June 9, 1952. NOTE following Diagnostic Code 7120; July 6, 1950. |
| | Diagnostic Code 7121—100 percent Criterion and Evaluation and 60 percent Criterion; March 10, 1976. Criteria for 30 percent and 10 percent and NOTE; July 6, 1950. |
| | Last sentence of NOTE following Diagnostic Code 7122; July 6, 1950. |
| | Diagnostic Codes 7304 and 7305—Evaluations; November 1, 1962. |
| | Diagnostic Code 7308—Evaluations; April 8, 1959. |
| | Diagnostic Code 7312—70% Evaluation and 50% Evaluation and Criterion; March 10, 1976. |
| | Diagnostic Code 7313—20% Evaluation; March 10, 1976. |
| | Diagnostic Code 7319—Evaluations; November 1, 1962. |
| | Diagnostic Code 7321—Evaluations and Note; July 6, 1950. |
| | Diagnostic Code 7328—Evaluations and Note; November 1, 1962. |
| | Diagnostic Code 7329—Evaluations and Note; November 1, 1962. |
| | Diagnostic Code 7330—60% Evaluation; November 1, 1962. |
| | Diagnostic Code 7332—60% Evaluation; November 1, 1962. |
| | Diagnostic Code 7334—50% and 30% Evaluations; July 6, 1950. |
| | Diagnostic Code 7334—10% Evaluation; November 1, 1962. |
| | Diagnostic Code 7339—Criterion for 20% Evaluation; March 10, 1976. |
| | Diagnostic Code 7343—Note; March 10, 1976. |
| | Diagnostic Code 7345—100%, 60% and 30% Evaluations; August 23, 1948. |
| | Diagnostic Code 7345—10% Evaluation; February 17, 1955. |
| | Diagnostic Code 7345—10% Evaluation; February 17, 1955. |
| | Diagnostic Code 7346—Evaluations; February 1, 1962. |
| | Diagnostic Code 7347; September 9, 1975. |
| | Diagnostic Code 7348; March 10, 1976. |
| | Diagnostic Code 7500—Note; July 6, 1950. |
| | Diagnostic Code 7519—20%, 40% and 60% Evaluations; March 10, 1976. |
| | Diagnostic Code 7524—Note; July 6, 1950. |
| | Diagnostic Code 7528—Note; March 10, 1976. |
| | Diagnostic Code 7530; September 9, 1975. |
| | Diagnostic Code 7531; September 9, 1975. |
| | Diagnostic Code 7627—Note; March 10, 1976. |
| | Diagnostic Code 7703—Evaluations; August 23, 1948. |
| Diagnostic Code 7709—Note; March 10, 1976. Evaluations; June 9, 1952. | |
| Diagnostic Code 7714; September 9, 1975. | |
| Diagnostic Code 7801—Note (2); July 6, 1950. | |
| Diagnostic Code 7804—Note; July 6, 1950. | |
| Diagnostic Code 7900—10% Evaluation; and Notes (2) and (3); August 13, 1981. | |
| Diagnostic Code 7902—20% Evaluation; August 13, 1981. | |
| Diagnostic Code 7903—10% Evaluation; August 13, 1981. | |
| Diagnostic Code 7905—10% Evaluation; August 13, 1981. | |

| Sec. | |
|-----------------|--|
| 4.122 4.124a | Diagnostic Code 7907—60% Evaluation; August 13, 1981. |
| | Diagnostic Code 7909—40% and 20% Evaluation; August 13, 1981. |
| | Diagnostic Code 7911—Evaluations and Note; March 1, 1963; 40% and 20% Evaluations; August 13, 1981. |
| | Diagnostic Code 7913—Note; September 9, 1975. |
| | Diagnostic Code 7914—Note; March 10, 1976. October 1, 1961. |
| | Diagnostic Code 8002, NOTE; |
| | Diagnostic Code 8021, NOTE; |
| | Diagnostic Code 8045; October 1, 1961. |
| | Diagnostic Code 8046; October 1, 1961. |
| | Diagnostic Code 8100—Evaluations; June 9, 1953. |
| 4.125– 4.132 | Diagnostic Codes 8910 through 8914; October 1, 1961. |
| | Diagnostic Codes 8910 through 8914 General Rating Formula—Criteria and Evaluations; September 9, 1975. |
| | All Diagnostic Codes under Mental Disorders; October 1, 1961, except as to evaluation for Diagnostic Codes 9500 through 9511; September 9, 1975. |

[29 FR 6718, May 22, 1964, as amended at 34 FR 5064, Mar. 11, 1969; 40 FR 42541, Sept. 15, 1975; 41 FR 11291, Mar. 18, 1976; 41 FR 34258, Aug. 13, 1976; 43 FR 45362, Oct. 2, 1978; 46 FR 43666, Aug. 31, 1981; 52 FR 44122, Nov. 18, 1987; 52 FR 46439, Dec. 7, 1987]

APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diagnostic Code No. | |
|---------------------|---|
| 5000 | Osteomyelitis, acute, subacute, or chronic. |
| 5001 | Bones and Joints, tuberculosis of. |
| 5002 | Arthritis, rheumatoid (atrophic). |
| 5003 | Arthritis, degenerative, hypertrophic, or osteoarthritis. |
| 5004 | Arthritis, gonorrheal. |
| 5005 | Arthritis, pneumococcal. |
| 5006 | Arthritis, typhoid. |
| 5007 | Arthritis, syphilitic. |
| 5008 | Arthritis, streptococcal. |
| 5009 | Arthritis, other types. |
| 5010 | Arthritis, due to trauma. |
| 5011 | Bones, caisson disease of. |
| 5012 | Bones, new growths of, malignant. |
| 5013 | Osteoporosis, with joint manifestations. |
| 5014 | Osteomalacia. |
| 5015 | Bones, new growths of, benign. |
| 5016 | Osteitis deformans. |
| 5017 | Gout. |
| 5018 | Hydrarthrosis, intermittent. |
| 5019 | Bursitis. |
| 5020 | Synovitis. |
| 5021 | Myositis. |
| 5022 | Periostitis. |
| 5023 | Myositis ossificans. |
| 5024 | Tenosynovitis. |

Pt. 4, App. B

38 CFR Ch. I (7-1-01 Edition)

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diag- nos- tic Code No. | |
|--|--|
| COMBINATIONS OF DISABILITIES | |
| 5100 | Anatomical loss of both hands and both feet. |
| 5101 | Loss of use of both hands and both feet. |
| 5102 | Anatomical loss of both hands and one foot. |
| 5103 | Anatomical loss of both feet and one hand. |
| 5104 | Loss of use of both hands and one foot. |
| 5105 | Loss of use of both feet and one hand. |
| 5106 | Anatomical loss of both hands. |
| 5107 | Anatomical loss of both feet. |
| 5108 | Anatomical loss of one hand and one foot. |
| 5109 | Loss of use of both hands. |
| 5110 | Loss of use of both feet. |
| 5111 | Loss of use of one hand and one foot. |
| AMPUTATIONS: UPPER EXTREMITY | |
| Arm, amputation of: | |
| 5120 | Disarticulation. |
| 5121 | Above insertion of deltoid. |
| 5122 | Below insertion of deltoid. |
| Forearm, amputation of: | |
| 5123 | Above insertion of pronator teres. |
| 5124 | Below insertion of pronator teres. |
| 5125 | Hand, loss of use of. |
| 5126 | Five digits of one hand, amputation of: |
| Four digits of one hand, amputation of: | |
| 5127 | Thumb, index, middle and ring. |
| 5128 | Thumb, index, middle and little. |
| 5129 | Thumb, index, ring and little. |
| 5130 | Thumb, middle, ring and little. |
| 5131 | Index, middle, ring and little. |
| Three digits of one hand, amputation of: | |
| 5132 | Thumb, index and middle. |
| 5133 | Thumb, index and ring. |
| 5134 | Thumb, index and little. |
| 5135 | Thumb, middle and ring. |
| 5136 | Thumb, middle and little. |
| 5137 | Thumb, ring and little. |
| 5138 | Index, middle and ring. |
| 5139 | Index, middle and little. |
| 5140 | Index, ring and little. |
| 5141 | Middle, ring and little. |
| Two digits of one hand, amputation of: | |
| 5142 | Thumb and index. |
| 5143 | Thumb and middle. |
| 5144 | Thumb and ring. |
| 5145 | Thumb and little. |
| 5146 | Index and middle. |
| 5147 | Index and ring. |
| 5148 | Index and little. |
| 5149 | Middle and ring. |
| 5150 | Middle and little. |
| 5151 | Ring and little. |
| 5152 | Thumb, amputation of. |
| 5153 | Index finger, amputation of. |
| 5154 | Middle finger, amputation of. |
| 5155 | Ring finger, amputation of. |
| 5156 | Little finger, amputation of. |

AMPUTATIONS: LOWER EXTREMITY

| | |
|-----------------------|-------------------------|
| Thigh, amputation of: | |
| 5160 | Disarticulation. |
| 5161 | Upper third. |
| 5162 | Middle or lower thirds. |
| Leg, amputation of: | |
| 5163 | With defective stump. |

| Diag- nos- tic Code No. | |
|-------------------------------------|--|
| 5164 | With loss of natural knee action. |
| 5165 | At a lower level. |
| 5166 | Forefoot, amputation proximal to metatarsal bones. |
| 5167 | Foot, loss of use of. |
| 5170 | Toes, all, amputation of, without metatarsal loss. |
| 5171 | Toe, great, amputation of. |
| 5172 | Toe, other, amputation of. |
| 5173 | Toes, three or more, amputation of, not including great toe. |

THE SHOULDER AND ARM

| | |
|------|--|
| 5200 | Scapulohumeral articulation, ankylosis of. |
| 5201 | Arm, limitation of motion of. |
| 5202 | Humerus, other impairment of. |
| 5203 | Clavicle or scapula, impairment of. |

THE ELBOW AND FOREARM

| | |
|------|--|
| 5205 | Elbow, ankylosis of. |
| 5206 | Forearm, limitation of flexion of. |
| 5207 | Forearm, limitation of extension of. |
| 5208 | Forearm, flexion limited to 100° and extension to 45°. |
| 5209 | Elbow, other impairment of. |
| 5210 | Radius and ulna, nonunion of, with flail false joint. |
| 5211 | Ulna, impairment of. |
| 5212 | Radius, impairment of. |
| 5213 | Supination and pronation, impairment of. |

THE WRIST AND HAND

| | |
|------|---|
| 5214 | Wrist, ankylosis. |
| 5215 | Wrist, limitation of motion of. |
| 5216 | Five digits of one hand, unfavorable ankylosis of. |
| 5217 | Four digits of one hand, unfavorable ankylosis of. |
| 5218 | Three digits of one hand, unfavorable ankylosis of. |
| 5219 | Two digits of one hand, unfavorable ankylosis of. |
| 5220 | Five digits of one hand, favorable ankylosis of. |
| 5221 | Four digits of one hand, favorable ankylosis of. |
| 5222 | Three digits of one hand, favorable ankylosis of. |
| 5223 | Two digits of one hand, favorable ankylosis of. |
| 5224 | Thumb, ankylosis of. |
| 5225 | Index finger, ankylosis of. |
| 5226 | Middle finger, ankylosis of. |
| 5227 | Finger, any other, ankylosis of. |

THE HIP AND THIGH

| | |
|------|------------------------------------|
| 5250 | Hip, ankylosis of. |
| 5251 | Thigh, limitation of extension of. |
| 5252 | Thigh, limitation of flexion of. |
| 5253 | Thigh, impairment of. |
| 5254 | Hip, flail joint. |
| 5255 | Femur, impairment of. |

THE KNEE AND LEG

| | |
|------|-----------------------------------|
| 5256 | Knee, ankylosis of. |
| 5257 | Knee, other impairment of. |
| 5258 | Cartilage, semilunar, dislocated. |
| 5259 | Cartilage, semilunar, removal of. |
| 5260 | Leg, limitation of flexion of. |
| 5261 | Leg, limitation of extension of. |
| 5262 | Tibia and fibula, impairment of. |
| 5263 | Genu recurvatum. |

Department of Veterans Affairs

Pt. 4, App. B

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag-
nos-
tic
Code
No.

Diag-
nos-
tic
Code
No.

THE ANKLE

5270 Ankle, ankylosis of.
5271 Ankle, limited motion of.
5272 Subastragalar or tarsal joint, ankylosis of.
5273 Os calcis or astragalus, malunion of.
5274 Astragalectomy.

5312 Group XII—Anterior muscles of the leg.
5313 Group XIII—Posterior thigh group.
5314 Group XIV—Anterior thigh group.
5315 Group XV—Mesial thigh group.
5316 Group XVI—Pelvic girdle group 1.
5317 Group XVII—Pelvic girdle group 2.
5318 Group XVIII—Pelvic girdle group 3.
5319 Group XIX—Muscles of the abdominal wall.
5320 Group XX—Spinal muscles.
5321 Group XXI—Muscles of respiration.
5322 Group XXII—Lateral, supra and infrahyoid group.
5323 Group XXIII—Lateral and posterior muscles of the neck.
5324 Diaphragm, rupture of.
5325 Muscle injury, facial muscles.
5326 Muscle hernia.

SHORTENING OF THE LOWER EXTREMITY

5275 Bones, of the lower extremity, shortening of.

THE FOOT

5276 Flatfoot, acquired.
5277 Weak foot, bilateral.
5278 Claw foot (pes cavus), acquired.
5279 Metatarsalgia, anterior (Morton's disease).
5280 Hallux valgus.
5281 Hallux rigidus.
5282 Hammer toe.
5283 Tarsal, or metatarsal bones, malunion of, or nonunion of.
5284 Foot injuries, other.

DISEASES OF THE EYE

6000 Uveitis.
6001 Keratitis.
6002 Scleritis.
6003 Iritis.
6004 Cyclitis.
6005 Choroiditis.
6006 Retinitis.
6007 Hemorrhage, intra-ocular, recent.
6008 Retina, detachment of.
6009 Eye, injury of, unhealed.
6010 Eye, tuberculosis of.
6011 Retina, localized scars.
6012 Glaucoma, congestive or inflammatory.
6013 Glaucoma, simple, primary, noncongestive.
6014 New growths, malignant, eyeball.
6015 New growths, benign, eyeball and adnexa.
6016 Nystagmus, central.
6017 Conjunctivitis, trachomatous, chronic.
6018 Conjunctivitis, other, chronic.
6019 Ptosis, eyelids.
6020 Ectropion.
6021 Entropion.
6022 Lagophthalmos.
6023 Eyebrows, loss of.
6024 Eyelashes, loss of.
6025 Epiphora.
6026 Neuritis, optic.
6027 Cataract, traumatic.
6028 Cataract, senile, and others.
6029 Aphakia.
6030 Accommodation, paralysis of.
6031 Dacryocystitis.
6032 Eyelids, loss of portion of.
6033 Lens, crystalline, dislocation of.
6034 Pterygium.

THE SPINE

5285 Vertebra, fracture of, residuals.
5286 Spine, complete bony fixation (ankylosis) of.
5287 Spine, ankylosis of, cervical.
5288 Spine, ankylosis of, dorsal.
5289 Spine, ankylosis of, lumbar.
5290 Spine, limitation of motion of, cervical.
5291 Spine, limitation of motion of, dorsal.
5292 Spine, limitation of motion of, lumbar.
5293 Intervertebral disc syndrome.
5294 Sacroiliac injury and weakness.
5295 Lumbosacral strain.

THE SKULL

5296 Skull, loss of part of, both inner and outer tables.

THE RIBS

5297 Ribs, removal of.

THE COCCYX

5298 Coccyx, removal of.

MUSCLE INJURIES

5301 Group I—Extrinsic muscles of shoulder girdle.
5302 Group II—Extrinsic muscles of shoulder girdle.
5303 Group III—Intrinsic muscles of shoulder girdle.
5304 Group IV—Intrinsic muscles of shoulder girdle.
5305 Group V—Flexor muscles of the elbow.
5306 Group VI—Extensor muscles of the elbow.
5307 Group VII—Muscles arising from internal condyle of humerus.
5308 Group VIII—Muscles arising mainly from external condyle of humerus.
5309 Group IX—Intrinsic muscles of the hand.
5310 Group X—Intrinsic muscles of the foot.
5311 Group XI—Posterior and lateral muscles of the leg.

COMBINATIONS OF DISABILITIES

6050 Blindness in both eyes having only light perception and anatomical loss of both hands and both feet.
6051 Blindness in both eyes having only light perception and loss of use of both hands and both feet.
6052 Blindness in both eyes having only light perception and anatomical loss of both hands.
6053 Blindness in both eyes having only light perception and anatomical loss of both feet.
6054 Blindness in both eyes having only light perception and anatomical loss of one hand and one foot.
6055 Blindness in both eyes having only light perception and loss of use of both hands.

Pt. 4, App. B

38 CFR Ch. I (7-1-01 Edition)

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diag- nos- tic Code No. | |
|-------------------------------------|---|
| 6056 | Blindness in both eyes having only light perception and loss of use of both feet. |
| 6057 | Blindness in both eyes having only light perception and loss of use of one hand and one foot. |
| 6058 | Blindness in both eyes having only light perception and anatomical loss of one hand. |
| 6059 | Blindness in both eyes having only light perception and anatomical loss of one foot. |
| 6060 | Blindness in both eyes having only light perception and loss of use of one hand. |
| 6061 | Blindness in both eyes having only light perception and loss of use of one foot. |
| 6062 | Blindness in both eyes having only light perception. |
| IMPAIRMENT OF CENTRAL VISUAL ACUITY | |
| | Blindness, anatomical loss, one eye: |
| 6063 | Other blind (5/200 or less). |
| 6064 | Other impaired (20/200 or less). |
| 6065 | Other impaired. |
| 6066 | Other normal. |
| | Blindness, light perception only one eye: |
| 6067 | Other blind (5/200 or less). |
| 6068 | Other impaired (20/200 or less). |
| 6069 | Other impaired. |
| 6070 | Other normal. |
| | Blindness, total (5/200 or less): |
| 6071 | Both eyes. |
| | Blindness, total one eye (5/200 or less): |
| 6072 | Other impaired (20/200 or less). |
| 6073 | Other impaired. |
| 6074 | Other normal. |
| | Blindness, partial (20/200 or less): |
| 6075 | Both eyes. |
| | One eye: |
| 6076 | Other impaired. |
| 6077 | Other normal. |
| | Blindness, partial: |
| 6078 | Both eyes. |
| 6079 | One eye only. |
| 6080 | Field vision, impairment of. |
| 6081 | Scotoma, pathological. |
| 6090 | Muscle function, ocular, impairment of. |
| 6091 | Symblepharon. |
| 6092 | Diplopia, due to limited muscle function. |
| IMPAIRMENT OF AUDITORY ACUITY | |
| 6100 | 0% evaluation based on Table VII |
| 6101 | 10% evaluation based on Table VII |
| 6102 | 20% evaluation based on Table VII |
| 6103 | 30% evaluation based on Table VII |
| 6104 | 40% evaluation based on Table VII |
| 6105 | 50% evaluation based on Table VII |
| 6106 | 60% evaluation based on Table VII |
| 6107 | 70% evaluation based on Table VII |
| 6108 | 80% evaluation based on Table VII |
| 6109 | 90% evaluation based on Table VII |
| 6110 | 100% evaluation based on Table VII. |
| DISEASES OF THE EAR | |
| 6200 | Otitis media, suppurative, chronic. |
| 6201 | Otitis media, catarrhal, chronic. |
| 6202 | Otosclerosis. |
| 6203 | Otitis interna. |
| 6204 | Labyrinthitis. |
| 6205 | Meniere's syndrome. |

| Diag- nos- tic Code No. | |
|-------------------------------------|--|
| 6206 | Mastoiditis. |
| 6207 | Auricle, loss or deformity. |
| 6208 | New growths, malignant, ear. |
| 6209 | New growths, benign, ear. |
| 6210 | Auditory canal, disease of. |
| 6211 | Tympanic membrane, perforation of. |
| 6260 | Tinnitus. |
| OTHER SENSE ORGANS | |
| 6275 | Smell, loss of sense of. |
| 6276 | Taste, loss of sense of. |
| SYSTEMIC DISEASES | |
| 6300 | Cholera, Asiatic. |
| 6301 | Kala-azar (visceral leishmaniasis). |
| 6302 | Leprosy. |
| 6304 | Malaria. |
| 6305 | Filariasis. |
| 6306 | Oroya fever. |
| 6307 | Plague. |
| 6308 | Relapsing fever. |
| 6309 | Rheumatic fever. |
| 6310 | Syphilis, unspecified. |
| 6311 | Tuberculosis, military. |
| 6313 | Avitaminosis. |
| 6314 | Beriberi. |
| 6315 | Pellagra. |
| 6316 | Brucellosis (Malta or undulant fever). |
| 6317 | Typhus, scrub. |
| 6350 | Lupus erythematosus, systemic. |
| RESPIRATORY SYSTEM | |
| THE NOSE AND THROAT | |
| 6501 | Rhinitis, atrophic, chronic. |
| 6502 | Septum, nasal, deflection of. |
| 6504 | Nose, loss of part of, or scars. |
| 6510 | Sinusitis, pansinusitis, chronic. |
| 6511 | Sinusitis, ethmoid, chronic. |
| 6512 | Sinusitis, frontal, chronic. |
| 6513 | Sinusitis, maxillary, chronic. |
| 6514 | Sinusitis, sphenoid, chronic. |
| 6515 | Laryngitis, tuberculous. |
| 6516 | Laryngitis, chronic. |
| 6517 | Larynx, injuries of, healed. |
| 6518 | Laryngectomy. |
| 6519 | Aphonia, organic. |
| 6520 | Larynx, stenosis of. |
| THE TRACHEA AND BRONCHI | |
| 6600 | Bronchitis, chronic. |
| 6601 | Bronchiectasis. |
| 6602 | Asthma, bronchial. |
| THE LUNGS AND PLEURA | |
| 6701 | Tuberculosis, pulmonary, chronic, far advanced, active. |
| 6702 | Tuberculosis, pulmonary, chronic, moderately advanced, active. |
| 6703 | Tuberculosis, pulmonary, chronic, minimal, active. |
| 6704 | Tuberculosis, pulmonary, chronic, active, advancement unspecified. |

Department of Veterans Affairs

Pt. 4, App. B

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diag- nos- tic Code No. | |
|-------------------------------------|--|
| 6707 | Tuberculosis, pulmonary, chronic, far advanced, active. |
| 6708 | Tuberculosis, pulmonary, chronic, moderately advanced, active. |
| 6709 | Tuberculosis, pulmonary, chronic, minimal, active. |
| 6710 | Tuberculosis, pulmonary, chronic, active, advancement unspecified. |
| 6721 | Tuberculosis, pulmonary, chronic, far advanced, inactive. |
| 6722 | Tuberculosis, pulmonary, chronic, moderately advanced, inactive. |
| 6723 | Tuberculosis, pulmonary, chronic, minimal, inactive. |
| 6724 | Tuberculosis, pulmonary, chronic, inactive, advancement unspecified. |
| 6725 | Tuberculosis, pulmonary, chronic, far advanced, inactive. |
| 6726 | Tuberculosis, pulmonary, chronic, moderately advanced, inactive. |
| 6727 | Tuberculosis, pulmonary, chronic, minimal, inactive. |
| 6728 | Tuberculosis, pulmonary, chronic, inactive, advancement unspecified. |
| 6732 | Pleurisy, tuberculous. |
| 6800 | Anthracosis. |
| 6801 | Silicosis. |
| 6802 | Pneumoconiosis, unspecified. |
| 6803 | Actinomycosis of lung. |
| 6804 | Streptotrichosis of lung. |
| 6805 | Blastomycosis of lung. |
| 6806 | Sporotrichosis of lung. |
| 6807 | Aspergillosis of lung. |
| 6808 | Mycosis of lung, unspecified. |
| 6809 | Lung, abscess of. |
| 6810 | Pleurisy, serofibrinous. |
| 6811 | Pleurisy, purulent (empyema). |
| 6812 | Fistula, bronchocutaneous, or bronchopleural. |
| 6813 | Lung, permanent collapse of. |
| 6814 | Pneumothorax, spontaneous. |
| 6815 | Pneumonectomy. |
| 6816 | Lobectomy. |
| 6817 | Lung, chronic passive congestion of. |
| 6818 | Pleural cavity, injuries, residuals of, including gunshot wounds. |
| 6819 | New growths, malignant, any specified part of respiratory system. |
| 6820 | New growths, benign, any specified part of respiratory system. |
| 6821 | Coccidioidomycosis. |
| THE CARDIOVASCULAR SYSTEM | |
| THE HEART | |
| 7000 | Rheumatic heart disease. |
| 7001 | Endocarditis, bacterial, subacute. |
| 7002 | Pericarditis, bacterial or rheumatic, acute. |
| 7003 | Adhesions, pericardial. |
| 7004 | Syphilitic heart disease. |
| 7005 | Arteriosclerotic heart disease. |
| 7006 | Myocardium, infarction of, due to thrombosis or embolism. |
| 7007 | Hypertensive heart disease. |
| 7008 | Hyperthyroid heart disease. |
| 7010 | Auricular flutter, paroxysmal. |
| 7011 | Auricular fibrillation, paroxysmal. |
| 7012 | Auricular fibrillation, permanent. |
| 7013 | Tachycardia, paroxysmal. |
| 7014 | Sinus tachycardia. |
| 7015 | Auriculoventricular block. |

| Diag- nos- tic Code No. | |
|-------------------------------------|---|
| THE ARTERIES AND VEINS | |
| 7100 | Arteriosclerosis, general. |
| 7101 | Hypertensive vascular disease (essential arterial hypertension). |
| 7110 | Aorta or branches, aneurysm of. |
| 7111 | Artery, any large artery, aneurysm of. |
| 7112 | Artery, small aneurysmal dilatation. |
| 7113 | Arteriovenous aneurysm, traumatic. |
| 7114 | Arteriosclerosis obliterans. |
| 7115 | Thrombo-angitis obliterans (Buerger's disease). |
| 7116 | Claudication, intermittent. |
| 7117 | Raynaud's disease. |
| 7118 | Angioneurotic edema. |
| 7119 | Erythromelalgia. |
| 7120 | Varicose veins. |
| 7121 | Phlebitis. |
| 7122 | Frozen feet, residuals of (Immersion foot). |
| THE DIGESTIVE SYSTEM | |
| 7200 | Mouth, injuries of. |
| 7201 | Lips, injuries of. |
| 7202 | Tongue, loss of, whole or part. |
| 7203 | Esophagus, stricture of. |
| 7204 | Esophagus, spasm of (cardiospasm). |
| 7205 | Esophagus, diverticulum of, acquired. |
| 7301 | Peritoneum, adhesions of. |
| 7304 | Ulcer, gastric. |
| 7305 | Ulcer, duodenal. |
| 7306 | Ulcer, marginal (gastrojejunal). |
| 7307 | Gastritis, hypertrophic. |
| 7308 | Postgastroectomy syndromes. |
| 7309 | Stomach, stenosis of. |
| 7310 | Stomach, injury of, residuals. |
| 7311 | Liver, injury of. |
| 7312 | Liver, cirrhosis of. |
| 7313 | Liver, abscess of, residuals. |
| 7314 | Cholecystitis, chronic. |
| 7315 | Cholelithiasis, chronic. |
| 7316 | Cholangitis, chronic. |
| 7317 | Gall bladder, injury of. |
| 7318 | Gall bladder, removal of. |
| 7319 | Irritable colon syndrome (spastic colitis, mucous colitis, etc.). |
| 7321 | Amebiasis. |
| 7322 | Dysentery, bacillary. |
| 7323 | Colitis, ulcerative. |
| 7324 | Distomiasis, intestinal or hepatic. |
| 7325 | Enteritis, chronic. |
| 7326 | Enterocolitis, chronic. |
| 7327 | Diverticulitis. |
| 7328 | Intestine, small, resection of. |
| 7329 | Intestine, large, resection of. |
| 7330 | Intestine, fistula of. |
| 7331 | Peritonitis, tuberculous, active. |
| 7332 | Rectum and anus, impairment of sphincter control. |
| 7333 | Rectum and anus, stricture of. |
| 7334 | Rectum, persistent prolapse of. |
| 7335 | Ano, fistula in. |
| 7336 | Hemorrhoids, external or internal. |
| 7337 | Pruritus ani. |
| 7338 | Hernia, inguinal. |
| 7339 | Hernia, ventral. |
| 7340 | Hernia, femoral. |
| 7341 | Wounds, incised, healed, abdominal wall. |
| 7342 | Visceroptosis. |
| 7343 | New growths, malignant, any specified part of digestive system. |

Pt. 4, App. B

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diag- nos- tic Code No. | |
|-------------------------------------|--|
| 7344 | New growths, benign, any specified part of digestive system. |
| 7345 | Hepatitis, infectious. |
| 7346 | Hernia, hiatal. |
| THE GENITOURINARY SYSTEM | |
| 7500 | Kidney, removal of. |
| 7501 | Kidney, abscess of. |
| 7502 | Nephritis, chronic. |
| 7503 | Pyelitis. |
| 7504 | Pyelonephritis, chronic. |
| 7505 | Kidney, tuberculosis of, active. |
| 7507 | Nephrosclerosis, arteriolar. |
| 7508 | Nephrolithiasis. |
| 7509 | Hydronephrosis. |
| 7510 | Ureterolithiasis. |
| 7511 | Ureter, stricture of. |
| 7512 | Cystitis, chronic. |
| 7513 | Cystitis, interstitial (Hunner), submucous or elusive ulcer. |
| 7514 | Bladder, tuberculosis of. |
| 7515 | Bladder, calculus in. |
| 7516 | Bladder, fistula of. |
| 7517 | Bladder, injury of. |
| 7518 | Urethra, stricture of. |
| 7519 | Urethra, fistula of. |
| 7520 | Penis, removal of half or more. |
| 7521 | Penis, removal of glans. |
| 7522 | Penis, deformity, with loss of erectile power. |
| 7523 | Testis, atrophy, complete. |
| 7524 | Testis, removal of. |
| 7525 | Epididymo-orchitis (tuberculous). |
| 7526 | Prostate gland, resection or removal. |
| 7527 | Prostate gland injuries, infectious hypertrophy, post-operative residuals. |
| 7528 | New growths, malignant, any specified part of genitourinary system. |
| 7529 | New growths, benign, any specified part of genitourinary system. |
| GYNECOLOGICAL CONDITIONS | |
| 7610 | Vulvovaginitis. |
| 7611 | Vaginitis. |
| 7612 | Cervicitis. |
| 7613 | Metritis. |
| 7614 | Salpingitis. |
| 7615 | Oophoritis. |
| 7617 | Uterus and ovaries, removal of, complete. |
| 7618 | Uterus, removal of, including corpus. |
| 7619 | Ovaries, removal of. |
| 7620 | Ovaries, atrophy of both. |
| 7621 | Uterus, prolapse. |
| 7622 | Uterus, displacement of. |
| 7623 | Pregnancy, surgical complications of. |
| 7624 | Fistula, rectovaginal. |
| 7625 | Fistula, urethrovaginal. |
| 7626 | Mammary glands, removal of. |
| 7627 | New growth, malignant, gynecological system, or mammary glands. |
| THE HEMIC AND LYMPHATIC SYSTEMS | |
| 7700 | Anemia, pernicious. |
| 7701 | Anemia, secondary. |
| 7702 | Agranulocytosis, acute. |
| 7703 | Leukemia. |

38 CFR Ch. I (7-1-01 Edition)

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diag- nos- tic Code No. | |
|--|---|
| 7704 | Polycythemia, primary. |
| 7705 | Purpura hemorrhagica. |
| 7706 | Splenectomy. |
| 7707 | Spleen, injury of, healed. |
| 7709 | Lymphogranulomatosis (Hodgkin's disease). |
| 7710 | Adenitis, cervical, tuberculous. |
| 7711 | Adenitis, axillary, tuberculous. |
| 7712 | Adenitis, inguinal, tuberculous. |
| 7713 | Adenitis, secondary. |
| THE SKIN | |
| 7800 | Scars, disfiguring, head, face or neck. |
| 7801 | Scars, burns, third degree. |
| 7802 | Scars, burns, second degree. |
| 7803 | Scars, superficial, poorly nourished. |
| 7804 | Scars, superficial, tender and painful. |
| 7805 | Scars, others. |
| 7806 | Eczema. |
| 7807 | Leishmaniasis, americana (mucocutaneous, espundia). |
| 7808 | Leishmaniasis, old world (cutaneous, oriental sore). |
| 7809 | Lupus erythematosus, discoid. |
| 7810 | Pinta. |
| 7811 | Tuberculosis luposa (lupus vulgaris). |
| 7812 | Verruga peruana. |
| 7813 | Dermatophytosis. |
| 7814 | Tinea barbae. |
| 7815 | Pemphigus. |
| 7816 | Psoriasis. |
| 7817 | Dermatitis exfoliativa. |
| 7818 | New growths, malignant, skin. |
| 7819 | New growths, benign, skin. |
| THE ENDOCRINE SYSTEM | |
| 7900 | Hyperthyroidism. |
| 7901 | Thyroid gland, toxic adenoma of. |
| 7902 | Thyroid gland, non-toxic adenoma of. |
| 7903 | Hypothyroidism. |
| 7904 | Hyperparathyroidism (osteitis fibrosa cystica). |
| 7905 | Hypoparathyroidism. |
| 7907 | Hyperpituitarism (pituitary basophilism, Cushing's syndrome). |
| 7908 | Hyperpituitarism (acromegaly or gigantism). |
| 7909 | Hypopituitarism (diabetes insipidus). |
| 7910 | Hyperadrenia (adrenogenital syndrome). |
| 7911 | Addison's disease. |
| 7912 | Pluriglandular syndromes. |
| 7913 | Diabetes mellitus. |
| 7914 | New growths, malignant, endocrine system. |
| 7915 | New growths, benign, endocrine system. |
| NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS | |
| 8000 | Encephalitis, epidemic, chronic. |
| | Brain, new growth of: |
| 8002 | Malignant. |
| 8003 | Benign. |
| 8004 | Paralysis agitans. |
| 8005 | Bulbar palsy. |
| 8007 | Brain, vessels, embolism of. |
| 8008 | Brain, vessels, thrombosis of. |
| 8009 | Brain, vessels, hemorrhage from. |
| 8010 | Myelitis. |
| 8011 | Poliomyelitis, anterior. |
| 8012 | Hematomyelia. |
| 8013 | Syphilis, cerebrospinal. |

Department of Veterans Affairs

Pt. 4, App. B

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diag- nos- tic Code No. | |
|-------------------------------------|---|
| 8014 | Syphilis, meningovascular. |
| 8015 | Tabes dorsalis. |
| 8017 | Amyotrophic lateral sclerosis. |
| 8018 | Multiple sclerosis. |
| 8019 | Meningitis, cerebrospinal, epidemic. |
| 8020 | Brain, abscess of. Spinal cord, new growths: |
| 8021 | Malignant. |
| 8022 | Benign. |
| 8023 | Progressive muscular atrophy. |
| 8024 | Syringomyelia. |
| 8025 | Myasthenia gravis. |
| 8045 | Brain disease due to trauma. |
| 8046 | Cerebral arteriosclerosis. |
| 8100 | Migraine. |
| 8103 | Tic, convulsive. |
| 8104 | Paramyoclonus multiplex (convulsive state, myoclonic type). |
| 8105 | Chorea, Sydenham's. |
| 8106 | Chorea, Huntington's. |
| 8107 | Athetosis, acquired. |
| 8108 | Narcolepsy. |

THE CRANIAL NERVES

| | |
|------|---|
| 8205 | Fifth (trigeminal) cranial nerve, paralysis of. |
| 8207 | Seventh (facial) cranial nerve, paralysis of. |
| 8209 | Ninth (glossopharyngeal) cranial nerve, paralysis of. |
| 8210 | Tenth (pneumogastric, vagus) cranial nerve, paralysis of. |
| 8211 | Eleventh (spinal accessory, external branch) cranial nerve, paralysis of. |
| 8212 | Twelfth (hypoglossal) cranial nerve, paralysis of. |
| 8305 | Fifth (trigeminal) cranial nerve, neuritis. |
| 8307 | Seventh (facial) cranial nerve, neuritis. |
| 8309 | Ninth (glossopharyngeal) cranial nerve, neuritis. |
| 8310 | Tenth (pneumogastric, vagus) cranial nerve, neuritis. |
| 8311 | Eleventh (spinal accessory, external branch) cranial nerve, neuritis. |
| 8312 | Twelfth (hypoglossal) cranial nerve, neuritis. |
| 8407 | Seventh (facial) cranial nerve, neuralgia. |
| 8409 | Ninth (glossopharyngeal) cranial nerve, neuralgia. |
| 8410 | Tenth (pneumogastric, vagus) cranial nerve, neuralgia. |
| 8411 | Eleventh (spinal accessory, external branch) cranial nerve, neuralgia. |
| 8412 | Twelfth (hypoglossal) cranial nerve, neuralgia. |

PERIPHERAL NERVES: PARALYSIS

| | |
|------|--|
| 8510 | Upper radicular group (fifth and sixth cervicals), paralysis of. |
| 8511 | Middle radicular group, paralysis of. |
| 8512 | Lower radicular group, paralysis of. |
| 8513 | All radicular groups, paralysis of. |
| 8514 | The musculospiral nerve (radial nerve), paralysis of. |
| 8515 | The median nerve, paralysis of. |
| 8516 | The ulnar nerve, paralysis of. |
| 8517 | Musculocutaneous nerve, paralysis of. |
| 8518 | Circumflex nerve, paralysis of. |
| 8519 | Long thoracic nerve, paralysis of. |
| 8520 | The sciatic nerve, paralysis of. |
| 8521 | External popliteal nerve (common peroneal), paralysis of. |
| 8522 | Musculocutaneous nerve (superficial peroneal), paralysis of. |
| 8523 | Anterior tibial nerve (deep peroneal), paralysis of. |
| 8524 | Internal popliteal nerve (tibial), paralysis of. |
| 8525 | Posterior tibial nerve, paralysis of. |

| Diag- nos- tic Code No. | |
|-------------------------------------|--|
| 8526 | Anterior crural nerve (femoral), paralysis of. |
| 8527 | Internal saphenous nerve, paralysis of. |
| 8528 | Obturator nerve, paralysis of. |
| 8529 | External cutaneous nerve of thigh, paralysis of. |
| 8530 | Ilio-inguinal nerve, paralysis of. |

PERIPHERAL NERVES: NEURITIS

| | |
|------|--|
| 8610 | Upper radicular group (fifth and sixth cervicals), neuritis. |
| 8611 | Middle radicular group, neuritis. |
| 8612 | Lower radicular group, neuritis. |
| 8613 | All radicular groups, neuritis. |
| 8614 | The musculospiral nerve (radial nerve), neuritis. |
| 8615 | The median nerve, neuritis. |
| 8616 | The ulnar nerve, neuritis. |
| 8617 | Musculocutaneous nerve, neuritis. |
| 8618 | Circumflex nerve, neuritis. |
| 8619 | Long thoracic nerve, neuritis. |
| 8620 | The sciatic nerve, neuritis. |
| 8621 | External popliteal nerve (common peroneal), neuritis. |
| 8622 | Musculocutaneous nerve (superficial peroneal), neuritis. |
| 8623 | Anterior tibial nerve (deep peroneal), neuritis. |
| 8624 | Internal popliteal nerve (tibial) neuritis. |
| 8625 | Posterior tibial nerve, neuritis. |
| 8626 | Anterior crural nerve (femoral), neuritis. |
| 8627 | Internal saphenous nerve, neuritis. |
| 8628 | Obturator nerve, neuritis. |
| 8629 | External cutaneous nerve of thigh, neuritis. |
| 8630 | Ilio-inguinal nerve, neuritis. |

PERIPHERAL NERVES: NEURALGIA

| | |
|------|---|
| 8710 | Upper radicular group (fifth and sixth cervicals), neuralgia. |
| 8711 | Middle radicular group, neuralgia. |
| 8712 | Lower radicular group, neuralgia. |
| 8713 | All radicular groups, neuralgia. |
| 8714 | The musculospiral nerve (radial nerve), neuralgia. |
| 8715 | The median nerve, neuralgia. |
| 8716 | The ulnar nerve, neuralgia. |
| 8717 | Musculocutaneous nerve, neuralgia. |
| 8718 | Circumflex nerve, neuralgia. |
| 8719 | Long thoracic nerve, neuralgia. |
| 8720 | The sciatic nerve, neuralgia. |
| 8721 | External popliteal nerve (common peroneal), neuralgia. |
| 8722 | Musculocutaneous nerve (superficial peroneal), neuralgia. |
| 8723 | Anterior tibial nerve (deep peroneal), neuralgia. |
| 8724 | Internal popliteal nerve (tibial), neuralgia. |
| 8725 | Posterior tibial nerve, neuralgia. |
| 8726 | Anterior crural nerve (femoral), neuralgia. |
| 8727 | Internal saphenous nerve, neuralgia. |
| 8728 | Obturator nerve, neuralgia. |
| 8729 | External cutaneous nerve of thigh neuralgia. |
| 8730 | Ilio-inguinal nerve, neuralgia. |

THE EPILEPSIES

| | |
|------|-------------------------|
| 8910 | Epilepsy, grand mal. |
| 8911 | Epilepsy, petit mal. |
| 8912 | Jacksonian type. |
| 8913 | Epilepsy, diencephalic. |
| 8914 | Epilepsy, psychomotor. |

Pt. 4, App. C

38 CFR Ch. I (7-1-01 Edition)

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

| Diag- nos- tic Code No. | |
|-------------------------------------|--|
| PSYCHOTIC DISORDERS | |
| 9200 | Schizophrenic reaction, simple type. |
| 9201 | Schizophrenic reaction, hebephrenic type. |
| 9202 | Schizophrenic reaction, catatonic type. |
| 9203 | Schizophrenic reaction, paranoid type. |
| 9204 | Schizophrenic reaction, chronic undifferentiated type. |
| 9205 | Schizophrenic reaction, other. |
| 9206 | Manic depressive reaction. |
| 9207 | Psychotic depressive reaction. |
| 9208 | Paranoid reaction (specify). |
| 9209 | Involuntal psychotic reaction. |
| 9210 | Psychotic reaction, other. |

| | |
|-------------------------|--|
| ORGANIC BRAIN DISORDERS | |
| 9300 | Acute brain syndrome (associated with infection, trauma, circulatory disturbance, etc.). |
| 9301 | Chronic brain syndrome associated with central nervous system syphilis (all forms). |
| 9302 | Chronic brain syndrome associated with intracranial infections other than syphilis. |
| 9303 | Chronic brain syndrome associated with intoxication. |
| 9304 | Chronic brain syndrome associated with brain trauma. |
| 9305 | Chronic brain syndrome associated with cerebral arteriosclerosis. |
| 9306 | Chronic brain syndrome associated with circulatory disturbance other than cerebral arteriosclerosis. |
| 9307 | Chronic brain syndrome associated with convulsive disorder (idiopathic epilepsy). |
| 9308 | Chronic brain syndrome associated with disturbance of metabolism, growth or nutrition. |
| 9309 | Chronic brain syndrome associated with intracranial neoplasm. |
| 9310 | Chronic brain syndrome associated with diseases of unknown or uncertain cause. |
| 9311 | Chronic brain syndrome of unknown cause. |

| | |
|--------------------------|---------------------------------|
| PSYCHONEUROTIC DISORDERS | |
| 9400 | Anxiety reaction. |
| 9401 | Dissociative reaction. |
| 9402 | Conversion reaction. |
| 9403 | Phobic reaction. |
| 9404 | Obsessive compulsive reaction. |
| 9405 | Depressive reaction. |
| 9406 | Psychoneurotic reaction, other. |

| | |
|-----------------------------|--|
| PSYCHOPHYSIOLOGIC DISORDERS | |
| 9500 | Psychophysiologic skin reaction. |
| 9501 | Psychophysiologic cardiovascular reaction. |
| 9502 | Psychophysiologic gastrointestinal reaction. |
| 9503 | Psychophysiologic nervous system reaction. |
| 9504 | Psychophysiologic reaction, other. |

| | |
|----------------------------|---|
| DENTAL AND ORAL CONDITIONS | |
| 9900 | Maxilla or mandible, osteomyelitis of. |
| 9901 | Mandible, loss of, complete, between angles. |
| 9902 | Mandible, loss of approximately one-half. |
| 9903 | Mandible, nonunion of. |
| 9904 | Mandible, malunion of. |
| 9905 | Temporomandibular articulation, limited motion of. |
| 9906 | Ramus, loss of whole or part of. |
| 9907 | Ramus, loss of less than one-half the substance of, not involving loss of continuity. |

| Diag- nos- tic Code No. | |
|-------------------------------------|---|
| 9908 | Condylod process, loss of, one or both sides. |
| 9909 | Coronoid process, loss of. |
| 9910 | Maxilla, loss of whole or part of substance of, non-union of, or malunion of. |
| 9911 | Hard palate, loss of half or more. |
| 9912 | Hard palate, loss of less than half of. |
| 9913 | Teeth, loss of, due to loss of substance of body of maxilla or mandible. |

[29 FR 6718, May 22, 1964, as amended at 34 FR 5064, Mar. 11, 1969, 52 FR 44122, Nov. 18, 1987; 53 FR 24938, July 1, 1988]

APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

| | Diag- nos- tic Code No. |
|--|-------------------------------------|
| Abscess: | |
| Brain | 8020 |
| Kidney | 7501 |
| Liver | 7313 |
| Lung | 6809 |
| Actinomycosis, lung | 6803 |
| Addison's disease | 7911 |
| Adenitis, secondary | 7713 |
| Adenoma, thyroid: | |
| Nontoxic | 7902 |
| Toxic | 7901 |
| Adhesions: | |
| Pericardial | 7003 |
| Peritoneum | 7301 |
| Agranulocytosis | 7702 |
| Amebiasis | 7321 |
| Amputation: | |
| Arm: | |
| Disarticulation | 5120 |
| Above deltoid | 5121 |
| Below deltoid | 5122 |
| Feet, both, and hand, one | 5103 |
| Feet, both | 5107 |
| Finger (digit) individual: | |
| Thumb | 5152 |
| Index | 5153 |
| Middle | 5154 |
| Ring | 5155 |
| Little | 5156 |
| Fingers (digits) of one hand: | |
| Five | 5126 |
| Four, thumb, index, middle, ring | 5127 |
| Four, thumb, index, middle, little | 5128 |
| Four, thumb, index, ring, little | 5129 |
| Four, thumb, middle, ring, little | 5130 |
| Four, index, middle, ring, little | 5131 |
| Three, thumb, index, middle | 5132 |
| Three, thumb, index, ring | 5133 |
| Three, thumb, index, little | 5134 |
| Three, thumb, middle, ring | 5135 |
| Three, thumb, middle, little | 5136 |
| Three, thumb, ring, little | 5137 |
| Three, index, middle, ring | 5138 |
| Three, index, middle, little | 5139 |
| Three, index, ring, little | 5140 |
| Three, middle, ring, little | 5141 |
| Two, thumb, index | 5142 |

Department of Veterans Affairs

Pt. 4, App. C

| | Diagnostic Code No. | | Diagnostic Code No. |
|--|---------------------|--|---------------------|
| Two, thumb, middle | 5143 | Other types | 5009 |
| Two, thumb, ring | 5144 | Pneumococcic | 5005 |
| Two, thumb, little | 5145 | Streptococcic | 5008 |
| Two, index, middle | 5146 | Syphilitic | 5007 |
| Two, index, ring | 5147 | Traumatic | 5010 |
| Two, index, little | 5148 | Typhoid | 5006 |
| Two, middle, ring | 5149 | Aspergillosis, lung | 6807 |
| Two, middle, little | 5150 | Asthma, bronchial | 6602 |
| Two, ring, little | 5151 | Astragalectomy | 5274 |
| Forearm: | | Athetosis | 8107 |
| Above pronator teres | 5123 | Atrophy: | |
| Below pronator teres | 5124 | Muscular, progressive | 8023 |
| Forefoot | 5166 | Ovaries, both | 7620 |
| Hand, one, and foot, one | 5108 | Testis, both | 7523 |
| Hands, both, and feet, both | 5100 | Auditory canal, disease | 6210 |
| Hands, both, and foot, one | 5102 | Avitaminosis | 6313 |
| Hands, both | 5106 | Beriberi | 6314 |
| Leg: | | Blastomycosis, lung | 6805 |
| With defective stump | 5163 | Blindness, anatomical loss, one eye: | |
| With loss of natural knee action | 5164 | Other blind (5/200 or less) | 6063 |
| At lower level | 5165 | Other impaired (20/200 or less) | 6064 |
| Thigh: | | Other impaired | 6065 |
| Disarticulation | 5160 | Other normal | 6066 |
| Upper third | 5161 | Blindness, light perception only: | |
| Middle or lower thirds | 5162 | Both eyes | 6062 |
| Toe, great | 5171 | One eye: | |
| Toe, other, with removal metatarsal head | 5172 | Other blind, 5/200 or less | 6067 |
| Toes, all | 5170 | Other impaired, 20/200 or less | 6068 |
| Toes, three or more | 5173 | Other impaired | 6069 |
| Anemia: | | Other normal | 6070 |
| Pernicious | 7700 | Blindness, light perception only and loss or loss of | |
| Secondary | 7701 | use of hands and/or feet | 6050- |
| Aneurysm: | | 6061 | |
| Aorta or branches | 7110 | Blindness, total (5/200 or less): | |
| Arteriovenous, traumatic | 7113 | Both eyes | 6071 |
| Artery | 7111 | One eye: | |
| Angioneurotic edema | 7118 | Other impaired, (20/200 or less) | 6072 |
| Ankylosis: | | Other impaired | 6073 |
| Ankle | 5270 | Other normal | 6074 |
| Elbow | 5205 | Blindness, partial (20/200 or less): | |
| Finger (digit) individual: | | Both eyes | 6075 |
| Thumb | 5224 | One eye: | |
| Index | 5225 | Other impaired | 6076 |
| Middle | 5226 | Other normal | 6077 |
| Other | 5227 | Blindness, partial: | |
| Fingers (digits) of one hand, unfavorable: | | Both eyes | 6078 |
| Five | 5216 | One eye only | 6079 |
| Four | 5217 | Block, auricular ventricular | 7015 |
| Three | 5218 | Bones, Caisson disease of | 5011 |
| Two | 5219 | Bones and joints, tuberculosis of | 5001 |
| Hip | 5250 | Bronchiectasis | 6601 |
| Knee | 5256 | Bronchitis | 6600 |
| Scapulohumeral | 5200 | Buerger's disease | 7115 |
| Spine: | | Brucellosis | 6316 |
| Complete | 5286 | Bursitis | 5019 |
| Cervical | 5287 | Caisson disease | 5011 |
| Dorsal | 5288 | Calculus, bladder | 7515 |
| Lumbar | 5289 | Cataract: | |
| Subastragular or Tarsal | 5272 | Senile and others | 6028 |
| Wrist | 5214 | Traumatic | 6027 |
| Anthraxosis | 6800 | Cervicitis | 7612 |
| Aphakia | 6029 | Cholangitis | 7316 |
| Aphonia, organic | 6519 | Cholecystitis | 7314 |
| Arteriosclerosis: | | Cholelithiasis | 7315 |
| Cerebral | 8046 | Cholera, Asiatic | 6300 |
| General | 7100 | Chorea: | |
| Obliterans | 7114 | Huntington's | 8106 |
| Arteriosclerotic heart disease | 7005 | Sydenham's | 8105 |
| Arthritis: | | Choroiditis | 6005 |
| Atrophic (rheumatoid) | 5002 | Claw-foot (pes cavus) acquired | 5278 |
| Gonorrheal | 5004 | Cirrhosis of liver | 7312 |
| Hypertrophic (degenerative) | 5003 | Claudication, intermittent | 7116 |

| | Diag- nostic Code No. | | Diag- nostic Code No. |
|---|--------------------------------|--|--------------------------------|
| Coccidioidomycosis | 6821 | Depressive reaction | 9405 |
| Colitis: | | Psychoneurotic reaction, other | 9406 |
| Mucous (See Colon syndrome, irritable) | 7319 | Psychophysiologic disorders: | |
| Spastic (See Colon syndrome, irritable) | 7319 | Psychophysiologic skin reaction | 9500 |
| Ulcerative | 7323 | Psychophysiologic cardiovascular reaction | 9501 |
| Collapse, lung, permanent | 6813 | Psychophysiologic gastrointestinal reac- tion | 9502 |
| Colon syndrome, irritable | 7319 | Psychophysiologic nervous system reac- tion | 9503 |
| Congestion, lung, passive | 6817 | Psychophysiologic reaction, other | 9504 |
| Conjunctivitis: | | Psychotic disorders: | |
| Trachomatous | 6017 | Schizophrenic reaction: | |
| Other | 6018 | Simple type | 9200 |
| Coccyx | 5298 | Hebephrenic type | 9201 |
| Cushing's syndrome | 7907 | Catatonic type | 9202 |
| Cyclitis | 6004 | Paranoid type | 9203 |
| Cystitis: | | Chronic undifferentiated type | 9204 |
| Chronic | 7512 | Other | 9205 |
| Interstitial (Hunner) | 7513 | Manic depressive reaction | 9206 |
| Dacryocystitis | 6031 | Psychotic depressive reaction | 9207 |
| Deafness | | Paranoid reaction | 9208 |
| 0% evaluation based on Table VII | 6100 | Involutional psychotic reaction | 9209 |
| 10% evaluation based on Table VII | 6101 | Psychotic reaction, other | 9210 |
| 20% evaluation based on Table VII | 6102 | Distomiasis, intestinal | 7324 |
| 30% evaluation based on Table VII | 6103 | Diverticulitis, intestinal | 7327 |
| 40% evaluation based on Table VII | 6104 | Diverticulum of esophagus | 7205 |
| 50% evaluation based on Table VII | 6105 | Dupuytren's contracture—see Ankylosis, fingers. | |
| 60% evaluation based on Table VII | 6106 | Dysentery, bacillary | 7322 |
| 70% evaluation based on Table VII | 6107 | Ectropion | 6020 |
| 80% evaluation based on Table VII | 6108 | Eczema | 7806 |
| 90% evaluation based on Table VII | 6109 | Edema, angioneurotic | 7118 |
| 100% evaluation based on Table VII | 6110 | Embolism, brain | 8007 |
| Deflection, nasal septum | 6502 | Emphysema (No DC; follows DC 6602). | |
| Dermatitis, exfoliativa | 7817 | Encephalitis | 8000 |
| Dermatophytosis | 7813 | Endocarditis, bacterial, subacute | 7001 |
| Diabetes mellitus | 7913 | Enteritis | 7325 |
| Diabetes insipidus | 7909 | Enterocolitis | 7326 |
| Diaphragm, rupture | 5324 | Entropion | 6021 |
| Dilation, aneurysmal artery | 7112 | Eucleation, eye, see Blindness. | |
| Diplopia | 6090 | Epilepsy: | |
| Disease: | | Grand mal | 8910 |
| Addison's | 7911 | Petit mal | 8911 |
| Hodgkin's | 7709 | Jacksonian | 8912 |
| Morton's | 5279 | Diencephalic | 8913 |
| Raynaud's | 7117 | Psychomotor | 8914 |
| Dislocation: | | Epiphora (lacrymal duct) | 6025 |
| Cartilage, semilunar | 5258 | Erythromelalgia | 7119 |
| Lens, crystalline | 6033 | Eyelids, loss of portion of | 6032 |
| Disorders, mental: | | Fever: | |
| Organic brain disorders: | | Hemoglobinuric, see Malaria. | |
| Acute brain syndrome | 9300 | Malta | 6316 |
| Chronic brain syndrome associated with: | | Oroya | 6306 |
| Central nervous system syphilis | 9301 | Relapsing | 6308 |
| Intracranial infections other than syphilis | 9302 | Rheumatic | 6309 |
| Intoxication | 9303 | Undulant | 6316 |
| Brain trauma | 9304 | Fibrillation, auricular: | |
| Cerebral arteriosclerosis | 9305 | Paroxysmal | 7011 |
| Circulatory disturbance other than cerebral arteriosclerosis | 9306 | Permanent | 7012 |
| Convulsive disorder (idiopathic epi- lepsy) | 9307 | Filariasis | 6305 |
| Disturbance of metabolism, growth or nutrition | 9308 | Fistula: | |
| Intracranial neoplasm | 9309 | Ano | 7335 |
| Diseases of unknown or uncertain cause | 9310 | Bladder | 7516 |
| Unknown cause | 9311 | Bronchocutaneous or bronchopleural | 6812 |
| Psychoneurotic disorders: | | Intestine | 7330 |
| Anxiety reaction | 9400 | Rectovaginal | 7624 |
| Dissociative reaction | 9401 | Urethra | 7625 |
| Conversion reaction | 9402 | Flail hip | 5254 |
| Phobic reaction | 9403 | Flatfoot (pes planus) acquired | 5276 |
| Obsessive compulsive reaction | 9404 | Flutter, auricular | 7010 |
| | | Fracture, vertebra, residuals of | 5285 |
| | | Frozen feet | 7122 |
| | | Gastritis, atrophic (see DC 7307). | |
| | | Gastritis, hypertrophic | 7307 |

Department of Veterans Affairs

Pt. 4, App. C

| | Diagnostic Code No. | | Diagnostic Code No. |
|--|---------------------|---|---------------------|
| Genu, recurvatum | 5263 | Thigh, motion | 5253 |
| Glaucoma: | | Tibia and fibula | 5262 |
| Congestive | 6012 | Ulna | 5211 |
| Noncongestive | 6013 | Visual acuity, <i>see</i> Blindness. | |
| Gout | 5017 | Infarction of myocardium | 7006 |
| Growths, new benign: | | Injury: | |
| Bones, joints and muscles | 5015 | Bladder | 7517 |
| Brain | 8003 | Gall bladder | 7317 |
| Digestive system | 7344 | Eye, unhealed | 6009 |
| Ear | 6209 | Foot | 5284 |
| Endocrine system | 7915 | Larynx | 6517 |
| Eyeball and adnexa | 6015 | Lips | 7201 |
| Genitourinary system | 7529 | Liver | 7311 |
| Respiratory | 6820 | Mouth | 7200 |
| Skin | 7819 | Muscle: | |
| Spinal cord | 8022 | Facial | 5325 |
| Growths, new, malignant: | | Group I | 5301 |
| Bones | 5012 | Group II | 5302 |
| Brain | 8002 | Group III | 5303 |
| Digestive system | 7343 | Group IV | 5304 |
| Ear | 6208 | Group V | 5305 |
| Endocrine system | 7914 | Group VI | 5306 |
| Eyeball only | 6014 | Group VII | 5307 |
| Genitourinary system | 7528 | Group VIII | 5308 |
| Gynecological system or mammary glands | 7627 | Group IX | 5309 |
| Respiratory | 6819 | Group X | 5310 |
| Skin | 7818 | Group XI | 5311 |
| Spinal cord | 8021 | Group XII | 5312 |
| Hallux rigidus | 5281 | Group XIII | 5313 |
| Hallux valgus | 5280 | Group XIV | 5314 |
| Hammer toe | 5282 | Group XV | 5315 |
| Hematomyelia | 8012 | Group XVI | 5316 |
| Hemorrhage: | | Group XVII | 5317 |
| Brain | 8009 | Group XVIII | 5318 |
| Intraocular | 6007 | Group XIX | 5319 |
| Hemorrhoids | 7336 | Group XX | 5320 |
| Hepatitis, infectious | 7345 | Group XXI | 5321 |
| Hernia: | | Group XXII | 5322 |
| Femoral | 7340 | Group XXIII | 5323 |
| Hiatal | 7346 | Pleural cavity | 6818 |
| Inguinal | 7338 | Prostate | 7527 |
| Muscle | 5326 | Sacroiliac | 5294 |
| Ventral | 7339 | Spleen | 7707 |
| Hodgkin's disease | 7709 | Stomach, residuals | 7310 |
| Hydrarthrosis, intermittent | 5018 | Tongue, whole or part | 7202 |
| Hydronephrosis | 7509 | Intervertebral disc | 5293 |
| Hyperadrenia | 7910 | Iritis | 6003 |
| Hyperparathyroidism | 7904 | Kala-azar | 6301 |
| Hyperpituitarism: | | Keratitis | 6001 |
| Acromegaly or gigantism | 7908 | Labyrinthitis | 6204 |
| Cushing's syndrome | 7907 | Lagophthalmos | 6022 |
| Hypertensive heart disease | 7007 | Laryngectomy | 6518 |
| Hypertensive vascular disease | 7101 | Laryngitis | 6516 |
| Hyperthyroid heart disease | 7008 | Leishmaniasis: | |
| Hyperthyroidism | 7900 | Americana | 7807 |
| Hypoadrenia | 7911 | Old World | 7808 |
| Hypoparathyroidism | 7905 | Lens, crystalline, dislocation of | 6033 |
| Hypopituitarism | 7909 | Leprosy | 6302 |
| Hypothyroidism | 7903 | Leukemia | 7703 |
| Immersion foot | 7122 | Limitation of extension: | |
| Impairment: | | Forearm | 5207 |
| Auditory acuity, <i>see</i> Deafness. | | Leg | 5261 |
| Clavicle | 5203 | Thigh | 5251 |
| Elbow | 5209 | Limitation of field vision | 6080 |
| Eye (field vision) | 6080 | Limitation of flexion: | |
| Eye (muscle function) | 6090 | Forearm | 5206 |
| Femur | 5255 | Leg | 5260 |
| Humerus | 5202 | Thigh | 5252 |
| Knee | 5257 | Limitation of flexion and extension: | |
| Radius | 5212 | Forearm | 5208 |
| Sphincter control | 7332 | Limitation of motion: | |
| Supination and pronation | 5213 | Ankle | 5271 |

| | Diagnostic Code No. | | Diagnostic Code No. |
|--|---------------------|--|---------------------|
| Arm | 5201 | Neuralgia: | |
| Cervical | 5290 | Cranial nerves: | |
| Dorsal | 5291 | Fifth (trigeminal) | 8405 |
| Lumbar | 5292 | Seventh (facial) | 8407 |
| Temporomandibular articulation | 9905 | Ninth (glossopharyngeal) | 8409 |
| Wrist | 5215 | Tenth (pneumogastric, vagus) | 8410 |
| Limitation, pronation | 5213 | Eleventh (spinal accessory, external branch) | 8411 |
| Limitation, supination | 5213 | Twelfth (hypoglossal) | 8412 |
| Limitation of muscle function, eye | 6090 | Peripheral nerves: | |
| Lobectomy | 6816 | Upper radicular group | 8710 |
| Loss: | | Middle radicular group | 8711 |
| Auricle or deformity | 6207 | Lower radicular group | 8712 |
| Condylod process | 9908 | All radicular groups | 8713 |
| Coronoid process | 9909 | Musculospiral | 8714 |
| Eyebrows | 6023 | Median | 8715 |
| Eyelashes | 6024 | Ulnar | 8716 |
| Mandible: | | Musculocutaneous | 8717 |
| Complete | 9901 | Circumflex | 8718 |
| One-half | 9902 | Long thoracic | 8719 |
| Maxilla | 9910 | Sciatic | 8720 |
| Teeth | 9913 | External popliteal | 8721 |
| Nose, loss of part, or scars | 6504 | Musculocutaneous (superficial peroneal) | 8722 |
| Palate, hard: | | Anterior tibial | 8723 |
| Half or more | 9911 | Internal popliteal | 8724 |
| Less than half | 9912 | Posterior tibial | 8725 |
| Ramus: | | Anterior crural | 8726 |
| Less than one-half substance | 9907 | Internal saphenous | 8727 |
| Whole or part | 9906 | Obturator | 8728 |
| Skull, part | 5296 | External cutaneous, thigh | 8729 |
| Smell, sense of | 6275 | Ilio-inguinal | 8730 |
| Taste, sense of | 6276 | Neuritis, optic | 6026 |
| Tongue or part | 7202 | Neuritis: | |
| Others, see Amputation, removal, etc. | | Cranial nerves: | |
| Loss of use: | | Fifth (trigeminal) | 8305 |
| Feet, both | 5110 | Seventh (facial) | 8307 |
| Feet, both, and hand, one | 5105 | Ninth (glossopharyngeal) | 8309 |
| Foot, one | 5167 | Tenth (pneumogastric, vagus) | 8310 |
| Hand, one | 5125 | Eleventh (spinal accessory, external branch) | 8311 |
| Hand, one, and foot, one | 5111 | Twelfth (hypoglossal) | 8312 |
| Hands, both, and feet, both | 5101 | Peripheral: | |
| Hands, both, and foot, one | 5104 | Upper radicular group | 8610 |
| Hands, both | 5109 | Middle radicular group | 8611 |
| Lupus, erythematosus, discoid | 7809 | Lower radicular group | 8612 |
| Lupus, erythematosus systemic (disseminated) | 6350 | All radicular groups | 8613 |
| Lupus, vulgaris | 7811 | Musculospiral | 8614 |
| Lymphogranulomatosis | 7709 | Median | 8615 |
| Malaria | 6304 | Ulnar | 8616 |
| Malunion: | | Musculocutaneous | 8617 |
| Clavicle | 5203 | Circumflex | 8618 |
| Os calcis (or astragalus) | 5273 | Long thoracic | 8619 |
| Mandible | 9904 | Sciatic | 8620 |
| Maxilla (or nonunion) | 9910 | External popliteal | 8621 |
| Scapula | 5203 | Musculocutaneous (superficial peroneal) | 8622 |
| Tarsal or metatarsal (or nonunion) | 5283 | Anterior tibial | 8623 |
| Others, see Impairment. | | Internal popliteal | 8624 |
| Mastoiditis | 6206 | Posterior tibial | 8625 |
| Meniere's disease | 6205 | Anterior crural | 8626 |
| Meningitis, cerebrospinal | 8019 | Internal saphenous | 8627 |
| Mental disorders—see Disorders, mental. | | Obturator | 8628 |
| Metatarsalgia | 5279 | External cutaneous, thigh | 8629 |
| Metritis | 7613 | Ilio-inguinal | 8630 |
| Migraine | 8100 | Non-union of bones: | |
| Muscle injury, see Injury, muscle. | | Mandible | 9903 |
| Myasthenia gravis | 8025 | Radius and Ulna | 5210 |
| Mycosis, lung, unspecified | 6808 | Tibia and fibula | 5262 |
| Myelitis | 8010 | Others, see Impairment. | |
| Myositis | 5021 | Nystagmus, central | 6016 |
| Myositis ossificans | 5023 | Oophoritis | 7615 |
| Narcolepsy | 8108 | Oroya fever | 6306 |
| Nephritis, chronic | 7502 | Osteitis deformans | 5016 |
| Nephrolithiasis | 7508 | | |
| Nephrosclerosis, arteriolar | 7507 | | |

Department of Veterans Affairs

Pt. 4, App. C

| | Diagnostic Code No. | | Diagnostic Code No. |
|--|---------------------|---|---------------------|
| Osteomalacia | 5014 | Psoriasis | 7816 |
| Osteomyelitis, jaw | 9900 | Psychiatric disorders, see Disorders, mental. | |
| Osteomyelitis | 5000 | Pterygium | 6034 |
| Osteoporosis | 5013 | Ptosis, eyelid | 6019 |
| Otitis externa | 6210 | Purpura, hemorrhagica | 7705 |
| Otitis interna | 6203 | Pyelitis | 7503 |
| Otitis media: | | Pyelonephritis, chronic | 7504 |
| Catarrhal | 6201 | Raynaud's disease | 7117 |
| Suppurative | 6200 | Removal: | |
| Otosclerosis | 6202 | Auricle or deformity | 6207 |
| Palsy, bulbar | 8005 | Cartilage, semilunar | 5259 |
| Paralysis: | | Coccyx | 5298 |
| Accommodation | 6030 | Gall bladder | 7318 |
| Agitans | 8004 | Kidney | 7500 |
| Paralysis, nerve: | | Mammary glands | 7626 |
| Cranial: | | Ovaries, both | 7619 |
| Fifth (trigeminal) | 8205 | Penis, half or more | 7520 |
| Seventh (facial) | 8207 | Penis, glans | 7521 |
| Ninth (glossopharyngeal) | 8209 | Prostate, or resection | 7526 |
| Tenth (pneumogastric, vagus) | 8210 | Ribs | 5297 |
| Eleventh (spiral accessory, external branch) | | Testis | 7524 |
| Twelfth (hypoglossal) | 8212 | Uterus | 7618 |
| Peripheral: | | Uterus and ovaries | 7617 |
| Upper radicular group | 8510 | Others, see Amputation, loss, etc. | |
| Middle radicular group | 8511 | Resection: | |
| Lower radicular group | 8512 | Intestine: | |
| All radicular groups | 8513 | Large | 7329 |
| Musculospiral | 8514 | Small | 7328 |
| Median | 8515 | Stomach | 7308 |
| Ulnar | 8516 | Retina, detachment of | 6008 |
| Musculocutaneous | 8517 | Retinitis | 6006 |
| Circumflex | 8518 | Rheumatic fever | 6309 |
| Long thoracic | 8519 | Rheumatic heart disease | 7000 |
| Sciatic | 8520 | Rhinitis: Atrophic | 6501 |
| External popliteal | 8521 | Rupture, diaphragm | 5324 |
| Musculocutaneous (superficial peroneal) | 8522 | Salpingitis | 7614 |
| Anterior tibial | 8523 | Scars: | |
| Internal popliteal | 8524 | Burns, second degree | 7802 |
| Posterior tibial | 8525 | Burns, third degree | 7801 |
| Anterior crural | 8526 | Head, etc., disfiguring | 7800 |
| Internal saphenous | 8527 | Retina | 6011 |
| Obturator | 8528 | Superficial, tender | 7804 |
| External cutaneous, thigh | 8529 | Superficial, with ulceration | 7803 |
| Ilio-inguinal | 8530 | Others | 7805 |
| Paramyoclonus multiplex | 8104 | Scleritis | 6002 |
| Pellagra | 6315 | Sclerosis: | |
| Pemphigus | 7815 | Amyotrophic, lateral | 8017 |
| Penis, deformity of | 7522 | Multiple | 8018 |
| Perforation: Tympanic membrane | 6211 | Scotoma, pathological | 6081 |
| Pericarditis | 7002 | Shortening, leg | 5275 |
| Periostitis | 5022 | Silicosis | 6801 |
| Pes cavus | 5278 | Sinusitis: | |
| Pes planus | 5276 | Ethmoid | 6511 |
| Phlebitis | 7121 | Frontal | 6512 |
| Pinta | 7810 | Maxillary | 6513 |
| Plague | 6307 | Pansinusitis | 6510 |
| Pleurisy: | | Sphenoid | 6514 |
| Purulent (empyema) | 6811 | Spasm, esophagus | 7204 |
| Serofibrinous | 6810 | Splenectomy | 7706 |
| Pluriglandular syndrome | 7912 | Sporotrichosis, lung | 6806 |
| Pneumoconiosis | 6802 | Stenosis: | |
| Pneumonectomy | 6815 | Larynx | 6520 |
| Pneumothorax, spontaneous | 6814 | Stomach | 7309 |
| Poliomyelitis, anterior | 8011 | Strain, lumbosacral | 5295 |
| Polycythemia | 7704 | Streptotrichosis, lung | 6804 |
| Pregnancy, surgical complications of | 7623 | Stricture: | |
| Prolapse: | | Esophagus | 7203 |
| Rectum | 7334 | Rectum, anus | 7333 |
| Uterus | 7621 | Ureter | 7511 |
| Pronation, limitation of | 5213 | Urethra | 7518 |
| Pruritis, ani | 7337 | Supination, limitation of | 5213 |
| | | Symblepharon | 6091 |

| | Diagnostic Code No. |
|---|---------------------|
| Syndrome: | |
| Cushing's | 7907 |
| Intervertebral disc | 5293 |
| Meniere's | 6205 |
| Pluriglandular | 7912 |
| Postgastrectomy | 7308 |
| Synovitis | 5020 |
| Syphilis: | |
| Cerebrospinal | 8013 |
| Meningovascular | 8014 |
| Unspecified | 6310 |
| Syphilitic heart disease | 7004 |
| Syringomyelia | 8024 |
| Tabes dorsalis | 8015 |
| Tachycardia: | |
| Paroxysmal | 7013 |
| Sinus | 7014 |
| Tenosynovitis | 5024 |
| Thrombo-anglitis obliterans | 7115 |
| Thrombophlebitis | 7121 |
| Thrombosis, brain | 8008 |
| Tic, convulsive | 8103 |
| Tinea barbae | 7814 |
| Tinnitus | 6260 |
| Tuberculosis: | |
| Adenitis, tuberculous: | |
| Axillary | 7711 |
| Cervical | 7710 |
| Inguinal | 7712 |
| Bladder | 7514 |
| Bones and joints | 5001 |
| Epididymo-orchitis, tuberculous | 7525 |
| Eye | 6010 |
| Kidney | 7505 |
| Laryngitis, tuberculous | 6515 |
| Luposa | 7811 |
| Miliary | 6311 |
| Nonpulmonary, inactive (see § 4.89) | |
| Peritonitis, tuberculous | 7331 |
| Pleurisy, tuberculous | 6732 |
| Pulmonary: | |
| Active: | |
| Far advanced | 6701 & 6707 |
| Moderately advanced | 6702 & 6708 |
| Minimal | 6703 & 6709 |
| Advancement unspecified | 6704 & 6710 |
| Inactive: | |
| Far advanced | 6721 & 6725 |
| Moderately advanced | 6722 & 6726 |
| Minimal | 6723 & 6727 |
| Advancement unspecified | 6724 & 6728 |
| Tympanic membrane, perforation of | 6211 |
| Typhus, scrub | 6317 |
| Ulcer: | |
| Duodenal | 7305 |
| Gastric | 7304 |
| Marginal | 7306 |
| Undescended testis (see Note under DC 7524) | |
| Uterus, displacement of | 7622 |
| Ureterolithiasis | 7510 |
| Uveitis | 6000 |
| Vaginitis | 7611 |
| Varicose veins | 7120 |
| Verruga peruana | 7812 |

| | Diagnostic Code No. |
|---------------------------------------|---------------------|
| Vertebra, fracture | 5285 |
| Visceroptosis | 7342 |
| Vision, impairment of, see Blindness. | |
| Vulvovaginitis | 7610 |
| Weak foot | 5277 |
| Wound, incised, abdominal wall | 7341 |

[29 FR 6718, May 22, 1964, as amended at 34 FR 5064, Mar. 11, 1969; 52 FR 44122, Nov. 18, 1987; 53 FR 24938, July 1, 1988]

PART 5 [RESERVED]

PART 6—UNITED STATES GOVERNMENT LIFE INSURANCE

AGE

Sec.

6.1 Misstatement of age.

PREMIUMS

6.2 Premium rate.

POLICIES

6.3 Incontestability of United States Government life insurance.

BENEFICIARY OF UNITED STATES GOVERNMENT LIFE INSURANCE

6.4 Proof of age, relationship and marriage.

6.5 Conditional designation of beneficiary.

6.6 Change of beneficiary.

6.7 Claims of creditors, taxation.

OPTIONAL SETTLEMENT

6.8 Selection, revocation and election.

6.9 Election of optional settlement by beneficiary.

6.10 Options.

DIVIDENDS

6.11 How dividends are paid.

6.12 Special dividends.

LOANS

6.13 Policy loans.

CASH VALUE

6.14 Cash value; other than special endowment at age 96 plan policy.

6.15 Cash value; special endowment at age 96 plan policy.

6.16 Payment of cash value in monthly installments.

INDEBTEDNESS

6.17 Collection of any indebtedness.