include an attempt to obtain additional evidence or the holding of an informal conference with the claimant. Upon the request of the claimant, the reviewer will conduct a hearing under §3.103(c).

(d) The reviewer may grant a benefit sought in the claim notwithstanding §3.105(b), but, except as provided in paragraph (e) of this section, may not revise the decision in a manner that is less advantageous to the claimant than the decision under review. A review decision made under this section will include a summary of the evidence, a citation to pertinent laws, a discussion of how those laws affect the decision, and a summary of the reasons for the decision.

(e) Notwithstanding any other provisions of this section, the reviewer may reverse or revise (even if disadvantageous to the claimant) prior decisions of an agency of original jurisdiction (including the decision being reviewed or any prior decision that has become final due to failure to timely appeal) on the grounds of clear and unmistakable error (see §3.105(a)).

(f) Review under this section does not limit the appeal rights of a claimant. Unless a claimant withdraws his or her Notice of Disagreement as a result of this review process, VA will proceed with the traditional appellate process by issuing a Statement of the Case.

(g) This section applies to all claims in which a Notice of Disagreement is filed on or after June 1, 2001.

(Authority: 38 U.S.C. 5109A and 7105(d))

[66 FR 21874, May 2, 2001, as amended at 67 FR 46868, July 17, 2002]

## PART 4—SCHEDULE FOR RATING **DISABILITIES**

## Subpart A—General Policy in Rating

## Sec.

- 4.1 Essentials of evaluative rating.
- 4.2 Interpretation of examination reports.
- 4.3 Resolution of reasonable doubt.
- Evaluation of evidence.
- Higher of two evaluations.
- 4.9 Congenital or developmental defects.
- 4.10 Functional impairment.
- 4.13 Effect of change of diagnosis.
- 4.14 Avoidance of pyramiding.
- 4.15 Total disability ratings.

- 4.16 Total disability ratings for compensation based on unemployability of the individual.
- 4.17 Total disability ratings for pension based on unemployability and age of the individual.
- 4.17a Misconduct etiology.
- 4.18 Unemployability.
- Age in service-connected claims. 4.19
- Analogous ratings. 4.20
- Application of rating schedule. 4.21
- Rating of disabilities aggravated by ac-4.22 tive service.
- 4.23 Attitude of rating officers.
- 4.24 Correspondence.
- 4.25 Combined ratings table.
- Bilateral factor.
- Use of diagnostic code numbers.
- Prestabilization rating from date of discharge from service.
- 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.
- 4.30 Convalescent ratings.
- 4.31 Zero percent evaluations.

### Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

- 4.40 Functional loss
- History of injury
- 4.42 Complete medical examination of injury cases.
- 4.43 Osteomyelitis.
- 4 44 The bones.
- 4.45 The joints.
- 4.46 Accurate measurement.
- 4.47-4.54 [Reserved]
- 4.55 Principles of combined ratings for muscle injuries.
- 4.56 Evaluation of muscle disabilities.
- 4.57 Static foot deformities. 4.58 Arthritis due to strain.
- Painful motion.
- 4.60 [Reserved]
- Examination. 4.61 4.62 Circulatory disturbances.
- Loss of use of hand or foot
- Loss of use of both buttocks.
- [Reserved]
- 4.66 Sacroiliac joint.
- Pelvic bones.
- 4.68 Amputation rule.
- 4.69 Dominant hand.
- 4.70 Inadequate examinations.
- Measurement of ankylosis and joint motion.
- 4.71a Schedule of ratings-musculoskeletal system.
- 4.72 [Reserved]
- 4.73 Schedule of ratings—muscle injuries.

## THE ORGANS OF SPECIAL SENSE

- 4.75 Examination of visual acuity.
- 4.76 Examination of field vision.
- 4.76a Computation of average concentric contraction of visual fields.

### 38 CFR Ch. I (7-1-07 Edition)

#### §4.1

- 4.77 Examination of muscle function.
- 4.78 Computing aggravation.
- 4.79 Loss of use of one eye, having only light perception.
- 4.80 Rating of one eye.
- 4.81-4.82 [Reserved]
- 4.83 Ratings at scheduled steps and distances.
- 4.83a Impairment of central visual acuity.
- 4.84 Differences between distant and near visual acuity.
- 4.84a Schedule of ratings—eye.

#### IMPAIRMENT OF AUDITORY ACUITY

- 4.85 Evaluation of hearing impairment.
- 4.86 Exceptional patterns of hearing impairment.
- 4.87 Schedule of ratings—ear.
- 4.87a Schedule of ratings—other sense organs.

# INFECTIOUS DISEASES, IMMUNE DISORDERS AND NUTRITIONAL DEFICIENCIES

- 4.88 [Reserved]
- 4.88a Chronic fatigue syndrome.
- 4.88b Schedule of ratings—infectious diseases, immune disorders and nutritional deficiencies.
- 4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19 1968
- 4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

## THE RESPIRATORY SYSTEM

- 4.96 Special provisions regarding evaluation of respiratory conditions.
- 4.97 Schedule of ratings—respiratory system.

## THE CARDIOVASCULAR SYSTEM

- 4.100 Application of the evaluation criteria for diagnostic codes 7000–7007, 7011, and 7015–7020.
- 4.101-4.103 [Reserved]
- 4.104 Schedule of ratings—cardiovascular system.

### THE DIGESTIVE SYSTEM

- 4.110 Ulcers.
- 4.111 Postgastrectomy syndromes.
- 4.112 Weight loss.
- 4.113 Coexisting abdominal conditions.
- 4.114 Schedule of ratings—digestive system.

#### THE GENITOURINARY SYSTEM

- 4.115 Nephritis.
- 4.115a Ratings of the genitourinary system—dysfunctions.
- 4.115b Ratings of the genitourinary system—diagnoses.

GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

#### THE HEMIC AND LYMPHATIC SYSTEMS

4.117 Schedule of ratings—hemic and lymphatic systems.

#### THE SKIN

4.118 Schedule of ratings-skin.

#### THE ENDOCRINE SYSTEM

4.119 Schedule of ratings—endocrine system.

# NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

- 4.120 Evaluations by comparison.
- 4.121 Identification of epilepsy.
- 4.122 Psychomotor epilepsy.
- 4.123 Neuritis, cranial or peripheral.
- 4.124 Neuralgia, cranial or peripheral.
- 4.124a Schedule of ratings—neurological conditions and convulsive disorders.

## MENTAL DISORDERS

- 4.125 Diagnosis of mental disorders.
- 4.126 Evaluation of disability from mental disorders.
- 4.127 Mental retardation and personality disorders.
- 4.128 Convalescence ratings following extended hospitalization.
- 4.129 Mental disorders due to traumatic stress.
- 4.130 Schedule of ratings-mental disorders.

#### DENTAL AND ORAL CONDITIONS

- 4.149 [Reserved]
- 4.150 Schedule of ratings—dental and oral conditions.

APPENDIX A TO PART 4—TABLE OF AMEND-MENTS AND EFFECTIVE DATES SINCE 1946

APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES

APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

AUTHORITY:  $38\ U.S.C.\ 1155,\ unless\ otherwise\ noted.$ 

Source:  $29 \, \mathrm{FR} \, 6718$ , May 22, 1964, unless otherwise noted.

# Subpart A—General Policy in Rating

## § 4.1 Essentials of evaluative rating.

This rating schedule is primarily a guide in the evaluation of disability resulting from all types of diseases and injuries encountered as a result of or

incident to military service. The percentage ratings represent as far as can practicably be determined the average impairment in earning capacity resulting from such diseases and injuries and their residual conditions in civil occupations. Generally, the degrees of disability specified are considered adequate to compensate for considerable loss of working time from exacerbations or illnesses proportionate to the severity of the several grades of disability. For the application of this schedule, accurate and fully descriptive medical examinations are required, with emphasis upon the limitation of activity imposed by the disabling condition. Over a period of many years, a veteran's disability claim may require reratings in accordance with changes in laws, medical knowledge and his or her physical or mental condition. It is thus essential, both in the examination and in the evaluation of disability, that each disability be viewed in relation to its history.

[41 FR 11292, Mar. 18, 1976]

# § 4.2 Interpretation of examination reports.

Different examiners, at different times, will not describe the same disability in the same language. Features of the disability which must have persisted unchanged may be overlooked or a change for the better or worse may not be accurately appreciated or described. It is the responsibility of the rating specialist to interpret reports of examination in the light of the whole recorded history, reconciling the various reports into a consistent picture so that the current rating may accurately reflect the elements of disability present. Each disability must be considered from the point of view of the veteran working or seeking work. If a diagnosis is not supported by the findings on the examination report or if the report does not contain sufficient detail, it is incumbent upon the rating board to return the report as inadequate for evaluation purposes.

[41 FR 11292, Mar. 18, 1976]

## §4.3 Resolution of reasonable doubt.

It is the defined and consistently applied policy of the Department of Vet-

erans Affairs to administer the law under a broad interpretation, consistent, however, with the facts shown in every case. When after careful consideration of all procurable and assembled data, a reasonable doubt arises regarding the degree of disability such doubt will be resolved in favor of the claimant. See §3.102 of this chapter.

[40 FR 42535, Sept. 15, 1975]

#### § 4.6 Evaluation of evidence.

The element of the weight to be accorded the character of the veteran's service is but one factor entering into the considerations of the rating boards in arriving at determinations of the evaluation of disability. Every element in any way affecting the probative value to be assigned to the evidence in each individual claim must be thoroughly and conscientiously studied by each member of the rating board in the light of the established policies of the Department of Veterans Affairs to the end that decisions will be equitable and just as contemplated by the requirements of the law.

### § 4.7 Higher of two evaluations.

Where there is a question as to which of two evaluations shall be applied, the higher evaluation will be assigned if the disability picture more nearly approximates the criteria required for that rating. Otherwise, the lower rating will be assigned.

# § 4.9 Congenital or developmental defects.

Mere congenital or developmental defects, absent, displaced or supernumerary parts, refractive error of the eye, personality disorder and mental deficiency are not diseases or injuries in the meaning of applicable legislation for disability compensation purposes.

[41 FR 11292, Mar. 18, 1976]

## §4.10 Functional impairment.

The basis of disability evaluations is the ability of the body as a whole, or of the psyche, or of a system or organ of the body to function under the ordinary conditions of daily life including employment. Whether the upper or

lower extremities, the back or abdominal wall, the eyes or ears, or the cardiovascular, digestive, or other system, or psyche are affected, evaluations are based upon lack of usefulness, of these parts or systems, especially in self-support. This imposes upon the medical examiner the responsibility of furnishing, in addition to the etiological, anatomical, pathological, laboratory and prognostic data required for ordinary medical classification, full description of the effects of disability upon the person's ordinary activity. In this connection, it will be remembered that a person may be too disabled to engage in employment although he or she is up and about and fairly comfortable at home or upon limited activity.

[41 FR 11292, Mar. 18, 1976]

## § 4.13 Effect of change of diagnosis.

The repercussion upon a current rating of service connection when change is made of a previously assigned diagnosis or etiology must be kept in mind. The aim should be the reconciliation and continuance of the diagnosis or etiology upon which service connection for the disability had been granted. The relevant principle enunciated in §4.125, entitled "Diagnosis of mental disorders," should have careful attention in this connection. When any change in evaluation is to be made, the rating agency should assure itself that there has been an actual change in the conditions, for better or worse, and not merely a difference in thoroughness of the examination or in use of descriptive terms. This will not, of course, preclude the correction of erroneous ratings, nor will it preclude assignment of a rating in conformity with §4.7.

 $[29\ FR\ 6718,\ May\ 22,\ 1964,\ as\ amended\ at\ 61\ FR\ 52700,\ Oct.\ 8,\ 1996]$ 

### §4.14 Avoidance of pyramiding.

The evaluation of the same disability under various diagnoses is to be avoided. Disability from injuries to the muscles, nerves, and joints of an extremity may overlap to a great extent, so that special rules are included in the appropriate bodily system for their evaluation. Dyspnea, tachycardia, nervousness, fatigability, etc., may result from

many causes; some may be service connected, others, not. Both the use of manifestations not resulting from service-connected disease or injury in establishing the service-connected evaluation, and the evaluation of the same manifestation under different diagnoses are to be avoided.

## §4.15 Total disability ratings.

The ability to overcome the handicap of disability varies widely among individuals. The rating, however, is based primarily upon the average impairment in earning capacity, that is, upon the economic or industrial handicap which must be overcome and not from individual success in overcoming it. However, full consideration must be given to unusual physical or mental effects in individual cases, to peculiar effects of occupational activities, to defects in physical or mental endowment preventing the usual amount of success in overcoming the handicap of disability and to the effect of combinations of disability. Total disability will be considered to exist when there is present any impairment of mind or body which is sufficient to render it impossible for the average person to follow a substantially gainful occupation; *Provided*, That permanent total disability shall be taken to exist when the impairment is reasonably certain to continue throughout the life of the disabled person. The following will be considered to be permanent total disability: the permanent loss of the use of both hands, or of both feet, or of one hand and one foot, or of the sight of both eyes, or becoming permanently helpless or permanently bedridden. Other total disability ratings are scheduled in the various bodily systems of this schedule.

#### § 4.16 Total disability ratings for compensation based on unemployability of the individual.

(a) Total disability ratings for compensation may be assigned, where the schedular rating is less than total, when the disabled person is, in the judgment of the rating agency, unable to secure or follow a substantially gainful occupation as a result of service-connected disabilities: *Provided* 

That, if there is only one such disability, this disability shall be ratable at 60 percent or more, and that, if there are two or more disabilities, there shall be at least one disability ratable at 40 percent or more, and sufficient additional disability to bring the combined rating to 70 percent or more. For the above purpose of one 60 percent disability, or one 40 percent disability in combination, the following will be considered as one disability: (1) Disabilities of one or both upper extremities, or of one or both lower extremities, including the bilateral factor, if applicable, (2) disabilities resulting from common etiology or a single accident, (3) disabilities affecting a single body system, e.g. orthopedic, digestive, rescardiovascular-renal, neuropsychiatric, (4) multiple injuries incurred in action, or (5) multiple disabilities incurred as a prisoner of war. It is provided further that the existence or degree of nonservice-connected previous disabilities or unemployability status will be dis-regarded where the percentages referred to in this paragraph for the service-connected disability or disabilities are met and in the judgment of the rating agency such service-connected disabilities render the veteran unemployable. Marginal employment shall not be considered substantially gainful employment. For purposes of this section, marginal employment generally shall be deemed to exist when a veteran's earned annual income does not exceed the amount established by the U.S. Department of Commerce, Bureau of the Census, as the poverty threshold for one person. Marginal employment may also be held to exist, on a facts found basis (includes but is not limited to employment in a protected environment such as a family business or sheltered workshop), when earned annual income exceeds the poverty threshold. Consideration shall be given in all claims to the nature of the employment and the reason for termination.

(Authority: 38 U.S.C. 501)

(b) It is the established policy of the Department of Veterans Affairs that all veterans who are unable to secure and follow a substantially gainful occupation by reason of service-connected

disabilities shall be rated totally disabled. Therefore, rating boards should submit to the Director, Compensation and Pension Service, for extra-schedular consideration all cases of veterans who are unemployable by reason of service-connected disabilities, but who fail to meet the percentage standards set forth in paragraph (a) of this section. The rating board will include a full statement as to the veteran's service-connected disabilities, employment history, educational and vocational attainment and all other factors having a bearing on the issue.

[40 FR 42535, Sept. 15, 1975, as amended at 54 FR 4281, Jan. 30, 1989; 55 FR 31580, Aug. 3, 1990; 58 FR 39664, July 26, 1993; 61 FR 52700, Oct. 8, 1996]

#### §4.17 Total disability ratings for pension based on unemployability and age of the individual.

All veterans who are basically eligible and who are unable to secure and follow a substantially gainful occupation by reason of disabilities which are likely to be permanent shall be rated as permanently and totally disabled. For the purpose of pension, the permanence of the percentage requirements of §4.16 is a requisite. When the percentage requirements are met, and the disabilities involved are of a permanent nature, a rating of permanent and total disability will be assigned if the veteran is found to be unable to secure and follow substantially gainful employment by reason of such disability. Prior employment or unemployment status is immaterial if in the judgment of the rating board the veteran's disabilities render him or her unemployable. In making such determinations, the following guidelines will be used:

(a) Marginal employment, for example, as a self-employed farmer or other person, while employed in his or her own business, or at odd jobs or while employed at less than half the usual remuneration will not be considered incompatible with a determination of unemployability, if the restriction, as to securing or retaining better employment, is due to disability.

(b) Claims of all veterans who fail to meet the percentage standards but who meet the basic entitlement criteria and are unemployable, will be referred by

### §4.17a

the rating board to the Veterans Service Center Manager under §3.321(b)(2) of this chapter.

(Authority: 38 U.S.C. 1155; 38 U.S.C. 3102)

[43 FR 45348, Oct. 2, 1978, as amended at 56 FR 57985, Nov. 15, 1991; 71 FR 28586, May 17, 2006]

#### § 4.17a Misconduct etiology.

A permanent and total disability rating under the provisions of §§ 4.15, 4.16 and 4.17 will not be precluded by reason of the coexistence of misconduct disability when:

(a) A veteran, regardless of employment status, also has innocently acquired 100 percent disability, or

(b) Where unemployable, the veteran has other disabilities innocently acquired which meet the percentage requirements of §§ 4.16 and 4.17 and would render, in the judgment of the rating agency, the average person unable to secure or follow a substantially gainful occupation.

[40 FR 42536, Sept. 15, 1975, as amended at 43 FR 45349, Oct. 2, 1978]

#### §4.18 Unemployability.

A veteran may be considered as unemployable upon termination of employment which was provided on account of disability, or in which special consideration was given on account of the same, when it is satisfactorily shown that he or she is unable to secure further employment. With amputations, sequelae of fractures and other residuals of traumatism shown to be of static character, a showing of continuous unemployability from date of incurrence, or the date the condition reached the stabilized level, is a general requirement in order to establish the fact that present unemployability is the result of the disability. However, consideration is to be given to the circumstances of employment in individual claims, and, if the employment was only occasional, intermittent, tryout or unsuccessful, or eventually terminated on account of the disability, present unemployability may be attributed to the static disability. Where unemployability for pension previously has been established on the basis of combined service-connected and nonservice-connected disabilities and the service-connected disability or disabilities have increased in severity, §4.16 is for consideration.

[40 FR 42536, Sept. 15, 1975, as amended at 43 FR 45349, Oct. 2, 1978]

#### § 4.19 Age in service-connected claims.

Age may not be considered as a factor in evaluating service-connected disability; and unemployability, in service-connected claims, associated with advancing age or intercurrent disability, may not be used as a basis for a total disability rating. Age, as such, is a factor only in evaluations of disability not resulting from service, i.e., for the purposes of pension.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

## § 4.20 Analogous ratings.

When an unlisted condition is encountered it will be permissible to rate under a closely related disease or injury in which not only the functions affected, but the anatomical localization and symptomatology are closely analogous. Conjectural analogies will be avoided, as will the use of analogous ratings for conditions of doubtful diagnosis, or for those not fully supported by clinical and laboratory findings. Nor will ratings assigned to organic diseases and injuries be assigned by analogy to conditions of functional origin.

#### §4.21 Application of rating schedule.

In view of the number of atypical instances it is not expected, especially with the more fully described grades of disabilities, that all cases will show all the findings specified. Findings sufficiently characteristic to identify the disease and the disability therefrom, and above all, coordination of rating with impairment of function will, however, be expected in all instances.

[41 FR 11293, Mar. 18, 1976]

# § 4.22 Rating of disabilities aggravated by active service.

In cases involving aggravation by active service, the rating will reflect only the degree of disability over and above the degree existing at the time of entrance into the active service, whether the particular condition was noted at the time of entrance into the active service, or it is determined upon the

evidence of record to have existed at that time. It is necessary therefore, in all cases of this character to deduct from the present degree of disability the degree, if ascertainable, of the disability existing at the time of entrance into active service, in terms of the rating schedule, except that if the disability is total (100 percent) no deduction will be made. The resulting difference will be recorded on the rating sheet. If the degree of disability at the time of entrance into the service is not ascertainable in terms of the schedule, no deduction will be made.

#### § 4.23 Attitude of rating officers.

It is to be remembered that the majority of applicants are disabled persons who are seeking benefits of law to which they believe themselves entitled. In the exercise of his or her functions, rating officers must not allow their personal feelings to intrude; an antagonistic, critical, or even abusive attitude on the part of a claimant should not in any instance influence the officers in the handling of the case. Fairness and courtesy must at all times be shown to applicants by all employees whose duties bring them in contact, directly or indirectly, with the Department's claimants.

[41 FR 11292, Mar. 18, 1976]

#### § 4.24 Correspondence.

All correspondence relative to the interpretation of the schedule for rating disabilities, requests for advisory opinions, questions regarding lack of clarity or application to individual cases involving unusual difficulties, will be addressed to the Director, Compensation and Pension Service. A clear statement will be made of the point or points upon which information is desired, and the complete case file will be simultaneously forwarded to Central Office. Rating agencies will assure themselves that the recent report of physical examination presents an adequate picture of the claimant's condition. Claims in regard to which the schedule evaluations are considered inadequate or excessive, and errors in the schedule will be similarly brought to attention.

[41 FR 11292, Mar. 18, 1976]

#### §4.25 Combined ratings table.

Table I, Combined Ratings Table, results from the consideration of the efficiency of the individual as affected first by the most disabling condition, then by the less disabling condition, then by other less disabling conditions, if any, in the order of severity. Thus, a person having a 60 percent disability is considered 40 percent efficient. Proceeding from this 40 percent efficiency, the effect of a further 30 percent disability is to leave only 70 percent of the efficiency remaining after consideration of the first disability, or 28 percent efficiency altogether. The individual is thus 72 percent disabled, as shown in table I opposite 60 percent and under 30 percent.

(a) To use table I, the disabilities will first be arranged in the exact order of their severity, beginning with the greatest disability and then combined with use of table I as hereinafter indicated. For example, if there are two disabilities, the degree of one disability will be read in the left column and the degree of the other in the top row, whichever is appropriate. The figures appearing in the space where the column and row intersect will represent the combined value of the two. This combined value will then be converted to the nearest number divisible by 10, and combined values ending in 5 will be adjusted upward. Thus, with a 50 percent disability and a 30 percent disability, the combined value will be found to be 65 percent, but the 65 percent must be converted to 70 percent to represent the final degree of disability. Similarly, with a disability of 40 percent, and another disability of 20 percent, the combined value is found to be 52 percent, but the 52 percent must be converted to the nearest degree divisible by 10, which is 50 percent. If there are more than two disabilities, the disabilities will also be arranged in the exact order of their severity and the combined value for the first two will be found as previously described for two disabilities. The combined value, exactly as found in table I, will be combined with the degree of the third disability (in order of severity). The combined value for the three disabilities will be found in the space where the column and row intersect, and if there

are only three disabilities will be converted to the nearest degree divisible by 10, adjusting final 5's upward. Thus, if there are three disabilities ratable at 60 percent, 40 percent, and 20 percent, respectively, the combined value for the first two will be found opposite 60 and under 40 and is 76 percent. This 76 will be combined with 20 and the combined value for the three is 81 percent. This combined value will be converted to the nearest degree divisible by 10 which is 80 percent. The same procedure will be employed when there are four or more disabilities. (See table I).

(b) Except as otherwise provided in this schedule, the disabilities arising from a single disease entity, e.g., arthritis, multiple sclerosis, cerebrovascular accident, etc., are to be rated separately as are all other disabiling conditions, if any. All disabilities are then to be combined as described in paragraph (a) of this section. The conversion to the nearest degree divisible by 10 will be done only once per rating decision, will follow the combining of all disabilities, and will be the last procedure in determining the combined degree of disability.

TABLE I—COMBINED RATINGS TABLE
[10 combined with 10 is 19]

	[10 COMMINION WITH TO IS 19]									
		10	20	30	40	50	60	70	80	90
19		27	35	43	51	60	68	76	84	92
		28	36	44	52	60	68	76	84	92
21 .		29	37	45	53	61	68	76	84	92
22 .		30	38	45	53	61	69	77	84	92
23 .		31	38	46	54	62	69	77	85	92
24 .		32	39	47	54	62	70	77	85	92
25 .		33	40	48	55	63	70	78	85	93
26 .		33	41	48	56	63	70	78	85	93
		34	42	49	56	64	71	78	85	93
		35	42	50	57	64	71	78	86	93
		36	43	50	57	65	72	79	86	93
		37	44	51	58	65	72	79	86	93
		38	45	52	59	66	72	79	86	93
		39	46	52	59	66	73	80	86	93
		40 41	46	53	60	67	73	80	87	93
		41	47	54 55	60	67	74 74	80	87	93 94
		42	48 49	55 55	61 62	68 68	74	81 81	87 87	94
		43	50	56	62	69	75	81	87	94
		44	50	57	63	69	75	81	88	94
		45	51	57	63	70	76	82	88	94
		46	52	58	64	70	76	82	88	94
		47	53	59	65	71	76	82	88	94
42 .		48	54	59	65	71	77	83	88	94
43 .		49	54	60	66	72	77	83	89	94
44 .		50	55	61	66	72	78	83	89	94
45 .		51	56	62	67	73	78	84	89	95
46 .		51	57	62	68	73	78	84	89	95
47 .		52	58	63	68	74	79	84	89	95
48 .		53	58	64	69	74	79	84	90	95
		54	59	64	69	75	80	85	90	95
		55	60	65	70	75	80	85	90	95
		56	61	66	71	76	80	85	90	95
		57	62	66	71	76	81	86	90	95
		58	62	67	72	77	81	86	91	95
		59 60	63 64	68 69	72 73	77 78	82 82	86 87	91 91	95 96
		60	65	69	74	78	82	87	91	96
		61	66	70	74	79	83	87	91	96
		62	66	71	75	79	83	87	92	96
		63	67	71	75	80	84	88	92	96
		64	68	72	76	80	84	88	92	96
		65	69	73	77	81	84	88	92	96
		66	70	73	77	81	85	89	92	96
		67	70	74	78	82	85	89	93	96
64 .		68	71	75	78	82	86	89	93	96
65 .		69	72	76	79	83	86	90	93	97
66 .		69	73	76	80	83	86	90	93	97
67 .		70	74	77	80	84	87	90	93	97
68 .		71	74	78	81	84	87	90	94	97
69 .		72	75	78	81	85	88	91	94	97

TABLE I—COMBINED RATINGS TABLE—Continued
[10 combined with 10 is 19]

	10	20	30	40	50	60	70	80	90
70	73	76	79	82	85	88	91	94	97
71	74	77	80	83	86	88	91	94	97
72	75	78	80	83	86	89	92	94	97
73	76	78	81	84	87	89	92	95	97
74	77	79	82	84	87	90	92	95	97
75	78	80	83	85	88	90	93	95	98
76	78	81	83	86	88	90	93	95	98
77	79	82	84	86	89	91	93	95	98
78	80	82	85	87	89	91	93	96	98
79	81	83	85	87	90	92	94	96	98
80	82	84	86	88	90	92	94	96	98
81	83	85	87	89	91	92	94	96	98
82	84	86	87	89	91	93	95	96	98
83	85	86	88	90	92	93	95	97	98
84	86	87	89	90	92	94	95	97	98
85	87	88	90	91	93	94	96	97	99
86	87	89	90	92	93	94	96	97	99
87	88	90	91	92	94	95	96	97	99
88	89	90	92	93	94	95	96	98	99
89	90	91	92	93	95	96	87	38	99
90	91	92	93	94	95	96	97	98	99
91	92	93	94	95	96	96	97	98	99
92	93	94	94	95	96	97	98	98	99
93	94	94	95	96	97	97	98	99	99
94	95	95	96	96	97	98	98	99	99

(Authority: 38 U.S.C. 1155)

[41 FR 11293, Mar. 18, 1976, as amended at 54 FR 27161, June 28, 1989; 54 FR 36029, Aug. 31, 1989]

### §4.26 Bilateral factor.

When a partial disability results from disease or injury of both arms, or of both legs, or of paired skeletal muscles, the ratings for the disabilities of the right and left sides will be combined as usual, and 10 percent of this value will be added (i.e., not combined) before proceeding with further combinations, or converting to degree of disability. The bilateral factor will be applied to such bilateral disabilities before other combinations are carried out and the rating for such disabilities including the bilateral factor in this section will be treated as 1 disability for the purpose of arranging in order of severity and for all further combinations. For example, with disabilities evaluated at 60 percent, 20 percent, 10 percent and 10 percent (the two 10's representing bilateral disabilities), the order of severity would be 60, 21 and 20. The 60 and 21 combine to 68 percent and the 68 and 20 to 74 percent, converted to 70 percent as the final degree of disability.

(a) The use of the terms "arms" and "legs" is not intended to distinguish

between the arm, forearm and hand, or the thigh, leg, and foot, but relates to the upper extremities and lower extremities as a whole. Thus with a compensable disability of the right thigh, for example, amputation, and one of the left foot, for example, pes planus, the bilateral factor applies, and similarly whenever there are compensable disabilities affecting use of paired extremities regardless of location or specified type of impairment.

- (b) The correct procedure when applying the bilateral factor to disabilities affecting both upper extremities and both lower extremities is to combine the ratings of the disabilities affecting the 4 extremities in the order of their individual severity and apply the bilateral factor by adding, not combining, 10 percent of the combined value thus attained.
- (c) The bilateral factor is not applicable unless there is partial disability of compensable degree in each of 2 paired extremities, or paired skeletal muscles.

#### §4.27 Use of diagnostic code numbers.

The diagnostic code numbers appearing opposite the listed ratable disabilities are arbitrary numbers for the purpose of showing the basis of the evaluation assigned and for statistical analysis in the Department of Veterans Affairs, and as will be observed, extend from 5000 to a possible 9999. Great care will be exercised in the selection of the applicable code number and in its citation on the rating sheet. No other numbers than these listed or hereafter furnished are to be employed for rating purposes, with an exception as described in this section, as to unlisted conditions. When an unlisted disease, injury, or residual condition is encountered, requiring rating by analogy, the diagnostic code number will be "builtup" as follows: The first 2 digits will be selected from that part of the schedule most closely identifying the part, or system, of the body involved; the last 2 digits will be "99" for all unlisted conditions. This procedure will facilitate a close check of new and unlisted conditions, rated by analogy. In the selection of code numbers, injuries will generally be represented by the number assigned to the residual condition on the basis of which the rating is determined. With diseases, preference is to be given to the number assigned to the disease itself; if the rating is determined on the basis of residual conditions, the number appropriate to the residual condition will be added, preceded by a hyphen. Thus, rheumatoid (atrophic) arthritis rated as ankylosis of the lumbar spine should be coded "5002-5240." In this way, the exact source of each rating can be easily identified. In the citation of disabilities on rating sheets, the diagnostic terminology will be that of the medical examiner, with no attempt to translate the terms into schedule nomenclature. Residuals of diseases or therapeutic procedures will not be cited without reference to the basic disease.

[41 FR 11293, Mar. 18, 1976, as amended at 70 FR 75399, Dec. 20, 2005]

# § 4.28 Prestabilization rating from date of discharge from service.

The following ratings may be assigned, in lieu of ratings prescribed

elsewhere, under the conditions stated for disability from any disease or injury. The prestabilization rating is not to be assigned in any case in which a total rating is immediately assignable under the regular provisions of the schedule or on the basis of individual unemployability. The prestabilization 50-percent rating is not to be used in any case in which a rating of 50 percent or more is immediately assignable under the regular provisions.

	Rating
Unstabilized condition with severe disability— Substantially gainful employment is not feasible or advisable Unhealed or incompletely healed wounds or injuries—	100
Material impairment of employability likely	50

NOTE (1): Department of Veterans Affairs examination is not required prior to assignment of prestabilization ratings; however, the fact that examination was accomplished will not preclude assignment of these benefits. Prestabilization ratings are for assignment in the immediate postdischarge period. They will continue for a 12-month period following discharge from service. However, prestabilization ratings may be changed to a regular schedular total rating or one authorizing a greater benefit at any time. In each prestabilization rating an examination will be requested to be accomplished not earlier than 6 months nor more than 12 months following discharge. In those prestabilization ratings in which following examination reduction in evaluation is found to be warranted, the higher evaluation will be continued to the end of the 12th month following discharge or to the end of the period provided under §3.105(e) of this chapter, whichever is later. Special monthly compensation should be assigned concurrently in these cases whenever records are adequate to establish entitlement.

NOTE (2): Diagnosis of disease, injury, or residuals will be cited, with diagnostic code number assigned from this rating schedule for conditions listed therein.

[35 FR 11906, July 24, 1970]

# § 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established that a service-connected disability has required hospital treatment in a Department of

Veterans Affairs or an approved hospital for a period in excess of 21 days or hospital observation at Department of Veterans Affairs expense for a service-connected disability for a period in excess of 21 days.

- (a) Subject to the provisions of paragraphs (d), (e), and (f) of this section this increased rating will be effective the first day of continuous hospitalization and will be terminated effective the last day of the month of hospital discharge (regular discharge or release to non-bed care) or effective the last day of the month of termination of treatment or observation for the service-connected disability. A temporary release which is approved by an attending Department of Veterans Affairs physician as part of the treatment plan will not be considered an absence.
- (1) An authorized absence in excess of 4 days which begins during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the first day of such authorized absence. An authorized absence of 4 days or less which results in a total of more than 8 days of authorized absence during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the ninth day of authorized absence.
- (2) Following a period of hospitalization in excess of 21 days, an authorized absence in excess of 14 days or a third consecutive authorized absence of 14 days will be regarded as the equivalent of hospital discharge and will interrupt hospitalization effective on the last day of the month in which either the authorized absence in excess of 14 days or the third 14 day period begins, except where there is a finding that convalescence is required as provided by paragraph (e) or (f) of this section. The termination of these total ratings will not be subject to §3.105(e) of this chapter.
- (b) Notwithstanding that hospital admission was for disability not connected with service, if during such hospitalization, hospital treatment for a service-connected disability is instituted and continued for a period in excess of 21 days, the increase to a total rating will be granted from the first day of such treatment. If service connection for the disability under treat-

ment is granted after hospital admission, the rating will be from the first day of hospitalization if otherwise in order.

- (c) The assignment of a total disability rating on the basis of hospital treatment or observation will not preclude the assignment of a total disability rating otherwise in order under other provisions of the rating schedule, and consideration will be given to the propriety of such a rating in all instances and to the propriety of its continuance after discharge. Particular attention, with a view to proper rating under the rating schedule, is to be given to the claims of veterans discharged from hospital, regardless of length of hospitalization, with indications on the final summary of expected confinement to bed or house, or to inability to work with requirement of frequent care of physician or nurse at home.
- (d) On these total ratings Department of Veterans Affairs regulations governing effective dates for increased benefits will control.
- (e) The total hospital rating if convalescence is required may be continued for periods of 1, 2, or 3 months in addition to the period provided in paragraph (a) of this section.

(f) Extension of periods of 1, 2 or 3 months beyond the initial 3 months may be made upon approval of the Veterans Service Center Manager.

(g) Meritorious claims of veterans who are discharged from the hospital with less than the required number of days but need post-hospital care and a prolonged period of convalescence will be referred to the Director, Compensation and Pension Service, under §3.321(b)(1) of this chapter.

[29 FR 6718, May 22, 1964, as amended at 41 FR 11294, Mar. 18, 1976; 41 FR 34256, Aug. 13, 1976; 54 FR 4281, Jan. 30, 1989; 54 FR 34981, Aug. 23, 1989; 71 FR 28586, May 17, 2006]

#### § 4.30 Convalescent ratings.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established by report at hospital discharge (regular discharge or release to non-bed care) or outpatient release that entitlement is warranted under paragraph (a) (1), (2) or (3) of this

section effective the date of hospital admission or outpatient treatment and continuing for a period of 1, 2, or 3 months from the first day of the month following such hospital discharge or outpatient release. The termination of these total ratings will not be subject to §3.105(e) of this chapter. Such total rating will be followed by appropriate schedular evaluations. When the evidence is inadequate to assign a schedular evaluation, a physical examination will be scheduled and considered prior to the termination of a total rating under this section.

- (a) Total ratings will be assigned under this section if treatment of a service-connected disability resulted in:
- (1) Surgery necessitating at least one month of convalescence (Effective as to outpatient surgery March 1, 1989.)
- (2) Surgery with severe postoperative residuals such as incompletely healed surgical wounds, stumps of recent amputations, therapeutic immobilization of one major joint or more, application of a body cast, or the necessity for house confinement, or the necessity for continued use of a wheelchair or crutches (regular weight-bearing prohibited). (Effective as to outpatient surgery March 1, 1989.)
- (3) Immobilization by cast, without surgery, of one major joint or more. (Effective as to outpatient treatment March 10, 1976.)

A reduction in the total rating will not be subject to §3.105(e) of this chapter. The total rating will be followed by an open rating reflecting the appropriate schedular evaluation; where the evidence is inadequate to assign the schedular evaluation, a physcial examination will be scheduled prior to the end of the total rating period.

- (b) A total rating under this section will require full justification on the rating sheet and may be extended as follows:
- (1) Extensions of 1, 2 or 3 months beyond the initial 3 months may be made under paragraph (a) (1), (2) or (3) of this section.
- (2) Extensions of 1 or more months up to 6 months beyond the initial 6 months period may be made under paragraph (a) (2) or (3) of this section

upon approval of the Veterans Service Center Manager.

[41 FR 34256, Aug. 13, 1976, as amended at 54 FR 4281, Jan. 30, 1989; 71 FR 28586, May 17, 2006]

### § 4.31 Zero percent evaluations.

In every instance where the schedule does not provide a zero percent evaluation for a diagnostic code, a zero percent evaluation shall be assigned when the requirements for a compensable evaluation are not met.

[58 FR 52018, Oct. 6, 1993]

## Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

#### § 4.40 Functional loss.

Disability of the musculoskeletal system is primarily the inability, due to damage or infection in parts of the system, to perform the normal working movements of the body with normal excursion, strength, speed, coordination and endurance. It is essential that the examination on which ratings are based adequately portray the anatomical damage, and the functional loss, with respect to all these elements. The functional loss may be due to absence of part, or all, of the necessary bones. joints and muscles, or associated structures, or to deformity, adhesions, defective innervation, or other pathology, or it may be due to pain, supported by adequate pathology and evidenced by the visible behavior of the claimant undertaking the motion. Weakness is as important as limitation of motion, and a part which becomes painful on use must be regarded as seriously disabled. A little used part of the musculoskeletal system may be expected to show evidence of disuse, either through atrophy, the condition of the skin, absence of normal callosity or the like.

### § 4.41 History of injury.

In considering the residuals of injury, it is essential to trace the medical-industrial history of the disabled person from the original injury, considering

the nature of the injury and the attendant circumstances, and the reguirements for, and the effect of, treatment over past periods, and the course of the recovery to date. The duration of the initial, and any subsequent, period of total incapacity, especially periods reflecting delayed union, inflammation, swelling, drainage, or operative intervention, should be given close attention. This consideration, or the absence of clear cut evidence of injury, may result in classifying the disability as not of traumatic origin, either reflecting congenital or developmental etiology, or the effects of healed dis-

# § 4.42 Complete medical examination of injury cases.

The importance of complete medical examination of injury cases at the time of first medical examination by the Department of Veterans Affairs cannot be overemphasized. When possible, this should include complete neurological and psychiatric examination, and other special examinations indicated by the physical condition, in addition to the required general and orthopedic or surgical examinations. When complete examinations are not conducted covering all systems of the body affected by disease or injury, it is impossible to visualize the nature and extent of the service connected disability. Incomplete examination is a common cause of incorrect diagnosis, especially in the neurological and psychiatric fields, and frequently leaves the Department of Veterans Affairs in doubt as to the presence or absence of disabling conditions at the time of the examination.

### § 4.43 Osteomyelitis.

Chronic, or recurring, suppurative osteomyelitis, once clinically identified, including chronic inflammation of bone marrow, cortex, or periosteum, should be considered as a continuously disabling process, whether or not an actively discharging sinus or other obvious evidence of infection is manifest from time to time, and unless the focus is entirely removed by amputation will entitle to a permanent rating to be combined with other ratings for residual conditions, however, not exceeding

amputation ratings at the site of election.

#### § 4.44 The bones.

The osseous abnormalities incident to trauma or disease, such as malunion with deformity throwing abnormal stress upon, and causing malalignment of joint surfaces, should be depicted from study and observation of all available data, beginning with inception of injury or disease, its nature, degree of prostration, treatment and duration of convalescence, and progress of recovery with development of permanent residuals. With shortening of a long bone, some degree of angulation is to be expected; the extent and direction should be brought out by X-ray and observation. The direction of angulation and extent of deformity should be carefully related to strain on the neighboring joints, especially those connected with weight-bearing.

## § 4.45 The joints.

As regards the joints the factors of disability reside in reductions of their normal excursion of movements in different planes. Inquiry will be directed to these considerations:

- (a) Less movement than normal (due to ankylosis, limitation or blocking, adhesions, tendon-tie-up, contracted scars, etc.).
- (b) More movement than normal (from flail joint, resections, nonunion of fracture, relaxation of ligaments, etc.).
- (c) Weakened movement (due to muscle injury, disease or injury of peripheral nerves, divided or lengthened tendons, etc.).
  - (d) Excess fatigability.
- (e) Incoordination, impaired ability to execute skilled movements smoothly.
- (f) Pain on movement, swelling, deformity or atrophy of disuse. Instability of station, disturbance of locomotion, interference with sitting, standing and weight-bearing are related considerations. For the purpose of rating disability from arthritis, the shoulder, elbow, wrist, hip, knee, and ankle are considered major joints; multiple involvements of the interphalangeal, metacarpal and carpal joints of

the upper extremities, the interphalangeal, metatarsal and tarsal joints of the lower extremities, the cervical vertebrae, the dorsal vertebrae, and the lumbar vertebrae, are considered groups of minor joints, ratable on a parity with major joints. The lumbosacral articulation and both sacroiliac joints are considered to be a group of minor joints, ratable on disturbance of lumbar spine functions.

#### § 4.46 Accurate measurement.

Accurate measurement of the length of stumps, excursion of joints, dimensions and location of scars with respect to landmarks, should be insisted on. The use of a goniometer in the measurement of limitation of motion is indispensable in examinations conducted within the Department of Veterans Affairs. Muscle atrophy must also be accurately measured and reported.

[41 FR 11294, Mar. 18, 1976]

## §§ 4.47-4.54 [Reserved]

# § 4.55 Principles of combined ratings for muscle injuries.

- (a) A muscle injury rating will not be combined with a peripheral nerve paralysis rating of the same body part, unless the injuries affect entirely different functions.
- (b) For rating purposes, the skeletal muscles of the body are divided into 23 muscle groups in 5 anatomical regions: 6 muscle groups for the shoulder girdle and arm (diagnostic codes 5301 through 5306); 3 muscle groups for the forearm and hand (diagnostic codes 5307 through 5309); 3 muscle groups for the foot and leg (diagnostic codes 5310 through 5312); 6 muscle groups for the pelvic girdle and thigh (diagnostic codes 5313 through 5318); and 5 muscle groups for the torso and neck (diagnostic codes 5319 through 5323).
- (c) There will be no rating assigned for muscle groups which act upon an ankylosed joint, with the following exceptions:
- (1) In the case of an ankylosed knee, if muscle group XIII is disabled, it will be rated, but at the next lower level than that which would otherwise be assigned.
- (2) In the case of an ankylosed shoulder, if muscle groups I and II are se-

verely disabled, the evaluation of the shoulder joint under diagnostic code 5200 will be elevated to the level for unfavorable ankylosis, if not already assigned, but the muscle groups themselves will not be rated.

- (d) The combined evaluation of muscle groups acting upon a single unankylosed joint must be lower than the evaluation for unfavorable ankylosis of that joint, except in the case of muscle groups I and II acting upon the shoulder.
- (e) For compensable muscle group injuries which are in the same anatomical region but do not act on the same joint, the evaluation for the most severely injured muscle group will be increased by one level and used as the combined evaluation for the affected muscle groups.
- (f) For muscle group injuries in different anatomical regions which do not act upon ankylosed joints, each muscle group injury shall be separately rated and the ratings combined under the provisions of §4.25.

(Authority: 38 U.S.C. 1155) [62 FR 30237, June 3, 1997]

# §4.56 Evaluation of muscle disabilities.

- (a) An open comminuted fracture with muscle or tendon damage will be rated as a severe injury of the muscle group involved unless, for locations such as in the wrist or over the tibia, evidence establishes that the muscle damage is minimal.
- (b) A through-and-through injury with muscle damage shall be evaluated as no less than a moderate injury for each group of muscles damaged.
- (c) For VA rating purposes, the cardinal signs and symptoms of muscle disability are loss of power, weakness, lowered threshold of fatigue, fatigue-pain, impairment of coordination and uncertainty of movement.
- (d) Under diagnostic codes 5301 through 5323, disabilities resulting from muscle injuries shall be classified as slight, moderate, moderately severe or severe as follows:
- (1) Slight disability of muscles—(i) Type of injury. Simple wound of muscle without debridement or infection.

(ii) History and complaint. Service department record of superficial wound with brief treatment and return to duty. Healing with good functional results. No cardinal signs or symptoms of muscle disability as defined in paragraph (c) of this section.

(iii) Objective findings. Minimal scar. No evidence of fascial defect, atrophy, or impaired tonus. No impairment of function or metallic fragments re-

tained in muscle tissue.

- (2) Moderate disability of muscles—(i) Type of injury. Through and through or deep penetrating wound of short track from a single bullet, small shell or shrapnel fragment, without explosive effect of high velocity missile, residuals of debridement, or prolonged infection.
- (ii) History and complaint. Service department record or other evidence of in-service treatment for the wound. Record of consistent complaint of one or more of the cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, particularly lowered threshold of fatigue after average use, affecting the particular functions controlled by the injured muscles.
- (iii) Objective findings. Entrance and (if present) exit scars, small or linear, indicating short track of missile through muscle tissue. Some loss of deep fascia or muscle substance or impairment of muscle tonus and loss of power or lowered threshold of fatigue when compared to the sound side.
- (3) Moderately severe disability of muscles—(i) Type of injury. Through and through or deep penetrating wound by small high velocity missile or large low-velocity missile, with debridement, prolonged infection, or sloughing of soft parts, and intermuscular scarring.
- (ii) History and complaint. Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section and, if present, evidence of inability to keep up with work requirements.
- (iii) *Objective findings.* Entrance and (if present) exit scars indicating track of missile through one or more muscle

groups. Indications on palpation of loss of deep fascia, muscle substance, or normal firm resistance of muscles compared with sound side. Tests of strength and endurance compared with sound side demonstrate positive evidence of impairment.

- (4) Severe disability of muscles—(i) Type of injury. Through and through or deep penetrating wound due to high-velocity missile, or large or multiple low velocity missiles, or with shattering bone fracture or open comminuted fracture with extensive debridement, prolonged infection, or sloughing of soft parts, intermuscular binding and scarring.
- (ii) History and complaint. Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, worse than those shown for moderately severe muscle injuries, and, if present, evidence of inability to keep up with work requirements.
- (iii) Objective findings. Ragged, depressed and adherent scars indicating wide damage to muscle groups in missile track. Palpation shows loss of deep fascia or muscle substance, or soft flabby muscles in wound area. Muscles swell and harden abnormally in contraction. Tests of strength, endurance, or coordinated movements compared with the corresponding muscles of the uninjured side indicate severe impairment of function. If present, the following are also signs of severe muscle disability:
- (A) X-ray evidence of minute multiple scattered foreign bodies indicating intermuscular trauma and explosive effect of the missile.
- (B) Adhesion of scar to one of the long bones, scapula, pelvic bones, sacrum or vertebrae, with epithelial sealing over the bone rather than true skin covering in an area where bone is normally protected by muscle.
- (C) Diminished muscle excitability to pulsed electrical current in electrodiagnostic tests.
- (D) Visible or measurable atrophy.(E) Adaptive contraction of an opposing group of muscles.
- (F) Atrophy of muscle groups not in the track of the missile, particularly of

the trapezius and serratus in wounds of the shoulder girdle.

(G) Induration or atrophy of an entire muscle following simple piercing by a projectile.

(Authority: 38 U.S.C. 1155 [62 FR 30238, June 3, 1997]

#### § 4.57 Static foot deformities.

It is essential to make an initial distinction between bilateral flatfoot as a congenital or as an acquired condition. The congenital condition, with depression of the arch, but no evidence of abnormal callosities, areas of pressure, strain or demonstrable tenderness, is a congenital abnormality which is not compensable or pensionable. In the acquired condition, it is to be remembered that depression of the longitudinal arch, or the degree of depression, is not the essential feature. The attention should be given to anatomical changes, as compared to normal, in the relationship of the foot and leg, particularly to the inward rotation of the superior portion of the os calcis, medial deviation of the insertion of the Achilles tendon, the medial tilting of the upper border of the astragalus. This is an unfavorable mechanical relationship of the parts. A plumb line dropped from the middle of the patella falls inside of the normal point. The forepart of the foot is abducted, and the foot everted. The plantar surface of the foot is painful and shows demonstrable tenderness, and manipulation of the foot produces spasm of the Achilles tendon, peroneal spasm due to adhesion about the peroneal sheaths, and other evidence of pain and limited motion. The symptoms should be apparent without regard to exercise. In severe cases there is gaping of bones on the inner border of the foot, and rigid valgus position with loss of the power of inversion and adduction. Exercise with undeveloped or unbalanced musculature, producing chronic irritation, can be an aggravating factor. In the absence of trauma or other definite evidence of aggravation, service connection is not in order for pes cavus which is a typically congenital or juvenile disease.

#### § 4.58 Arthritis due to strain.

With service incurred lower extremity amputation or shortening, a disabling arthritis, developing in the same extremity, or in both lower extremities, with indications of earlier, or more severe, arthritis in the injured extremity, including also arthritis of the lumbosacral joints and lumbar spine, if associated with the leg amputation or shortening, will be considered as service incurred, provided, however, that arthritis affecting joints not directly subject to strain as a result of the service incurred amputation will not be granted service connection. This will generally require separate evaluation of the arthritis in the joints directly subject to strain. Amputation, or injury to an upper extremity, is not considered as a causative factor with subsequently developing arthritis, except in joints subject to direct strain or actually injured.

## §4.59 Painful motion.

With any form of arthritis, painful motion is an important factor of disability, the facial expression, wincing, etc., on pressure or manipulation, should be carefully noted and definitely related to affected joints. Muscle spasm will greatly assist the identification. Sciatic neuritis is not uncommonly caused by arthritis of the spine. The intent of the schedule is to recognize painful motion with joint or periarticular pathology as productive of disability. It is the intention to recognize actually painful, unstable, or malaligned joints, due to healed injury, as entitled to at least the minimum compensable rating for the joint. Crepitation either in the soft tissues such as the tendons or ligaments, or crepitation within the joint structures should be noted carefully as points of contact which are diseased. Flexion elicits such manifestations. The joints involved should be tested for pain on both active and passive motion, in weight-bearing and nonweight-bearing and, if possible, with the range of the opposite undamaged joint.

### §4.60 [Reserved]

#### § 4.61 Examination.

With any form of arthritis (except traumatic arthritis) it is essential that the examination for rating purposes cover all major joints, with especial reference to Heberden's or Haygarth's nodes.

#### § 4.62 Circulatory disturbances.

The circulatory disturbances, especially of the lower extremity following injury in the popliteal space, must not be overlooked, and require rating generally as phlebitis.

#### § 4.63 Loss of use of hand or foot.

Loss of use of a hand or a foot, for the purpose of special monthly compensation, will be held to exist when no effective function remains other than that which would be equally well served by an amputation stump at the site of election below elbow or knee with use of a suitable prosthetic appliance. The determination will be made on the basis of the actual remaining function of the hand or foot, whether the acts of grasping, manipulation, etc., in the case of the hand, or of balance and propulsion, etc., in the case of the foot, could be accomplished equally well by an amputation stump with prosthesis.

(a) Extremely unfavorable complete ankylosis of the knee, or complete ankylosis of 2 major joints of an extremity, or shortening of the lower extremity of 3½ inches (8.9 cms.) or more, will be taken as loss of use of the hand or foot involved.

(b) Complete paralysis of the external popliteal nerve (common peroneal) and consequent, footdrop, accompanied by characteristic organic changes including trophic and circulatory disturbances and other concomitants confirmatory of complete paralysis of this nerve, will be taken as loss of use of the foot.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

## § 4.64 Loss of use of both buttocks.

Loss of use of both buttocks shall be deemed to exist when there is severe damage to muscle Group XVII, bilat-

eral (diagnostic code number 5317) and additional disability rendering it impossible for the disabled person, without assistance, to rise from a seated position and from a stooped position (fingers to toes position) and to maintain postural stability (the pelvis upon head of femur). The assistance may be rendered by the person's own hands or arms, and, in the matter of postural stability, by a special appliance.

#### § 4.65 [Reserved]

### § 4.66 Sacroiliac joint.

The common cause of disability in this region is arthritis, to be identified in the usual manner. The lumbosacral and sacroiliac joints should be considered as one anatomical segment for rating purposes. X-ray changes from arthritis in this location are decrease or obliteration of the joint space, with the appearance of increased bone density of the sacrum and ilium and sharpening of the margins of the joint. Disability is manifest from erector spinae spasm (not accounted for by other pathology), tenderness on deep palpation and percussion over these joints, loss of normal quickness of motion and resiliency, and postural defects often accompanied by limitation of flexion and extension of the hip. Traumatism is a rare cause of disability in this connection, except when superimposed upon congenital defect or upon an existent arthritis; to permit assumption of pure traumatic origin, objective evidence of damage to the joint, and history of trauma sufficiently severe to injure this extremely strong and practically immovable joint is required. There should be careful consideration of lumbosacral sprain, and the various symptoms of pain and paralysis attributable to disease affecting the lumbar vertebrae and the intervertebral disc.

## §4.67 Pelvic bones.

The variability of residuals following these fractures necessitates rating on specific residuals, faulty posture, limitation of motion, muscle injury, painful motion of the lumbar spine, manifest by muscle spasm, mild to moderate sciatic neuritis, peripheral nerve injury, or limitation of hip motion.

### §4.68 Amputation rule.

The combined rating for disabilities of an extremity shall not exceed the rating for the amputation at the elective level, were amputation to be performed. For example, the combined evaluations for disabilities below the knee shall not exceed the 40 percent evaluation, diagnostic code 5165. This 40 percent rating may be further combined with evaluation for disabilities above the knee but not to exceed the above the knee amputation elective level. Painful neuroma of a stump after amputation shall be assigned the evaluation for the elective site of reamputation.

#### § 4.69 Dominant hand.

Handedness for the purpose of a dominant rating will be determined by the evidence of record, or by testing on VA examination. Only one hand shall be considered dominant. The injured hand, or the most severely injured hand, of an ambidextrous individual will be considered the dominant hand for rating purposes.

(Authority: 38 U.S.C. 1155) [62 FR 30239, June 3, 1997]

#### § 4.70 Inadequate examinations.

If the report of examination is inadequate as a basis for the required consideration of service connection and evaluation, the rating agency may request a supplementary report from the examiner giving further details as to the limitations of the disabled person's ordinary activity imposed by the disease, injury, or residual condition, the prognosis for return to, or continuance of, useful work. When the best interests of the service will be advanced by personal conference with the examiner, such conference may be arranged through channels.

# § 4.71 Measurement of ankylosis and joint motion.

Plates I and II provide a standardized description of ankylosis and joint motion measurement. The anatomical position is considered as  $0^{\circ}$ , with two major exceptions: (a) Shoulder rotation—arm abducted to 90°, elbow flexed to  $90^{\circ}$  with the position of the forearm reflecting the midpoint 0° between internal and external rotation of the shoulder; and (b) supination and pronation—the arm next to the body. elbow flexed to 90°, and the forearm in midposition 0° between supination and pronation. Motion of the thumb and fingers should be described by appropriate reference to the joints (See Plate III) whose movement is limited, with a statement as to how near, in centimeters, the tip of the thumb can approximate the fingers, or how near the tips of the fingers can approximate the proximal transverse crease of palm.




PLATE I




PLATE II

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978; 67 FR 48785, July 26, 2002]

# $\$\,4.71a$ Schedule of ratings—musculo-skeletal system.

ACUTE, SUBACUTE, OR CHRONIC DISEASES

	Rat- ing
5000 Osteomyelitis, acute, subacute, or chronic:	
Of the pelvis, vertebrae, or extending into major	
joints, or with multiple localization or with long history of intractability and debility, anemia, amyloid liver changes, or other continuous	
constitutional symptoms	100
Frequent episodes, with constitutional symptoms	60
With definite involucrum or sequestrum, with or without discharging sinus	30
With discharging sinus or other evidence of active infection within the past 5 years	20
Inactive, following repeated episodes, without evidence of active infection in past 5 years	10

ACUTE, SUBACUTE, OR CHRONIC DISEASES—Continued

Rat-

NOTE (1): A rating of 10 percent, as an exception to the amputation rule, is to be assigned in any case of active osteomyelitis where the amputation rating for the affected part is no percent. This 10 percent rating and the other partial ratings of 30 percent or less are to be combined with ratings for ankylosis, limited motion, nonunion or malunion, shortening, etc., subject, of course, to the amputation rule. The 60 percent rating, as it is based on constitutional symptoms, is not subject to the amputation rule. A rating for osteomyelitis will not be applied following cure by removal or radical resection of the affected bone.

## **Department of Veterans Affairs**

§4.71a

10

### ACUTE, SUBACUTE, OR CHRONIC DISEASES— Continued

Continued		
	Rat- ing	
NOTE (2): The 20 percent rating on the basis of		W
activity within the past 5 years is not assign-		
able following the initial infection of active osteomyelitis with no subsequent reactivation.		
The prerequisite for this historical rating is an		W
established recurrent osteomyelitis. To qualify		
for the 10 percent rating, 2 or more episodes		
following the initial infection are required. This 20 percent rating or the 10 percent rating,		NOTE on X
when applicable, will be assigned once only to		with
cover disability at all sites of previously active		Note
infection with a future ending date in the case		on X
of the 20 percent rating.  5001 Bones and joints, tuberculosis of, active or in-		ratin
active:		5013 5004 Arth
Active	100	5005 Arth
Inactive: See §§ 4.88b and 4.89		5006 Arth
5002 Arthritis rheumatoid (atrophic) As an active		5007 Arth
process: With constitutional manifestations associated		5008 Arth
with active joint involvement, totally incapaci-		5009 Arth With th
tating	100	thro
Less than criteria for 100% but with weight loss		arthi
and anemia productive of severe impairment		5010 Arth
of health or severely incapacitating exacer- bations occurring 4 or more times a year or a		ray findir
lesser number over prolonged periods	60	5011 Bon
Symptom combinations productive of definite im-		cord inve severity
pairment of health objectively supported by ex-		5012 Bon
amination findings or incapacitating exacer-	40	Note:
bations occurring 3 or more times a year  One or two exacerbations a year in a well-estab-	40	for '
lished diagnosis	20	X-ra thera
For chronic residuals:		has
For residuals such as limitation of motion or an-		the i
kylosis, favorable or unfavorable, rate under		5013 Ost
the appropriate diagnostic codes for the spe- cific joints involved. Where, however, the limi-		5014 Ost
tation of motion of the specific joint or joints in-		5015 Bor 5016 Ost
volved is noncompensable under the codes a		5016 Ost 5017 Gou
rating of 10 percent is for application for each		5017 God
such major joint or group of minor joints af- fected by limitation of motion, to be combined,		5019 Burs
not added under diagnostic code 5002. Limita-		5020 Syn
tion of motion must be objectively confirmed		5021 Myd
by findings such as swelling, muscle spasm,		5022 Peri 5023 Myd
or satisfactory evidence of painful motion.		5023 Myc
NOTE: The ratings for the active process will not be combined with the residual ratings for limi-		The o
tation of motion or ankylosis. Assign the high-		thro
er evaluation.		tion
5003 Arthritis, degenerative (hypertrophic or osteo-		tive, agno
arthritis):  Degenerative arthritis established by X-ray find-		5025 Fibr
ings will be rated on the basis of limitation of		syndrom
motion under the appropriate diagnostic codes		With v
for the specific joint or joints involved (DC		der
5200 etc.). When however, the limitation of		slee head
motion of the specific joint or joints involved is noncompensable under the appropriate diag-		sion
nostic codes, a rating of 10 pct is for applica-		Th
tion for each such major joint or group of		
minor joints affected by limitation of motion, to		Th
be combined, not added under diagnostic		
code 5003. Limitation of motion must be objectively confirmed by findings such as swell-		
ing, muscle spasm, or satisfactory evidence of		
painful motion. In the absence of limitation of motion, rate as below:		Th

# ACUTE, SUBACUTE, OR CHRONIC DISEASES—Continued

	Rat- ing
With X-ray evidence of involvement of 2 or	
more major joints or 2 or more minor joint	
groups, with occasional incapacitating ex-	
acerbations	20
With X-ray evidence of involvement of 2 or more major joints or 2 or more minor joint	
groups	10
Note (1): The 20 pct and 10 pct ratings based	
on X-ray findings, above, will not be combined	
with ratings based on limitation of motion.	
NOTE (2): The 20 pct and 10 pct ratings based	
on X-ray findings, above, will not be utilized in	
rating conditions listed under diagnostic codes 5013 to 5024, inclusive.	
5004 Arthritis, gonorrheal.	
5005 Arthritis, pneumococcic.	
5006 Arthritis, typhoid.	
5007 Arthritis, syphilitic.	
5008 Arthritis, streptococcic.	
5009 Arthritis, other types (specify).	
With the types of arthritis, diagnostic codes 5004	
through 5009, rate the disability as rheumatoid arthritis.	
5010 Arthritis, due to trauma, substantiated by X-	
ray findings: Rate as arthritis, degenerative.	
5011 Bones, caisson disease of: Rate as arthritis,	
cord involvement, or deafness, depending on the	
severity of disabling manifestations.	
5012 Bones, new growths of, malignant	100
NOTE: The 100 percent rating will be continued for 1 year following the cessation of surgical,	
X-ray, antineoplastic chemotherapy or other	
therapeutic procedure. At this point, if there	
has been no local recurrence or metastases,	
the rating will be made on residuals.	
5013 Osteoporosis, with joint manifestations. 5014 Osteomalacia.	
5015 Bones, new growths of, benign.	
5016 Osteitis deformans.	
5017 Gout.	
5018 Hydrarthrosis, intermittent.	
5019 Bursitis.	
5020 Synovitis.	
5021 Myositis.	
5022 Periostitis. 5023 Myositis ossificans.	
5024 Tenosynovitis.	
The diseases under diagnostic codes 5013	
through 5024 will be rated on limitation of mo-	
tion of affected parts, as arthritis, degenera-	
tive, except gout which will be rated under di-	
agnostic code 5002. 5025 Fibromyalgia (fibrositis, primary fibromyalgia	
syndrome)	
With widespread musculoskeletal pain and ten-	
der points, with or without associated fatigue,	
sleep disturbance, stiffness, paresthesias,	
headache, irritable bowel symptoms, depres-	
sion, anxiety, or Raynaud's-like symptoms:	
That are constant, or nearly so, and refractory to therapy	40
That are episodic, with exacerbations often	70
precipitated by environmental or emo-	
precipitated by environmental or emo- tional stress or by overexertion, but that	
are present more than one-third of the	
That require continuous medication for con	20
That require continuous medication for con-	10

# 38 CFR Ch. I (7-1-07 Edition)

# §4.71a

# ACUTE, SUBACUTE, OR CHRONIC DISEASES— Continued

	Rat- ing
NOTE: Widespread pain means pain in both the left and right sides of the body, that is both above and below the waist, and that affects both the axial skeleton (i.e., cervical spine, anterior chest, thoracic spine, or low back) and the extremities.	

## PROSTHETIC IMPLANTS

	Rating	
	Major	Minor
5051 Shoulder replacement (prosthesis). Prosthetic replacement of the shoulder joint:		
For 1 year following implantation of prosthesis	100	100
severe, painful motion or weak- ness in the affected extremity With intermediate degrees of resid- ual weakness, pain or limitation	60	50
of motion, rate by analogy to diagnostic codes 5200 and 5203.  Minimum rating	30	20
joint:  For 1 year following implantation of prosthesis	100	100
ness in the affected extremity With intermediate degrees of resid-	50	40
ual weakness, pain or limitation of motion rate by analogy to di- agnostic codes 5205 through 5208.		
Minimum evaluation	30	20
prosthesis	100	100
ness in the affected extremity With intermediate degrees of residual weakness, pain or limitation of motion, rate by analogy to diagnostic code 5214.	40	30
Minimum rating	20	20
prosthesis		100

## PROSTHETIC IMPLANTS—Continued

	Rat	ing
	Major	Mino
Following implantation of prosthesis		
with painful motion or weakness		
such as to require the use of		
crutches		1 90
Markedly severe residual weak-		
ness, pain or limitation of motion		
following implantation of pros-		7/
thesis		70
Moderately severe residuals of weakness, pain or limitation of		
motion		50
Minimum rating		30
5055 Knee replacement (prosthesis).		
Prosthetic replacement of knee joint:		
For 1 year following implantation of		
prosthesis		100
With chronic residuals consisting of		
severe painful motion or weak-		
ness in the affected extremity		60
With intermediate degrees of resid-		
ual weakness, pain or limitation		
of motion rate by analogy to di-		
agnostic codes 5256, 5261, or		
5262.		30
Minimum rating5056 Ankle replacement (prosthesis).		30
5056 Ankle replacement (prosthesis).  Prosthetic replacement of ankle joint:		
For 1 year following implantation of		
prosthesis		100
With chronic residuals consisting of		
severe painful motion or weak-		
ness		40
With intermediate degrees of resid-		
ual weakness, pain or limitation		
of motion rate by analogy to		
5270 or 5271.		
Minimum rating		20
NOTE (1): The 100 pct rating for 1 year		
following implantation of prosthesis will commence after initial grant of the		
1-month total rating assigned under		
§ 4.30 following hospital discharge.		
NOTE (2): Special monthly compensa-		
tion is assignable during the 100 pct		
rating period the earliest date perma-		
nent use of crutches is established.		
COMBINATIONS OF DISABILITIES		
5104 Anatomical loss of one hand and loss		
of use of one foot		1100
5105 Anatomical loss of one foot and loss		
of use of one hand		1100
5106 Anatomical loss of both hands		1100
5107 Anatomical loss of both feet		<sup>1</sup> 100
5108 Anatomical loss of one hand and one		
foot		1100
5109 Loss of use of both hands		1100
5110 Loss of use of both feet		1100
5111 Loss of use of one hand and one		
foot		<sup>1</sup> 100

<sup>&</sup>lt;sup>1</sup> Also entitled to special monthly compensation.

TABLE II—RATINGS FOR MULTIPLE LOSSES OF EXTREMITIES WITH DICTATOR'S RATING CODE AND 38 CFR CITATION

		· ·	or it on the						
	Impairment of other extremity								
Impairment of one extremity	Anatomical loss or loss of use below elbow	Anatomical loss or loss of use below knee	Anatomical loss or loss of use above elbow (preventing use of prosthesis)	Anatomical loss or loss of use above knee (preventing use of prosthesis)	Anatomical loss near shoulder (preventing use of prosthesis)	Anatomical loss near hip (pre- venting use of prosthesis)			
Anatomical loss or loss of use below elbow.	M Codes M-1 a, b, or c, 38 CFR 3.350 (c)(1)(i).	L Codes L-1 d, e, f, or g, 38 CFR 3.350(b).	M½ Code M-5, 38 CFR 3.350 (f)(1)(x).	L½ Code L-2 c, 38 CFR 3.350 (f)(1)(vi).	N Code N-3, 38 CFR 3.350 (f)(1)(xi).	M Code M-3 c, 38 CFR 3.350 (f)(1)(viii)			
Anatomical loss or loss of use below knee.		L Codes L-1 a, b, or c, 38 CFR 3.350(b).	L½ Code L-2 b, 38 CFR 3.350	L½ Code L-2 a, 38 CFR 3.350 (f)(1)(i).	M Code M–3 b, 38 CFR 3.350	M Code M-3 a, 38 CFR 3.350 (f)(1)(ii)			
Anatomical loss or loss of use above elbow			(f)(1)(iii). N Code N-1, 38 CFR 3.350 (d)(1).	M Code M-2 a, 38 CFR 3.350	(f)(1)(iv). N½ Code N-4, 38 CFR 3.350	M½ Code M-4 c, 38 CFR 3.350			
(preventing use of prosthesis). Anatomical loss or loss of use above knee (preventing use				(c)(1)(iii).  M Code M-2 a, 38 CFR 3.350 (c)(1)(ii).	(f)(1)(ix). M½ Code M-4 b, 38 CFR 3.350 (f)(1)(vii).	(f)(1)(xi) M½ Code M-4 a, 38 CFR 3.350 (f)(1)(v)			
of prosthesis). Anatomical loss near shoulder (preventing use					O Code O-1, 38 CFR 3.350 (e)(1)(i).	N Code N-2 b, 38 CFR 3.350 (d)(3)			
of prosthesis). Anatomical loss near hip (pre- venting use of prosthesis).						N Code N-2 a, 38 CFR 3.350 (d)(2)			

Note.—Need for aid attendance or permanently bedridden qualifies for subpar. L. Code L–1 h, i (38 CFR 3.350(b)). Paraplegia with loss of use of both lower extremities and loss of anal and bladder sphincter control qualifies for subpar. O. Code O–2 (38 CFR 3.350(e)(2)). Where there are additional disabilities rated 50% or 100%, or anatomical or loss of use of a third extremity see 38 CFR 3.350(f) (3), (4) or (5).

# (Authority: 38 U.S.C. 1115)

## AMPUTATIONS: UPPER EXTREMITY

		Rati	ing
		Major	Minor
A	rm, amputation of:		
5120	Disarticulation	1 90	1 90
5121	Above insertion of deltoid	1 90	1 80
5122	Below insertion of deltoid	180	170
F	orearm, amputation of:		
5123	Above insertion of pronator teres	<sup>1</sup> 80	<sup>1</sup> 70
5124	Below insertion of pronator teres	170	<sup>1</sup> 60
5125	Hand, loss of use of	170	<sup>1</sup> 60
	MULTIPLE FINGER AMPUTATIONS		
5126	Five digits of one hand, amputation		
of .		170	<sup>1</sup> 60
F	our digits of one hand, amputation of:		
5127	Thumb, index, long and ring	170	<sup>1</sup> 60
5128	Thumb, index, long and little	170	<sup>1</sup> 60
5129	Thumb, index, ring and little	170	<sup>1</sup> 60
5130	Thumb, long, ring and little	170	<sup>1</sup> 60
5131	Index, long, ring and little	60	50
Т	hree digits of one hand, amputation of:		
5132	Thumb, index and long	60	50
5133	Thumb, index and ring	60	50
5134	Thumb, index and little	60	50
5135	Thumb, long and ring	60	50
5136	Thumb, long and little	60	50

## AMPUTATIONS: UPPER EXTREMITY—Continued

		Rati	ng
		Major	Minor
5137	Thumb, ring and little	60	50
5138	Index, long and ring	50	40
5139	Index, long and little	50	40
5140	Index, ring and little	50	40
5141	Long, ring and little	40	30
T	wo digits of one hand, amputation of:		
5142	Thumb and index	50	40
5143	Thumb and long	50	40
5144	Thumb and ring	50	40
5145	Thumb and little	50	40
5146	Index and long	40	30
5147	Index and ring	40	30
5148	Index and little	40	30
5149	Long and ring	30	20
5150	Long and little	30	20
5151	Ring and little	30	20
(a	a) The ratings for multiple finger ampu-		
	tations apply to amputations at the		
	proximal interphalangeal joints or		
	through proximal phalanges		
(b	) Amputation through middle pha-		
	langes will be rated as prescribed for		
	unfavorable ankylosis of the fingers	1	

§4.71a

# 38 CFR Ch. I (7-1-07 Edition)

AMPUTATIONS: UPPER EXTREMITY—Continued

	Rati	ing
	Major	Mino
(c) Amputations at distal joints, or through distal phalanges, other than negligible losses, will be rated as prescribed for favorable ankylosis of the fingers.  (d) Amputation or resection of metacarpal bones (more than one-half the bone lost) in multiple fingers injuries will require a rating of 10 percent added to (not combined with) the ratings, multiple finger amputations, subject to the amputation rule applied to the forearm.  (e) Combinations of finger amputations at various levels, or finger amputations with ankylosis or limitation of motion of the fingers will be rated on the basis of the grade of disability, i.e., amputation, unfavorable ankylosis, most representative of the levels or combinations. With an even number of fingers involved, and adjacent grades of disability, select the higher of the two grades.  (f) Loss of use of the hand will be held to exist when no effective function remains other than that which would be equally well served by an amputation stump with a suitable prosthetic applicance.		

5152 Thumb, amputation of:

AMPUTATIONS: UPPER EXTREMITY—Continued

	Rati	ing
	Major	Minor
With metacarpal resection At metacarpophalangeal joint or through	40	30
proximal phalanx	30	20
At distal joint or through distal phalanx 5153 Index finger, amputation of With metacarpal resection (more than	20	20
one-half the bone lost)	30	20
thereto	20	20
Through middle phalanx or at distal joint 5154 Long finger, amputation of:	10	10
With metacarpal resection (more than one-half the bone lost)	20	20
mal interphalangeal joint or proximal thereto	10	10
With metacarpal resection (more than one-half the bone lost)	20	20
mal interphalangeal joint or proximal thereto	10	10
With metacarpal resection (more than one-half the bone lost)	20	20
mal interphalangeal joint or proximal thereto	10	10
ings are the only applicable ratings for amputations of whole or part of single fingers.		

<sup>&</sup>lt;sup>1</sup> Entitled to special monthly compensation.

### SINGLE FINGER AMPUTATIONS




PLATE III

# §4.71a

# 38 CFR Ch. I (7-1-07 Edition)

## AMPUTATIONS: LOWER EXTREMITY

	Rat- ing
Thigh, amputation of:	
5160 Disarticulation, with loss of extrinsic pelvic gir- dle muscles	290
5161 Upper third, one-third of the distance from perineum to knee joint measured from perineum	<sup>2</sup> 80
5162 Middle or lower thirds Leg, amputation of:	<sup>2</sup> 60
5163 With defective stump, thigh amputation recommended	² 60
5164 Amputation not improvable by prosthesis controlled by natural knee action	<sup>2</sup> 60
5165 At a lower level, permitting prosthesis	<sup>2</sup> 40
bones (more than one-half of metatarsal loss) 5167 Foot, loss of use of	<sup>2</sup> 40 <sup>2</sup> 40

# AMPUTATIONS: LOWER EXTREMITY—Continued

	Rat- ing
5170 Toes, all, amputation of, without metatarsal	
loss	30
5171 Toe, great, amputation of:	
With removal of metatarsal head	30
Without metatarsal involvement	10
5172 Toes, other than great, amputation of, with re-	
moval of metatarsal head:	
One or two	20
Without metatarsal involvement	0
5173 Toes, three or four, amputation of, without	
metatarsal involvement:	
Including great toe	20
Not including great toe	10

<sup>&</sup>lt;sup>2</sup> Also entitled to special monthly compensation.

## AMPUTATIONS: LOWER EXTREMITY




PLATE IV

# 38 CFR Ch. I (7-1-07 Edition)

# §4.71a

## THE SHOULDER AND ARM

## THE ELBOW AND FOREARM—Continued

	Rati	ing
	Major	Minor
5200 Scapulohumeral articulation, anky-		
losis of:		
Note: The scapula and humerus move		
as one piece.		
Unfavorable, abduction limited to 25°		
from side	50	40
Intermediate between favorable and un-		
favorable	40	30
Favorable, abduction to 60°, can reach		
mouth and head	30	20
5201 Arm, limitation of motion of:		
To 25° from side	40	30
Midway between side and shoulder		
level	30	20
At shoulder level	20	20
5202 Humerus, other impairment of:		
Loss of head of (flail shoulder)	80	70
Nonunion of (false flail joint)	60	50
Fibrous union of	50	40
Recurrent dislocation of at		
scapulohumeral joint.		
With frequent episodes and guard-		
ing of all arm movements	30	20
With infrequent episodes, and		
guarding of movement only at		
shoulder level	20	20
Malunion of:		
Marked deformity	30	20
Moderate deformity	20	20
5203 Clavicle or scapula, impairment of:		
Dislocation of	20	20
Nonunion of:		
With loose movement	20	20
Without loose movement	10	10
Malunion of	10	10
Or rate on impairment of function of		
contiguous joint.		

# THE ELBOW AND FOREARM

	Rating	
	Major	Minor
5205 Elbow, ankylosis of:		
Unfavorable, at an angle of less than		
50° or with complete loss of		
supination or pronation	60	50
Intermediate, at an angle of more than		
90°, or between 70° and 50°	50	40
Favorable, at an angle between 90° and		
70°	40	30
5206 Forearm, limitation of flexion of:		
Flexion limited to 45°	50	40
Flexion limited to 55°	40	30
Flexion limited to 70°	30	20
Flexion limited to 90°	20	20
Flexion limited to 100°	10	10
Flexion limited to 110°	0	0
5207 Forearm, limitation of extension of:		
Extension limited to 110°	50	40
Extension limited to 100°	40	30
Extension limited to 90°	30	20
Extension limited to 75°	20	20
Extension limited to 60°	10	10
Extension limited to 45°	10	10
5208 Forearm, flexion limited to 100° and		
extension to 45°	20	20
5209 Elbow, other impairment of Flail joint	60	50 t

	Rati	ing
	Major	Minor
Joint fracture, with marked cubitus varus or cubitus valgus deformity or with ununited fracture of head of ra- dius	20	20
5210 Radius and ulna, nonunion of, with		
flail false joint	50	40
Nonunion in upper half, with false movement:		
With loss of bone substance (1 inch (2.5 cms.) or more) and marked		
deformity	40	30
deformity	30	20
Nonunion in lower half	20	20
Malunion of, with bad alignment 5212 Radius, impairment of:	10	10
Nonunion in lower half, with false movement:		
With loss of bone substance (1 inch (2.5 cms.) or more) and marked		
deformity	40	30
deformity	30	20
	20	20
Nonunion in upper half		
Malunion of, with bad alignment 5213 Supination and pronation, impairment	10	10
of: Loss of (bone fusion):		
The hand fixed in supination or		
hyperpronation	40	30
The hand fixed in full pronation	30	20
The hand fixed near the middle of		
the arc or moderate pronation Limitation of pronation:	20	20
Motion lost beyond middle of arc Motion lost beyond last quarter of	30	20
arc, the hand does not approach full pronation	20	20
Limitation of supination: To 30° or less	10	10
NOTE: In all the forearm and wrist injuries, codes 5205 through 5213, mul-	10	10
tiple impaired finger movements due		
to tendon tie-up, muscle or nerve in-		
jury, are to be separately rated and		
combined not to exceed rating for		
loss of use of hand.		

## THE WRIST

	Rating	
	Major	Minor
5214 Wrist, ankylosis of:     Unfavorable, in any degree of palmar flexion, or with ulnar or radial deviation	50 40 30	40 30 20
5215 Wrist, limitation of motion of:  Dorsiflexion less than 15°	10	10
arm	10	10

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND—Continued

THE TIME			-	D-4	
	Rat	_		Rat	
	Major	Minor		Major	Mino
1) For the index, long, ring, and little fingers			(iv) If only the metacarpophalangeal		
(digits II, III, IV, and V), zero degrees of			or proximal interphalangeal joint		
flexion represents the fingers fully ex-			is ankylosed, and there is a gap of two inches (5.1 cm.) or less		
tended, making a straight line with the rest			between the fingertip(s) and the		
of the hand. The position of function of the			proximal transverse crease of the		
hand is with the wrist dorsiflexed 20 to 30			palm, with the finger(s) flexed to		
degrees, the metacarpophalangeal and proximal interphalangeal joints flexed to			the extent possible, evaluate as		
30 degrees, and the thumb (digit I) ab-			favorable ankylosis		
ducted and rotated so that the thumb pad			(4) Evaluation of ankylosis of the thumb:		
faces the finger pads. Only joints in these			(i) If both the carpometacarpal and		
positions are considered to be in favorable			interphalangeal joints are		
position. For digits II through V, the			ankylosed, and either is in exten-		
metacarpophalangeal joint has a range of			sion or full flexion, or there is ro-		
zero to 90 degrees of flexion, the proximal			tation or angulation of a bone,		
interphalangeal joint has a range of zero			evaluate as amputation at metacarpophalangeal joint or		
to 100 degrees of flexion, and the distal			through proximal phalanx		
(terminal) interphalangeal joint has a			(ii) If both the carpometacarpal and		
range of zero to 70 or 80 degrees of flex-			interphalangeal joints are		
ion			ankylosed, evaluate as unfavor-		
<ol><li>When two or more digits of the same hand are affected by any combination of</li></ol>			able ankylosis, even if each joint		
amputation, ankylosis, or limitation of mo-			is individually fixed in a favorable		
tion that is not otherwise specified in the			position		
rating schedule, the evaluation level as-			(iii) If only the carpometacarpal or		
signed will be that which best represents			interphalangeal joint is		
the overall disability (i.e., amputation, un-			ankylosed, and there is a gap of		
favorable or favorable ankylosis, or limita-			more than two inches (5.1 cm.) between the thumb pad and the		
tion of motion), assigning the higher level			fingers, with the thumb attempt-		
of evaluation when the level of disability is			ing to oppose the fingers, evalu-		
equally balanced between one level and			ate as unfavorable ankylosis		l
the next higher level			(iv) If only the carpometacarpal or		
3) Evaluation of ankylosis of the index,			interphalangeal joint is		
long, ring, and little fingers:			ankylosed, and there is a gap of		
<ul><li>(i) If both the metacarpophalangeal and proximal interphalangeal</li></ul>			two inches (5.1 cm.) or less be-		
joints of a digit are ankylosed,			tween the thumb pad and the fin-		
and either is in extension or full			gers, with the thumb attempting		
flexion, or there is rotation or an-			to oppose the fingers, evaluate		
gulation of a bone, evaluate as			as favorable ankylosis(5) If there is limitation of motion of two or		
amputation without metacarpal			more digits, evaluate each digit separately		
resection, at proximal inter-			and combine the evaluations		
phalangeal joint or proximal					
thereto			I. Multiple Digits: Unfavorable Ank	cylosis	
(ii) If both the metacarpophalangeal			5040 5" " " ( )   ( )		
and proximal interphalangeal			5216 Five digits of one hand, unfavorable		, ا
joints of a digit are ankylosed,			ankylosis of	60	5
evaluate as unfavorable anky-			amputation is warranted.		
losis, even if each joint is individ- ually fixed in a favorable position.			5217 Four digits of one hand, unfavorable		
(iii) If only the metacarpophalangeal			ankylosis of:		
or proximal interphalangeal joint			Thumb and any three fingers	60	
is ankylosed, and there is a gap			Index, long, ring, and little fingers	50	4
of more than two inches (5.1			Note: Also consider whether evaluation as		
cm.) between the fingertip(s) and			amputation is warranted.		
the proximal transverse crease of			5218 Three digits of one hand, unfavorable		
the palm, with the finger(s) flexed			ankylosis of:		
to the extent possible, evaluate			Thumb and any two fingers	50	4
as unfavorable ankylosis	l	l	Index, long, and ring; index, long, and little; or index, ring, and little		
			fingers	40	] 3
			Long, ring, and little fingers	30	
			Note: Also consider whether evaluation as		1
			amputation is warranted.		
			5219 Two digits of one hand, unfavorable		
			ankylosis of:	I	l
			Thumb and any finger	40	3

# 38 CFR Ch. I (7-1-07 Edition)

# §4.71a

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND—Continued

	Rati	ing
	Major	Minor
Index and long; index and ring; or index and little fingers	30	20
ring and little fingers  Note: Also consider whether evaluation as amputation is warranted.	20	20
II. Multiple Digits: Favorable Anky	losis	
5220 Five digits of one hand, favorable ankylosis of	50	40
Thumb and any three fingers	50	40
Index, long, ring, and little fingers 5222 Three digits of one hand, favorable ankylosis of:	40	30
Thumb and any two fingers Index, long, and ring; index, long, and little; or index, ring, and little	40	30
fingers	30	20
Long, ring and little fingers	20	20
Thumb and any finger	30	20
Index and long; index and ring; or index and little fingers	20	20
ring and little fingers	10	10
III. Ankylosis of Individual Dig	its	
5224 Thumb, ankylosis of:		
Unfavorable	20	20
Favorable	10	10
Note: Also consider whether evaluation as amputation is warranted and whether an additional evaluation is warranted for re- sulting limitation of motion of other digits or interference with overall function of the hand.		
5225 Index finger, ankylosis of:		
Unfavorable or favorable	10	10
Note: Also consider whether evaluation as amputation is warranted and whether an additional evaluation is warranted for re- sulting limitation of motion of other digits or interference with overall function of the hand.		
5226 Long finger, ankylosis of: Unfavorable or favorable	10	10
Note: Also consider whether evaluation as amputation is warranted and whether an additional evaluation is warranted for resulting limitation of motion of other digits or interference with overall function of the hand.	10	. 10
nanu.	1	
5227 Ring or little finger, ankylosis of:		

EVALUATION OF ANKYLOSIS OR LIMITATION OF MOTION OF SINGLE OR MULTIPLE DIGITS OF THE HAND—Continued

		Rati	ing
		Major	Mino
amputation is wa additional evalua sulting limitation	er whether evaluation as arranted and whether an tion is warranted for re- of motion of other digits ith overall function of the		
IV. Limitati	on of Motion of Individua	al Digits	
	ation of motion:		
(5.1 cm.) and the attemptir With a ga (2.5 to	of more than two inches ) between the thumb pad fingers, with the thumb go to oppose the fingers p of one to two inches 5.1 cm.) between the ad and the fingers, with	20	21
the thun the finge	nb attempting to oppose ersp of less than one inch	10	10
and the attemptir 5229 Index or long	) between the thumb pad fingers, with the thumb ng to oppose the fingers g finger, limitation of mo-	0	,
more be the proxi the palm	o of one inch (2.5 cm.) or between the fingertip and imal transverse crease of the with the finger flexed to ont possible, or; with ex-		
degrees With a ga (2.5 cm and the crease o	limited by more than 30	10	11
and; ext more tha	tension is limited by no an 30 degrees	0	,
tion: Any limitati	ion of motion	0	,
	THE SPINE		
			Rat-
General Rating Fo	rmula for Diseases and I	niuries	9
	of the Spine	-	
evaluated unde Intervertebral Dis tating Episodes):	les 5235 to 5243 unlesser the Formula for c Syndrome Based on Ir	Rating ncapaci-	
(whther aching ir residuals	vithout symptoms such a or not it radiates), stiffn n the area of the spine affe s of injury or disease	ess, or ected by	
	nfavorable ankylosis of the spinenfavorable ankylosis of the		10
U	thoracolumbar spine		5

THE SPINE—Continued

## THE SPINE—Continued

	Rat- ing		Rat- ing
Unfavorable ankylosis of the entire cervical spine; or, forward flexion of the thoracolumbar spine 30 degrees or less; or, favorable ankylosis of the entire thoracolumbar spine		Note (2): (See also Plate V.) For VA compensation purposes, normal forward flexion of the cervical spine is zero to 45 degrees, extension is zero to 45 degrees, left and right lateral flexion are zero to 45 degrees, and left and right lateral rotation are zero to 80 degrees. Normal forward flexion of the thoracolumbar spine is zero to 90 degrees, extension is zero to 30 degrees, left and right lateral flexion are zero to 30 degrees, left and right lateral flexion are zero to 30 degrees, and left and right lateral rotation are zero to 30 degrees. The combined range of motion refers to the sum of the range of forward flexion, extension, left and right lateral flexion, and left and right rotation. The normal combined range of motion of the cervical spine is 340 degrees and of the thoracolumbar spine is 240 degrees. The normal ranges of motion for each component of spinal motion provided in this note are the maximum that can be used for calculation of the combined range of motion.  Note (3): In exceptional cases, an examiner may state that because of age, body habitus, neurologic disease, or other factors not the result of disease or injury of the spine, the range of motion of the spine in a particular individual, even though it does not conform to the normal range of motion stated in Note (2). Provided that the examiner supplies an explanation, the examiner's assessment that the range of motion is normal for that individual will be accepted.  Note (4): Round each range of motion measurement to the nearest five degrees.  Note (5): For VA compensation purposes, unfavorable ankylosis is a condition in which the entire cervical spine, the entire thoracolumbar spine, or the entire spine is fixed in flexion or extension, and the ankylosis results in one or more of the following: difficulty walking because of a limited line of vision; restricted opening of the mouth and chewing; breathing limited to diaphragmatic respiration; gastrointestinal symptoms due to pressure of the costal margin on the abdomen; dyspnea or	
muscle spasm, guarding, or lo- calized tenderness not resulting in abnormal gait or abnormal spi- nal contour; or, vertebral body fracture with loss of 50 percent or more of the height	10	dysphagia; atlantoaxial or cervical subluxation or dislocation; or neurologic symptoms due to nerve root stretching. Fixation of a spinal segment in neutral position (zero degrees) always represents favorable ankylosis.  Note (6): Separately evaluate disability of the thoracolumbar and cervical spine segments, ex-	
neurologic abnormalities, including, but not limited to, bowel or bladder impairment, separately, under an appropriate diagnostic code.		cept when there is unfavorable ankylosis of both segments, which will be rated as a single disability. 5235 Vertebral fracture or dislocation 5236 Sacrolilac injury and weakness 5237 Lumbosacral or cervical strain 5238 Spinal stenosis 5239 Spondylolisthesis or segmental instability 5240 Ankylosing spondylitis 5241 Spinal fusion 5242 Degenerative arthritis of the spine (see also diagnostic code 5003) 5243 Intervertebral disc syndrome Evaluate intervertebral disc syndrome (preoperatively or postoperatively) either under the General Rating Formula for Diseases and Injuries of the Spine or under the Formula for Rating Intervertebral Disc Syndrome Based on Incapacitating Episodes, whichever method results in the higher evaluation when all disabilities are combined under § 4.25.	

# §4.71a


# 38 CFR Ch. I (7-1-07 Edition)

## THE SPINE—Continued

	Rat- ing
Formula for Rating Intervertebral Disc Syndrome Based on Incapacitating Episodes	
With incapacitating episodes having a total duration of at least 6 weeks during the past 12 months	60
With incapacitating episodes having a total duration of at least 4 weeks but less than 6 weeks during the past 12 months	40
With incapacitating episodes having a total duration of at least 2 weeks but less than 4 weeks during the past 12 months	20
With incapacitating episodes having a total duration of at least one week but less than 2 weeks during	
the past 12 months	10

## THE SPINE—Continued

	ing
Note (1): For purposes of evaluations under diagnostic code 5243, an incapacitating episode is a period of acute signs and symptoms due to intervertebral disc syndrome that requires bed rest prescribed by a physician and treatment by a physician.	
Note (2): If intervertebral disc syndrome is present in more than one spinal segment, provided that the effects in each spinal segment are clearly distinct, evaluate each segment on the basis of incapacitating episodes or under the General Rating Formula for Diseases and Injuries of the Spine, whichever method results in a higher evaluation for that segment.	






PLATE V
RANGE OF MOTION OF CERVICAL AND THORACOLUMBAR SPINE

# §4.71a

# 38 CFR Ch. I (7-1-07 Edition)

## THE HIP AND THIGH

	Rat- ing
5250 Hip, ankylosis of:	
Unfavorable, extremely unfavorable ankylosis,	
the foot not reaching ground, crutches neces-	
sitated	3 90
Intermediate	70
Favorable, in flexion at an angle between 20°	
and 40°, and slight adduction or abduction 5251 Thigh, limitation of extension of:	60
Extension limited to 5°	10
5252 Thigh, limitation of flexion of:	10
Flexion limited to 10°	40
Flexion limited to 20°	30
Flexion limited to 30°	20
Flexion limited to 45°	10
5253 Thigh, impairment of:	
Limitation of abduction of, motion lost beyond	
10°	20
Limitation of adduction of, cannot cross legs	10
Limitation of rotation of, cannot toe-out more	
than 15°, affected leg	10
5254 Hip, flail joint	80
5255 Femur, impairment of:	
Fracture of shaft or anatomical neck of:	
With nonunion, with loose motion (spiral or oblique fracture)	80
With nonunion, without loose motion.	00
weightbearing preserved with aid of brace	60
Fracture of surgical neck of, with false joint	60
Malunion of:	30
With marked knee or hip disability	30
With moderate knee or hip disability	20
With slight knee or hip disability	10

<sup>&</sup>lt;sup>3</sup>Entitled to special monthly compensation.

## THE KNEE AND LEG

	Rat- ing
5256 Knee, ankylosis of:	
Extremely unfavorable, in flexion at an angle of	
45° or more	60
In flexion between 20° and 45°	50
In flexion between 10° and 20°	40
Favorable angle in full extension, or in slight	
flexion between 0° and 10°	30
5257 Knee, other impairment of:	
Recurrent subluxation or lateral instability:	
Severe	30
Moderate	20
Slight	10
5258 Cartilage, semilunar, dislocated, with frequent	
episodes of "locking," pain, and effusion into the	
joint	20
5259 Cartilage, semilunar, removal of, symptomatic	10
5260 Leg, limitation of flexion of:	
Flexion limited to 15°	30
Flexion limited to 30°	20
Flexion limited to 45°	10
Flexion limited to 60°	0
5261 Leg, limitation of extension of:	
Extension limited to 45°	50
Extension limited to 30°	40
Extension limited to 20°	30
Extension limited to 15°	20
Extension limited to 10°	10
Extension limited to 5°	0
5262 Tibia and fibula, impairment of:	١
Nonunion of, with loose motion, requiring brace	40
Malunion of:	
With marked knee or ankle disability	30

## THE KNEE AND LEG—Continued

	Rat- ing
With moderate knee or ankle disability	20
With slight knee or ankle disability	10
tively demonstrated)	10

## THE ANKLE

	Rat- ing
5270 Ankle, ankylosis of:	
In plantar flexion at more than 40°, or in	
dorsiflexion at more than 10° or with abduc-	
tion, adduction, inversion or eversion deformity	40
In plantar flexion, between 30° and 40°, or in	
dorsiflexion, between 0° and 10°	30
In plantar flexion, less than 30°	20
5271 Ankle, limited motion of:	
Marked	20
Moderate	10
5272 Subastragalar or tarsal joint, ankylosis of:	
In poor weight-bearing position	20
In good weight-bearing position	10
5273 Os calcis or astragalus, malunion of:	
Marked deformity	20
Moderate deformity	10
5274 Astragalectomy	20

## SHORTENING OF THE LOWER EXTREMITY

	Rat- ing
5275 Bones, of the lower extremity, shortening of:	
Over 4 inches (10.2 cms.)	<sup>3</sup> 60
31/2 to 4 inches (8.9 cms. to 10.2 cms.)	<sup>3</sup> 50
3 to 31/2 inches (7.6 cms. to 8.9 cms.)	40
21/2 to 3 inches (6.4 cms. to 7.6 cms.)	30
2 to 21/2 inches (5.1 cms. to 6.4 cms.)	20
11/4 to 2 inches (3.2 cms. to 5.1 cms.)	10
NOTE: Measure both lower extremities from ante-	
rior superior spine of the ilium to the internal	
malleolus of the tibia. Not to be combined with	
other ratings for fracture or faulty union in the	
same extremity.	

<sup>&</sup>lt;sup>3</sup> Also entitled to special monthly compensation.

## THE FOOT

	ing
5276 Flatfoot, acquired: Pronounced; marked pronation, extreme tender-	
ness of plantar surfaces of the feet, marked inward displacement and severe spasm of the tendo achillis on manipulation, not improved	
by orthopedic shoes or appliances.	
Bilateral	50
Unilateral	30
Severe; objective evidence of marked deformity	
(pronation, abduction, etc.), pain on manipula-	
tion and use accentuated, indication of swell-	
ing on use, characteristic callosities:	
Bilateral	30
Unilateral	20

## **Department of Veterans Affairs**

## THE FOOT—Continued

	Rat- ing
Moderate; weight-bearing line over or medial to great toe, inward bowing of the tendo achillis, pain on manipulation and use of the feet, bilat-	
eral or unilateral	10
Mild; symptoms relieved by built-up shoe or arch support	0
5277 Weak foot, bilateral:  A symptomatic condition secondary to many constitutional conditions, characterized by atrophy of the musculature, disturbed circulation, and weakness:  Rate the underlying condition, minimum rating	10
5278 Claw foot (pes cavus), acquired: Marked contraction of plantar fascia with dropped forefoot, all toes hammer toes, very painful callosities, marked varus deformity:	
Bilateral Unilateral All toes tending to dorsiflexion, limitation of dorsiflexion at ankle to right angle, shortened plantar fascia, and marked tenderness under metatarsal heads:	50 30
Bilateral	30 20
Bilateral	10 10 0
5279 Metatarsalgia, anterior (Morton's disease), unilateral, or bilateral	10
5280 Hallux valgus, unilateral:  Operated with resection of metatarsal head Severe, if equivalent to amputation of great toe  5281 Hallux rigidus, unilateral, severe: Rate as hallux valgus, severe.  Note: Not to be combined with claw foot ratings.	10 10
5282 Hammer toe: All toes, unilateral without claw foot	10 0
nonunion of: Severe	30 20 10
percent. 5284 Foot injuries, other: Severe	30 20 10

## THE SKULL

	Rat- ing
5296 Skull, loss of part of, both inner ar	nd outer ta-
bles:	
With brain hernia	80
Without brain hernia:	
Area larger than size of a 50-ce	nt piece or
1.140 in <sup>2</sup> (7.355 cm <sup>2</sup> )	50
Area intermediate	30
Area smaller than the size of	a 25-cent
piece or 0.716 in 2 (4.619 cm 2)	10

## THE SKULL—Continued

	Rat- ing
NOTE: Rate separately for intracranial complications.	

## THE RIBS

	Rat- ing
5297 Ribs, removal of:	
More than six	50
Five or six	40
Three or four	30
Two	20
One or resection of two or more ribs without regeneration	10

## THE COCCYX

	Rat- ing
5298 Coccyx, removal of:	
Partial or complete, with painful residuals	10
Without painful residuals	0

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 40 FR 42536, Sept. 15, 1975; 41 FR 11294, Mar. 18, 1976; 43 FR 45350, Oct. 2, 1978; 51 FR 6411, Feb. 24, 1986; 61 FR 20439, May 7, 1996; 67 FR 48785, July 26, 2002; 67 FR 54349, Aug. 22, 2002; 68 FR 51456, Aug. 27, 2003; 69 FR 32450, June 10, 2004]

## §4.72 [Reserved]

# § 4.73 Schedule of ratings—muscle injuries.

NOTE: When evaluating any claim involving muscle injuries resulting in loss of use of any extremity or loss of use of both buttocks (diagnostic code 5317, Muscle Group XVII), refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation.

# 38 CFR Ch. I (7-1-07 Edition)

# §4.73

## THE SHOULDER GIRDLE AND ARM

## THE FOREARM AND HAND

	Rating			Rating	
	Domi- nant	Non- domi- nant		Domi- nant	Non- domi- nant
5301 Group I. Function: Upward rotation of scapula; elevation of arm above shoulder level. Extrinsic muscles of shoulder girdle: (1) Trapezius; (2) levator scapulae; (3) serratus magnus.  Severe	40	30	5307 Group VII. Function: Flexion of wrist and fingers. Muscles arising from internal condyle of humerus: Flexors of the carpus and long flexors of fingers and thumb; pronator.  Severe	40	30
Moderately Severe	30	20	Moderately Severe	30	20
Moderate	10	10	Moderate	10	10
Slight	0	0	Slight	0	0
sic muscles of shoulder girdle: (1) Pectoralis major II (costostemal); (2) latissimus dorsi and teres major (teres major, although technically an intrinsic muscle, is included with latissimus dorsi); (2) entendin misor (1) templaid.			Severe Moderately Severe Moderately Severe Slight Forearm Tx.	30 20 10 0	20 20 10 0
(3) pectoralis minor; (4) rhomboid.  Severe	40	30	muscles act in strong grasping move- ments and are supplemented by the intrin-		
Moderately Severe	30	20	sic muscles in delicate manipulative		
Moderate	20	20	movements. Intrinsic muscles of hand:		
Slight	0	0	Thenar eminence; short flexor, opponens, abductor and adductor of thumb; hypothenar eminence; short flexor, opponens and abductor of little finger; 4 lumbricales; 4 dorsal and 3 palmar interossei.		
of shoulder girdle: (1) Pectoralis major I (clavicular); (2) deltoid. Severe	40 30	30 20	NOTE: The hand is so compact a structure that isolated muscle injuries are rare, being nearly always complicated with inju-		
Moderate	20	20	ries of bones, joints, tendons, etc. Rate on		
Slight	0	0	limitation of motion, minimum 10 percent.		
5304 Group IV. Function: Stabilization of shoulder against injury in strong movements, holding head of humerus in socket; abduction; outward rotation and inward ro-			THE FOOT AND LEG		
tation of arm. Intrinsic muscles of shoulder girdle: (1) Supraspinatus; (2) infraspinatus and teres minor; (3) subscapularis; (4)			5310 Group X. Function: Movements of	forefeet	Rat- ing
coracobrachialis.			and toes; propulsion thrust in walking.	Intrinsic	
Severe	30 20	20 20	muscles of the foot: Plantar: (1) Flexor di	gitorum	
Moderately Severe  Moderate	10	20 10	brevis; (2) abductor hallucis; (3) abductor minimi; (4) quadratus plantae; (5) lumbrica	or digiti	
Slight	0	0	flexor hallucis brevis; (7) adductor hallucis;		
5305 Group V. Function: Elbow supination (1) (long head of biceps is stabilizer of shoulder joint); flexion of elbow (1, 2, 3). Flexor muscles of elbow: (1) Biceps; (2) brachialis; (3) brachioradialis.			or digiti minimi brevis; (9) dorsal and interossei. Other important plantar structure tar aponeurosis, long plantar calcaneonavicular ligament, tendons of p tibial, peroneus longus, and long flexors of the control of	plantar s: Plan- and osterior	
Severe	40	30	and little toes. Severe		30
Moderately Severe	30	20	Moderately Severe		20
Moderate Slight	10 0	10 0	Moderate		10
5306 Group VI. Function: Extension of elbow (long head of triceps is stabilizer of shoulder joint). Extensor muscles of the elbow: (1) Triceps; (2) anconeus	0	0	Slight	extensor uctures: its; ten-	0
Severe	40	30	dons of long extensors of toes and peron cles.	eı mus-	
Moderately Severe	30	20	Severe		20
Moderate	10 0	10 0	Moderately Severe		10
Slight					

#### THE FOOT AND LEG—Continued

	Rat- ing
NOTE: Minimum rating for through-and-through wounds of the foot—10.	
5311 Group XI. Function: Propulsion, plantar flexion of foot (1); stabilization of arch (2, 3); flexion of toes (4, 5); Flexion of knee (6). Posterior and lateral crural muscles, and muscles of the calf: (1) Triceps surae (gastrocnemius and soleus); (2) tibialis posterior; (3) peroneus longus; (4) peroneus brevis; (5) flexor hallucis longus; (6) flexor digitorum longus; (7) popliteus; (8) plantaris.  Severe	30 20 10 0
Severe Moderately Severe Moderate Slight	30 20 10 0

#### THE PELVIC GIRDLE AND THIGH

THE TELVIO CHABLE THAT THICH	
	Rat- ing
5313 Group XIII. Function: Extension of hip and flexion of knee; outward and inward rotation of flexed knee; acting with rectus femoris and sartorius (see XIV, 1, 2) synchronizing simultaneous flexion of hip and knee and extension of hip and knee by belt-over-pulley action at knee joint. Posterior thigh group, Hamstring complex of 2-joint muscles: (1) Biceps femoris; (2) semimembranosus; (3) semitendinosus.  Severe Moderately Severe Moderate Slight  5314 Group XIV. Function: Extension of knee (2, 3, 4, 5); simultaneous flexion of hip and flexion of knee (1); tension of fascia lata and iliotibial (Maissiat's) band, acting with XVII (1) in postural support of body (6); acting with hamstrings in synchronizing hip and knee (1, 2). Anterior thigh group: (1) Sartorius; (2) rectus femoris; (3) vastus	40 30 10 0
externus; (4) vastus intermedius; (5) vastus internus; (6) tensor vaginae femoris.  Severe  Moderately Severe  Slight  Soroup XV. Function: Adduction of hip (1, 2, 3, 4); flexion of hip (1, 2); flexion of knee (4). Mesial	40 30 10 0
thigh group: (1) Adductor longus; (2) adductor brevis; (3) adductor magnus; (4) gracilis.  Severe  Moderately Severe  Moderate  Slight  5316 Group XVI. Function: Flexion of hip (1, 2, 3).  Pelvic girdle group 1: (1) Psoas; (2) iliacus; (3)	30 20 10 0
pectineus. Severe	40 30 10 0

#### THE PELVIC GIRDLE AND THIGH—Continued

	Rat- ing
5317 Group XVII. Function: Extension of hip (1); abduction of thigh; elevation of opposite side of pelvis (2, 3); tension of fascia lata and iliotibial (Maissiat's) band, acting with XIV (6) in postural support of body steadying pelvis upon head of femur and condyles of femur on tibia (1). Pelvic girdle group 2: (1) Gluteus maximus; (2) gluteus medius; (3) gluteus minimus.  Severe	*50 40 20 0
Severe	30
Moderately Severe	20
Moderate	10
Slight	0

\*If bilateral, see §3.350(a)(3) of this chapter to determine whether the veteran may be entitled to special monthly compensation.

	Rat- ing
5319 Group XIX. Function: Support and compression of abdominal wall and lower thorax; flexion	
and lateral motions of spine; synergists in strong downward movements of arm (1). Muscles of the abdominal wall: (1) Rectus abdominis; (2) external oblique; (3) internal oblique; (4) transversalis; (5)	
quadratus lumborum.	
Severe	50
Moderately Severe	30
Moderate	10
Slight	0
5320 Group XX. Function: Postural support of body;	
extension and lateral movements of spine. Spinal	
muscles: Sacrospinalis (erector spinae and its pro-	
longations in thoracic and cervical regions).	
Cervical and thoracic region:.	
Severe	40
Moderately Severe	20
Moderate	10
Slight Lumbar region:.	0
Severe	60
Moderately Severe	40
Moderate	20
Slight	-0
5321 Group XXI. Function: Respiration. Muscles of	·
respiration: Thoracic muscle group.	
Severe or Moderately Severe	20
Moderate	10
Slight	0
5322 Group XXII. Function: Rotary and forward	
movements of the head; respiration; deglutition.	
Muscles of the front of the neck: (Lateral, supra-,	
and infrahyoid group.) (1) Trapezius I (clavicular in-	
sertion); (2) sternocleidomastoid; (3) the "hyoid" muscles; (4) sternothyroid; (5) digastric.	
, ( )	30
Severe  Moderately Severe	20
	10
Moderate	

THE TORSO AND NECK—Continued

	Rat- ing
5323 Group XXIII. Function: Movements of the head; fixation of shoulder movements. Muscles of the side and back of the neck: Suboccipital; lateral vertebral and anterior vertebral muscles.  Severe  Moderately Severe  Moderate  Slight	30 20 10 0

#### **MISCELLANEOUS**

Rat- ing

5324 Diaphragm, rupture of, with herniation. Rate under diagnostic code 7346.

under diagnostic code 7346. 5325 Muscle injury, facial muscles. Evaluate functional impairment as seventh (facial) cranial nerve neuropathy (diagnostic code 8207), disfiguring scar (diagnostic code 7800), etc. Minimum, if interfering to any extent with mastication—10.

to any extent with mastication—10.
5326 Muscle hernia, extensive. Without other injury to the muscle—10.

5327 Muscle, neoplasm of, malignant (excluding soft tissue sarcoma)—100.

NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residual impairment of function.

5328 Muscle, neoplasm of, benign, postoperative. Rate on impairment of function, i.e., limitation of motion, or scars, diagnostic code 7805, etc.

5329 Sarcoma, soft tissue (of muscle, fat, or fibrous connective tissue)—100.

NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residual impairment of function.

(Authority: 38 U.S.C. 1155) [62 FR 30239, June 3, 1997]

THE ORGANS OF SPECIAL SENSE

#### § 4.75 Examination of visual acuity.

Ratings on account of visual impairments considered for service connection are, when practicable, to be based only on examination by specialists. Such special examinations should include uncorrected and corrected cen-

tral visual acuity for distance and near, with record of the refraction. Snellen's test type or its equivalent will be used. Mydriatics should be routine, except when contraindicated. Funduscopic and ophthalmological findings must be recorded. The best distant vision obtainable after best correction by glasses will be the basis of rating, except in cases keratoconus in which contact lenses are medically required. Also, if there exists a difference of more than 4 diopters of spherical correction between the two eyes, the best possible visual acuity of the poorer eye without glasses, or with a lens of not more than 4 diopters difference from that used with the better eye will be taken as the visual acuity of the poorer eye. When such a difference exists, close attention will be given to the likelihood of congenital origin in mere refractive error.

[40 FR 42537, Sept. 15, 1975]

#### §4.76 Examination of field vision.

Measurement of the visual field will be made when there is disease of the optic nerve or when otherwise indicated. The usual perimetric methods will be employed, using a standard perimeter and 3 mm. white test object. At least 16 meridians 22½ degrees apart will be charted for each eye. (See Figure 1. For the 8 principal meridians, see table III.) The charts will be made a part of the report of examination. Not less than 2 recordings, and when possible, 3 will be made. The minimum limit for this function is established as a concentric central contraction of the visual field to 5°. This type of contraction of the visual field reduces the visual efficiency to zero. Where available the examination for form field should be supplemented, when indicated, by the use of tangent screen or campimeter. This last test is especially valuable in detection of scotoma.

[43 FR 45352, Oct. 2, 1978]

#### § 4.76a Computation of average concentric contraction of visual fields.


The extent of contraction of visual field in each eye is determined by recording the extent of the remaining visual fields in each of the eight 45 degree principal meridians. The number

§4.76a

of degrees lost is determined at each meridian by subtracting the remaining degrees from the normal visual fields given in table III. The degrees lost are then added together to determine total degrees lost. This is subtracted from 500. The difference represents the total remaining degrees of visual field. The difference divided by eight represents the average contraction for rating purposes.

TABLE III—NORMAL VISUAL FIELD EXTENT AT 8
PRINCIPAL MERIDIANS

Meridian	Normal de- grees
Temporally	85
Down temporally	85
Down	65
Down nasally	50
Nasally	60
Up nasally	55
Up	45
Up temporally	55
Total	500



TS-19 52a

Example of computation of concentric contraction under the schedule with abnormal findings taken from Figure 1.

Loss	Degrees	
Temporally	55	
Down temporally	55	
Down	45	

Loss	Degrees
Down nasally	30
Nasally	40
Up nasally	35
Up	25
Up temporally	35
Total loss	320

Remaining field 500° minus 320° =  $180^{\circ}$ .  $180^{\circ} \div 8 = 221/2^{\circ}$  average concentric contraction

[43 FR 45352, Oct. 2, 1978]

#### §4.77 Examination of muscle function.

The measurement of muscle function will be undertaken only when the history and findings reflect disease or injury of the extrinsic muscles of the eye, or of the motor nerves supplying these muscles. The measurement will

be performed using a Goldmann Perimeter Chart as in Figure 2 below. The chart identifies four major quadrants, (upward, downward, and two lateral) plus a central field (20° or less). The examiner will chart the areas in which diplopia exists, and such plotted chart will be made a part of the examination report. Muscle function is considered normal (20/40) when diplopia does not exist within  $40^{\circ}$  in the lateral or downward quadrants, or within 30° in the upward quadrant. Impairment of muscle function is to be supported in each instance by record of actual appropriate pathology. Diplopia which is only occasional or correctable is not considered a disability.




Figure 2. Goldmann Perimeter Chart

52c

[53 FR 30262, Aug. 11, 1988]

#### § 4.78 Computing aggravation.

In determining the effect of aggravation of visual disability, even though the visual impairment of only one eye is service connected, evaluate the vision of both eyes, before and after suffering the aggravation, and subtract the former evaluation from the latter except when the bilateral vision amounts to total disability. In the event of subsequent increase in the disability of either eye, due to intercurrent disease or injury not associated with the service, the condition of the eves before suffering the subsequent increase will be taken as the basis of compensation subject to the provisions of §3.383(a) of this chapter.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45354, Oct. 2, 1978]

## § 4.79 Loss of use of one eye, having only light perception.

Loss of use or blindness of one eye, having only light perception, will be held to exist when there is inability to recognize test letters at 1 foot (.30m.) and when further examination of the eyes reveals that perception of objects, hand movements or counting fingers cannot be accomplished at 3 feet (.91m.), lesser extents of visions, particularly perception of objects, hand movements, or counting fingers at distances less than 3 feet (.91 m.), being considered of negligible utility. With visual acuity 5/200 (1.5/60) or less or the visual field reduced to 5° concentric contraction, in either event in both eyes, the question of entitlement on account of regular aid and attendance will be determined on the facts in the individual case.

[43 FR 45354, Oct. 2, 1978]

#### § 4.80 Rating of one eye.

Combined ratings for disabilities of the same eye should not exceed the amount for total loss of vision of that eye unless there is an enucleation or a serious cosmetic defect added to the total loss of vision.

#### §§ 4.81-4.82 [Reserved]

## §4.83 Ratings at scheduled steps and distances.

In applying the ratings for impairment of visual acuity, a person not having the ability to read at any one of the scheduled steps or distances, but reading at the next scheduled step or distance, is to be rated as reading at this latter step or distance. That is, a person who can read at 20/100 (6/30) but who cannot at 20/70 (6/21), should be rated as seeing at 20/100 (6/30).

[41 FR 34257, Aug. 13, 1976, as amended at 43 FR 45354, Oct. 2, 1978]

## §4.83a Impairment of central visual acuity.

The percentage evaluation will be found from table V by intersecting the horizontal row appropriate for the Snellen index for one eye and the vertical column appropriate to the Snellen index of the other eye. For example, if one eye has a Snellen index of 5/200 (1.5/60) and the other eye has a Snellen index of 20/70 (6/21), the percentage evaluation is found in the third horizontal row from the bottom and the fourth vertical column from the left. The evaluation is 50 percent and the diagnostic code 6073.

[41 FR 11297, Mar. 18, 1976, as amended at 43 FR 45354, Oct. 2, 1978]

## §4.84 Differences between distant and near visual acuity.

Where there is a substantial difference between the near and distant corrected vision, the case should be referred to the Director, Compensation and Pension Service.

[40 FR 42537, Sept. 15, 1975]

#### § 4.84a Schedule of ratings—eye.

#### DISEASES OF THE EYE

		Rat- ing
6000	Uveitis	
6001	Keratitis	
6002	Scleritis	
6003	Iritis	
6004	Cyclitis	
6005	Choroiditis	
6006	Retinitis	
6007	Hemorrhage, intra-ocular, recent	
6008	Retina, detachment of	
6009	Eve. injury of, unhealed:	

### §4.84a

#### DISEASES OF THE EYE-Continued

### DISEASES OF THE EYE—Continued

		BIGEAGES OF THE ETE CONTINUES	
	Rat- ing		Rat- ing
The above disabilities, in chronic form, are to be		Unileteral	10
rated from 10 percent to 100 percent for im-		Unilateral	10
pairment of visual acuity or field loss, pain,		6026 Neuritis, optic:	
rest-requirements, or episodic incapacity, com-		Rate underlying disease, and combine impair-	
bining an additional rating of 10 percent during		ment of visual acuity or field loss.	
continuance of active pathology. Minimum rat-		6027 Cataract, traumatic:	
ing during active pathology	10	Preoperative.	
6010 Eye, tuberculosis of, active or inactive:		Rate on impairment of vision.	
Active	100	Postoperative.	
Inactive: See §§ 4.88b and 4.89.		Rate on impairment of vision and aphakia.	
6011 Retina, localized scars, atrophy, or irregular-		6028 Cataract, senile, and others:	
ities of, centrally located, with irregular, duplicated		Preoperative.	
enlarged or diminished image:		Rate on impairment of vision.	
Unilateral or bilateral	10		
6012 Glaucoma, congestive or inflammatory:		Postoperative.	
Frequent attacks of considerable duration; during		Rate on impairment of vision and aphakia.	
continuance of actual total disability	100	6029 Aphakia:	
Or, rate as iritis, diagnostic Code 6003.		Bilateral or unilateral	30
6013 Glaucoma, simple, primary, noncongestive:		NOTE: The 30 percent rating prescribed for	
Rate on impairment of visual acuity or field loss.		aphakia is a minimum rating to be applied to	
Minimum rating	10	the unilateral or bilateral condition and is not	
6014 New growths, malignant (eyeball only):		to be combined with any other rating for im-	
Pending completion of operation or other indi-		paired vision. When only one eye is aphakic,	
cated treatment	100	the eye having poorer corrected visual acuity	
Healed; rate on residuals.		will be rated on the basis of its acuity without	
6015 New growths, benign (eyeball and adnexa,		correction. When both eyes are aphakic, both	
other than superficial)		will be rated on corrected vision. The cor-	
Rate on impaired vision, minimum	10	rected vision of one or both aphakic eyes will	
Healed; rate on residuals.		be taken one step worse than the ascertained	
6016 Nystagmus, central	10	value, however, not better than 20/70 (6/21).	
6017 Conjunctivitis, trachomatous, chronic:		Combined ratings for disabilities of the same	
Active; rate for impairment of visual acuity; min-		eye should not exceed the amount for total	
imum rating while there is active pathology	30	loss of vision of that eye unless there is an	
Healed; rate on residuals, if no residuals	0	enucleation or a serious cosmetic defect	
6018 Conjunctivitis, other, chronic:		added to the total loss of vision.	
Active, with objective symptoms	10	6030 Accommodation, paralysis of	20
Healed; rate on residuals, if no residuals	0	6031 Dacryocystitis	
6019 Ptosis, unilateral or bilateral:		Rate as epiphora.	
Pupil wholly obscured.		6032 Eyelids, loss of portion of:	
Rate equivalent to 5/200 (1.5/60). Pupile one-half or more obscured.		Rate as disfigurement. (See diseases of the	
Rate equivalent to 20/100 (6/30).		skin.)	
With less interference with vision.		6033 Lens, crystalline, dislocation of:	
Rate as disfigurement.		Rate as aphakia.	
6020 Ectropion:		6034 Pterygium:	
Bilateral	20	Rate for loss of vision, if any.	
Unilateral	10		
6021 Entropion:	'0	6035 Keratoconus: To be evaluated on impairment	
Bilateral	20	of corrected visual acuity using contact lenses.	
Unilateral	10	NOTE: When contact lenses are medically re-	
6022 Lagophthalmos:	"	quired for keratoconus, either unilateral or bi-	
Bilateral	20	lateral, the minimum rating will be 30 percent.	
Unilateral	10		
6023 Eyebrows, loss of, complete, unilateral or bi-	'		
lateral	10		
6024 Eyelashes, loss of, complete, unilateral or bi-			
lateral	10		
6025 Epiphora (lacrymal duct, interference with,			
from any cause):			
Bilateral	20		

TABLE IV—TABLE FOR RATING BILATERAL BLINDNESS OR BLINDNESS COMBINED WITH HEARING LOSS WITH DICTATOR'S CODE AND 38 CFR CITATIONS

	Vision other eye			Plus service-connected Hearing loss				
Vision one eye	5/200 (1.5/60) or less	Light perception only	No light perception or anatomical loss	Total deafness one ear	10% or 20% at least one ear SC	30% at least one ear SC	40% at least one ear SC	60% or more at least one ear SC
5/200 (1.5/60) or less.	L <sup>1</sup> Code LB-1 38 CFR 3.350(b)(2).	L+½¹ Code LB-2 38 CFR 3.350(f)(2)(i).	M Code MB–2 a or b 38 CFR 3.350(f)(2)(ii).	Add ½ step Code PB-1 38 CFR 3.350(f)(2)(iv).	No additional SMC.	Add a full step Code PB-3 38 CFR 3.350(f)(2)(vi).	Add a full step Code PB-3 38 CFR 3.350(f)(2)(vi).	O Code OB-1 38 CFR 3.350(e)(1)(iii)
Light perception only.		M Code MB-1 a 38 CFR 3.350(c)(1)((iv).	M+½ Code MB-3 a or b 38 CFR 3.350(f)(iii).	O Code OB-2 38 CFR 3.350(e)(1)(iv).	Add ½ step Code PB-2 38 CFR 3.350(f)(2)(v).	Add a full step Code PB-3 38 CFR 3.350(f)(2)(iv).	O Code OB-2 38 CFR 3.350(e)(1)(iv).	O Code OB-1 38 CFR 3.350(e)(1)(iii)
No light perception or anatomical loss.			N Code NB-1 a-b or c 38 CFR 3.350(d)(4).	O Code OB-2 38 CFR 3.350(e)(1)(iv).	Add ½ step Code PB-2 38 CFR 3.350(f)(2)(v).	Add full step Code PB-3 38 CFR 3.350(f)(2)(vi).	O Code OB-2 38 CFR 3.350(e)(1)(iv).	O Code OB-1 38 CFR 3.350(e)(1)(iii)

<sup>&</sup>lt;sup>1</sup>With need for aid and attendance qualifies for Subpar. m. code MB-1, b; 38 CFR 3.350(c)(1)(v).

NOTE. (1) Any of the additional SMC payable under Dictator's Codes PB-1, PB-2, or PB-3 is not to exceed the rate payable under Subpar. O. (2) If in addition to any of the above the veteran has the service-connected loss or loss of use of an extremity, additional SMC is payable, not to exceed the rate payable under Subpar. O. See Dictator's Codes PB-4, PB-5, PB-6, and 38 CFR 3.350(f)(2)(viii) (A), (B), (C).

**Department of Veterans Affairs** 

### (Authority: 38 U.S.C. 1115) IMPAIRMENT OF CENTRAL VISUAL ACUITY

		Rat- ing
6061	Anatomical loss both eyes	5 100
6062	Blindness in both eyes having only light per-	
	tion	5 100
Α	natomical loss of 1 eye:	
6063	In the other eye 5/200 (1.5/60)	5 100
6064	In the other eye 10/200 (3/60)	6 90
6064	In the other eye 15/200 (4.5/60)	6 80
6064	In the other eye 20/200 (6/60)	670
6065	In the other eye 20/100 (6/30)	6 60
6065	In the other eye 20/70 (6/21)	6 60
6065	In the other eye 20/50 (6/15)	<sup>6</sup> 50
6066	In the other eye 20/40 (6/12)	<sup>6</sup> 40
	lindness in 1 eye, having only light perception:	
6067	In the other eye 5/200 (1.5/60)	5 100
6068	In the other eye 10/200 (3/60)	5 90
6068	In the other eye 15/200 (4.5/60)	5 80
6068	In the other eye 20/200 (6/60)	<sup>5</sup> 70
6069	In the other eye 20/100 (6/30)	5 60
6069	In the other eye 20/70 (6/21)	<sup>5</sup> 50
6069	In the other eye 20/50 (6/15)	5 40
6070	In the other eye 20/40 (6/12)	5 30
V	ision in 1 eye 5/200 (1.5/60):	
6071	In the other eye 5/200 (1.5/60)	5 100
6072	In the other eye 10/200 (3/60)	90
6072	In the other eye 15/200 (4.5/60)	80
6072	In the other eye 20/200 (6/60)	70
6073	In the other eye 20/100 (6/30)	60
6073	In the other eye 20/70 (6/21)	50
6073	In the other eye 20/50 (6/15)	40
6074	In the other eye 20/40 (6/12)	30
	ision in 1 eye 10/200 (3/60):	
6075	In the other eye 10/200 (3/60)	90
6075	In the other eye 15/200 (4.5/60)	80
6075	In the other eye 20/200 (6/60)	70

#### IMPAIRMENT OF CENTRAL VISUAL ACUITY— Continued

§4.84a

		Rat- ing
6076	In the other eye 20/100 (6/30)	60
6076	In the other eye 20/70 (6/21)	50
6076	In the other eye 20/50 (6/15)	40
6077	In the other eye 20/40 (6/12)	30
V	ision in 1 eye 15/200 (4.5/60):	
6075	In the other eye 15/200 (4.5/60)	80
6075	In the other eye 20/200 (6/60)	70
6076	In the other eye 20/100 (6/30)	60
6076	In the other eye 20/70 (6/21)	40
6076	In the other eye 20/50 (6/15)	30
6077		20
V	ision in 1 eye 20/200 (6/60):	
6075	In the other eye 20/200 (6/60)	70
6076	In the other eye 20/100 (6/30)	60
6076	In the other eye 20/70 (6/21)	40
6076	In the other eye 20/50 (6/15)	30
6077	In the other eye 20/40 (6/12)	20
	ision in 1 eye 20/100 (6/30):	
6078	In the other eye 20/100 (6/30)	50
6078	In the other eye 20/70 (6/21)	30
6078	In the other eye 20/50 (6/15)	20
6079		10
	ision in 1 eye 20/70 (6/21):	
6078	In the other eye 20/70 (6/21)	30
6078	, - = -, - = -, (-,,	20
6079	In the other eye 20/40 (6/12)	10
	ision in 1 eye 20/50 (6/15):	
6078		10
6079		10
	ision in 1 eye 20/40 (6/12):	_
In	the other eye 20/40 (6/12)	0

TABLE V—RATINGS FOR CENTRAL VISUAL ACUITY IMPAIRMENT [With Diagnostic Code]

	Vision in other eye												
Vision in one eye	20/40 (6/ 12)	20/50 (6/ 15)	20/70 (6/ 21)	20/100 (6/30)	20/200 (6/60)	15/200 (4.5/60)	10/200 (3/60)	5/200 (1.5/60)	Light perception only/anatomical loss				
20/40 (6/12)	0												
20/50 (6/15)	10 (6079)	10 (6078)											
20/70 (6/21)	10 (6079)	20 (6078)	30 (6078)										
20/100 (6/30)	10 (6079)	20 (6078)	30 (6078)	50 (6078)									
20/200 (6/60)	20 (6077)	30 (6076)	40 (6076)	60 (6076)	70 (6075)								
15/200 (4.5/60)	20 (6077)	30 (6076)	40 (6076)	60 (6076)	70 (6075)	80 (6075)							
10/200 (3/60)	30 (6077)	40 (6076)	50 (6076)	60 (6076)	70 (6075)	80 (6075)	90 (6075)						
5/200 (1.5/60)	30 (6074)	40 (6073)	50 (6073)	60 (6073)	70 (6072)	80 (6072)	90 (6072)	<sup>5</sup> 100 (6071)					

<sup>&</sup>lt;sup>5</sup> Also entitled to special monthly compensation. <sup>6</sup> Add 10% if artificial eye cannot be worn; also entitled to special monthly compensation.

### §4.84a

#### TABLE V—RATINGS FOR CENTRAL VISUAL ACUITY IMPAIRMENT—Continued [With Diagnostic Code]

		Vision in other eye												
Vision in one eye	20/40 (6/ 12)	20/50 (6/ 15)	20/70 (6/ 21)	20/100 (6/30)	20/200 (6/60)	15/200 (4.5/60)	10/200 (3/60)	5/200 (1.5/60)	Light percep- tion only/ana- tomical loss					
Light per- ception														
only	<sup>5</sup> 30 (6070)	<sup>5</sup> 40 (6069)	<sup>5</sup> 50 (6069)	<sup>5</sup> 60 (6069)	<sup>5</sup> 70 (6068)	<sup>5</sup> 80 (6068)	<sup>5</sup> 90 (6068)	<sup>5</sup> 100 (6067)	<sup>5</sup> 100 (6062)					
Anatomical loss of														
one eye	<sup>6</sup> 40 (6066)	<sup>6</sup> 50 (6065)	<sup>6</sup> 60 (6065)	<sup>6</sup> 60 (6065)	<sup>6</sup> 70 (6064)	<sup>6</sup> 80 (6064)	<sup>6</sup> 90 (6064)	<sup>5</sup> 100 (6063)	<sup>5</sup> 100 (6061)					

Rat-ing

30 30 10

20 10

100 30

70 20

50 10

30

10

20 10

#### RATINGS FOR IMPAIRMENT OF FIELD VISION

6080 Field vision, impairment of:
Homonymous hemianopsia
Field, visual, loss of temporal half:
Bilateral
Unilateral
Or rate as 20/70 (6/21).
Field, visual, loss of nasal half:
Bilateral
Unilateral
Or rate as 20/50 (6/15).
Field, visual, concentric contraction of:
To 5°:
Bilateral
Unilateral
Or rate as 5/200 (1.5/60).
To 15° but not to 5°:
Bilateral
Unilateral
Or rate as 20/200 (6/60).
To 30° but not to 15°:
Bilateral
Unilateral
Or rate as 20/100 (6/30).
To 45° but not to 30°:
Bilateral
Unilateral
Or rate as 20/70 (6/21):
To 60° but not to 45°:
Bilateral
Unilateral
Or rate as 20/50 (6/15).
Note (1): Correct diagnosis reflecting disease or
injury should be cited

#### RATINGS FOR IMPAIRMENT OF FIELD VISION— Continued

	Rat- ing
Note (2): Demonstrable pathology commensurate with the functional loss will be required. The concentric contraction ratings require contraction within the stated degrees, temporally; the nasal contraction may be less. The alternative ratings are to be employed when there is ratable defect of visual acuity, or a different impairment of the visual field in the other eye. Concentric contraction resulting from demonstrable pathology to 5 degrees or less will be considered on a parity with reduction of central visual acuity to 5/200 (1.5/60) or less for all purposes including entitlement under §3.350(b)(2) of this chapter; not however, for the purpose of §3.350(a) of this chapter. Entitlement on account of blindness requiring regular aid and attendance, §3.350(c) of this chapter, will continue to be determined on the facts in the individual case	
Large or centrally located, minimum	10

#### RATINGS FOR IMPAIRMENT OF MUSCLE FUNCTION [6090 Diplopia (double vision)]

Degree of diplopia						
(a) Central 20°	5/200					
(b) 21° to 30°:						
(1) Down	15/200					
(2) Lateral	20/100					
(3) Up	20/70					
(c) 31° to 40°:						
(1) Down	20/200					
(2) Lateral	20/70					
(3) Up	20/40					
Note: (1) Correct diagnosis reflecting disease or injury should be cited						

Also entitled to special monthly compensation.
 Add 10 percent if artificial eye cannot be worn; also entitled to special monthly compensation.

RATINGS FOR IMPAIRMENT OF MUSCLE FUNCTION—Continued

[6090 Diplopia (double vision)]

Degree of diplopia	Equiva lent visual acuity
Note: (2) The above ratings will be applied to only one eye. Ratings will not be applied for both diplopia and decreased visual acuity or field of vision in the same eye. When diplopia is present and there is also ratable impairment of visual acuity or field of vision of both eyes the above diplopia ratings will be applied to the poorer eye while the better eye is rated according to the best corrected visual acuity or visual field  Note: (3) When the diplopia field extends beyond more than one quadrant or more than one range of degrees, the evaluation for diplopia will be based on the quadrant and degree range that provide the highest evaluation  Note: (4) When diplopia exists in two individual and separate areas of the same eye, the equivalent visual acuity will be taken one step worse, but no worse than 5/200  6091 Symblepharon  Rate as limited muscle function, diagnostic code 6090  6092 Diplopia, due to limited muscle function	

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 40 FR 42537, Sept. 15, 1975; 41 FR 11297, Mar. 18, 1976; 43 FR 45354, Oct. 2, 1978; 51 FR 6411, Feb. 24, 1986; 53 FR 30264, Aug. 11, 1988; 53 FR 50955, Dec. 19, 1988; 57 FR 24364, June 9, 1992]

IMPAIRMENT OF AUDITORY ACUITY

## § 4.85 Evaluation of hearing impairment.

(a) An examination for hearing impairment for VA purposes must be conducted by a state-licensed audiologist and must include a controlled speech discrimination test (Maryland CNC) and a puretone audiometry test. Examinations will be conducted without the use of hearing aids.

(b) Table VI, "Numeric Designation

(b) Table VI, "Numeric Designation of Hearing Impairment Based on Puretone Threshold Average and Speech Discrimination," is used to determine a Roman numeral designation (I through XI) for hearing impairment based on a combination of the percent of speech discrimination (horizontal

rows) and the puretone threshold average (vertical columns). The Roman numeral designation is located at the point where the percentage of speech discrimination and puretone threshold average intersect.

(c) Table VIa, "Numeric Designation of Hearing Impairment Based Only on Puretone Threshold Average," is used to determine a Roman numeral designation (I through XI) for hearing impairment based only on the puretone threshold average. Table VIa will be used when the examiner certifies that use of the speech discrimination test is not appropriate because of language difficulties, inconsistent speech discrimination scores, etc., or when indicated under the provisions of §4.86.

(d) "Puretone threshold average," as used in Tables VI and VIa, is the sum of the puretone thresholds at 1000, 2000, 3000 and 4000 Hertz, divided by four. This average is used in all cases (including those in §4.86) to determine the Roman numeral designation for hearing impairment from Table VI or VIa.

- (e) Table VII, "Percentage Evaluations for Hearing Impairment," is used to determine the percentage evaluation by combining the Roman numeral designations for hearing impairment of each ear. The horizontal rows represent the ear having the better hearing and the vertical columns the ear having the poorer hearing. The percentage evaluation is located at the point where the row and column intersect.
- (f) If impaired hearing is service-connected in only one ear, in order to determine the percentage evaluation from Table VII, the non-service-connected ear will be assigned a Roman Numeral designation for hearing impairment of I, subject to the provisions of §3.383 of this chapter.
- (g) When evaluating any claim for impaired hearing, refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation due either to deafness, or to deafness in combination with other specified disabilities.
  - (h) Numeric tables VI, VIA\*, and VII.

#### TABLE VI

## NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ON PURETONE THRESHOLD AVERAGE AND SPEECH DISCRIMINATION

#### **Puretone Threshold Average**

% of discrim- ination	0-41	42-49	50-57	58-65	66-73	74-81	82-89	90-97	98+
92-100	I	I	I	II	II	II	III	III	IV
84-90	II	II	II	III	III	III	IV	IV	IV
76-82	III	III	ÏV	IV	IV	V	V	V	V
68-74	IV	IV	V	V	VI	VI	VII	VII	VII
60-66	V	v	VI	VI	VII	VII	VIII	VIII	VIII
52-58	VI	VI	VII	VII	VIII	VIII	VIII	VIII	IX
44-50	VII	VII	VIII	VIII	VIII	IX	IX	IX	X
36-42	VIII	VIII	VIII	IX	IX	IX	X	X	X
0-34	IX	X	XI	XI	XI	XI	XI	XI	XI

#### TABLE VIA\*

## NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ONLY ON PURETONE THRESHOLD AVERAGE

#### **Puretone Threshold Average**

0-41	42-48	49-55	56-62	63-69	70-76	77-83	84-90	91-97	98-104	105+
I	II	III	IV	V	VI	VII	VIII	IX	X	XI

<sup>\*</sup> This table is for use only as specified in §§ 4.85 and 4.86.

TABLE VII

## PERCENTAGE EVALUATION FOR HEARING IMPAIRMENT (DIAGNOSTIC CODE 6100)

#### **Poorer Ear**

XI	100*										
X	90	80									
IX	80	70	60					-			
VIII	70	60	50	50							
VII	60	60	50	40	40						
VI	50	50	40	40	30	30	-				
v	40	40	40	30	30	20	20				
IV	30	30	30	20	20	20	10	10			
III	20	20	20	20	20	10	10	10	0		
II	10	10	10	10	10	10	10	0 .	0	0	
I	10	10	0	0	0	0	O	0	0	0	0
L	XI	X	IX	VIII	VII	VI	V	IV	Ш	II	I

<sup>\*</sup> Review for entitlement to special monthly compensation under §3.350 of this chapter.

[64 FR 25206, May 11, 1999]

## § 4.86 Exceptional patterns of hearing impairment.

(a) When the puretone threshold at each of the four specified frequencies (1000, 2000, 3000, and 4000 Hertz) is 55 decibels or more, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher nu-

meral. Each ear will be evaluated separately.

(b) When the puretone threshold is 30 decibels or less at 1000 Hertz, and 70 decibels or more at 2000 Hertz, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. That numeral will then be elevated to the next higher

only):

Rate on impairment of function.

6210 Chronic otitis externa:

Rating

10

30

100

60 30

10

Roman numeral. Each ear will be evaluated separately.

(Authority: 38 U.S.C. 1155) [64 FR 25209, May 11, 1999]

#### §4.87 Schedule of ratings—ear.

#### DISEASES OF THE EAR

DISEASES OF THE LAR	
6200 Chronic suppurative otitis media, mastoiditis, or cholesteatoma (or any combination):  During suppuration, or with aural polyps  NOTE: Evaluate hearing impairment, and complications such as labyrinthitis, tinnitus, facial	
nerve paralysis, or bone loss of skull, separately.	
6201 Chronic nonsuppurative otitis media with effusion (serous otitis media):	
Rate hearing impairment 6202 Otosclerosis: Rate hearing impairment	
6204 Peripheral vestibular disorders:	
Dizziness and occasional staggering  Occasional dizziness	
NOTE: Objective findings supporting the diagnosis of vestibular disequilibrium are required before a compensable evaluation can be assigned under this code. Hearing impairment or suppuration shall be separately rated and	
combined.	
6205 Meniere's syndrome (endolymphatic hydrops): Hearing impairment with attacks of vertigo and	
cerebellar gait occurring more than once weekly, with or without tinnitus	
Hearing impairment with attacks of vertigo and cerebellar gait occurring from one to four	
times a month, with or without tinnitus	
Hearing impairment with vertigo less than once a month, with or without tinnitus	
Note: Evaluate Meniere's syndrome either under	
these criteria or by separately evaluating vertigo (as a peripheral vestibular disorder),	
hearing impairment, and tinnitus, whichever	
method results in a higher overall evaluation.	
But do not combine an evaluation for hearing impairment, tinnitus, or vertigo with an evalua-	
tion under diagnostic code 6205. 6207 Loss of auricle:	
Complete loss of both	
Complete loss of one	
of the substance	
skin only)	
NOTE: A rating of 100 percent shall continue be- yond the cessation of any surgical, radiation treatment, antineoplastic chemotherapy or	
other therapeutic procedure. Six months after	
discontinuance of such treatment, the appro- priate disability rating shall be determined by	
mandatory VA examination. Any change in	
evaluation based on that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on re-	
siduals. 6209 Benign neoplasms of the ear (other than skin	

38 CFR Ch. I (7-1-07 Edition)

	Rat- ing
Swelling, dry and scaly or serous discharge, and itching requiring frequent and prolonged treatment	10 0 10

(Authority: 38 U.S.C. 1155)

[64 FR 25210, May 11, 1999, as amended at 68 FR 25823, May 14, 2003]

#### §4.87a Schedule ratings-other sense organs.

	Rat- ing
6275 Sense of smell, complete loss	10 10

(Authority: 38 U.S.C. 1155) [64 FR 25210, May 11, 1999]

INFECTIOUS DISEASES, IMMUNE DIS-ORDERS AND NUTRITIONAL Defi-

§4.88 [Reserved]

#### § 4.88a Chronic fatigue syndrome.

(a) For VA purposes, the diagnosis of chronic fatigue syndrome requires:

(1) new onset of debilitating fatigue severe enough to reduce daily activity to less than 50 percent of the usual level for at least six months; and

(2) the exclusion, by history, physical examination, and laboratory tests, of all other clinical conditions that may produce similar symptoms; and

- (3) six or more of the following:
- (i) acute onset of the condition,
- (ii) low grade fever,
- (iii) nonexudative pharyngitis,

418

## DISEASES OF THE EAR—Continued

CIENCIES

§4.88b

- (iv) palpable or tender cervical or axillary lymph nodes,
- (v) generalized muscle aches or weakness,
- (vi) fatigue lasting 24 hours or longer after exercise,
- (vii) headaches (of a type, severity, or pattern that is different from headaches in the pre-morbid state),

  - (viii) migratory joint pains, (ix) neuropsychologic symptoms,
  - (x) sleep disturbance.
  - (b) [Reserved]

[59 FR 60902, Nov. 29, 1994]

#### §4.88b Schedule of ratings—infectious diseases, immune disorders and nutritional deficiencies.

	V-0-1-4-1 - 4-0-1-0-1-0-1-0-1-0-1-0-1-0-1-0-1-0-1-0-	
		Rating
6300	Cholera, Asiatic: As active disease, and for 3 months convalescence	100
6301	Visceral Leishmaniasis:  During treatment for active disease	100
	NOTE: A 100 percent evaluation shall continue beyond the cessation of treatment for active disease. Six discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory vition. Any change in evaluation based upon that or any subsequent examination shall be subject to the §3.105(e) of this chapter. Rate residuals such as liver damage or lymphadenopathy under the appropria	/A examina- provisions of
6302	Leprosy (Hansen's Disease): As active disease	100
	NOTE: A 100 percent evaluation shall continue beyond the date that an examining physician has determined has become inactive. Six months after the date of inactivity, the appropriate disability rating shall be demandatory VA examination. Any change in evaluation based upon that or any subsequent examination spect to the provisions of §3.105(e) of this chapter. Rate residuals such as skin lesions or periphera under the appropriate system.	etermined by shall be sub-
6304	Malaria: As active disease	100
	Note: The diagnosis of malaria depends on the identification of the malarial parasites in blood smears. It served in an endemic area and presents signs and symptoms compatible with malaria, the diagnosis m on clinical grounds alone. Relapses must be confirmed by the presence of malarial parasites in blood so Thereafter rate residuals such as liver or spleen damage under the appropriate system	ay be based
	Lymphatic Filariasis: As active disease Thereafter rate residuals such as epididymitis or lymphangitis under the appropriate system	100
6306	Bartonellosis: As active disease, and for 3 months convalescence	100
6307	Plague: As active disease	100
6308	Thereafter rate residuals such as lymphadenopathy under the appropriate system Relapsing Fever:	400
	As active disease  Thereafter rate residuals such as liver or spleen damage or central nervous system involvement under the system	100 appropriate
6309	Rheumatic fever: As active disease	100
6310	Thereafter rate residuals such as heart damage under the appropriate system Syphilis, and other treponemal infections:	
	Rate the complications of nervous system, vascular system, eyes or ears. (See DC 7004, syphilitic heart 8013, cerebrospinal syphilis, DC 8014, meningovascular syphilis, DC 8015, tabes dorsalis, and DC 930 associated with central nervous system syphilis)	
6311	Tuberculosis, miliary: As active disease Inactive: See §§ 4.88c and 4.89.	100
6313	Avitaminosis:  Marked mental changes, moist dermatitis, inability to retain adequate nourishment, exhaustion, and	400
	cachexia  With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor	100 60
	With stomatitis, diarrhea, and symmetrical dermatitis	40
	With stomatitis, or achlorhydria, or diarrhea	20
6314	discomfort, weakness, inability to concentrate and irritability	10
	As active disease: With congestive heart failure, anasarca, or Wernicke-Korsakoff syndrome	100

		Rating
	With cardiomegaly, or; with peripheral neuropathy with footdrop or atrophy of thigh or calf muscles With peripheral neuropathy with absent knee or ankle jerks and loss of sensation, or; with symptoms such as weakness, fatigue, anorexia, dizziness, heaviness and stiffness of legs, headache or sleep	60
	disturbance Thereafter rate residuals under the appropriate body system.	30
6315	Marked mental changes, moist dermatitis, inability to retain adequate nourishment, exhaustion, and	
	cachexia	100
	With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor	60 40
	With stomatitis, or achlorhydria, or diarrhea	20
	Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discomfort, weakness, inability to concentrate and irritability	10
316	Brucellosis:	
	As active disease	100
317	Thereafter rate residuals such as liver or spleen damage or meningitis under the appropriate system Typhus, scrub:	
	As active disease, and for 3 months convalescence	100
318	Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system Melioidosis:	
	As active disease	100
319	Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system Lyme Disease:	400
	As active disease	100
320		
	As active disease	100
350	Thereafter rate residuals such as spleen or liver damage under the appropriate system Lupus erythematosus, systemic (disseminated):	
	Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health	100
	Exacerbations lasting a week or more, 2 or 3 times per year	60 10
	NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate sys	stem, or by
	evaluating DC 6350, whichever method results in a higher evaluation.	, , , ,
351		
	AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remis-	
	sions	100
	development of AIDS-related opportunistic infection or neoplasm	60
	rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis	30
	Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations	10
	Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count	0
	NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protoc credited medical institution.	ol at an ac-
	NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms m separately under appropriate codes if higher overall evaluation results, but not in combination with perce erwise assignable above.	
354	Chronic Fatigue Syndrome (CFS):	
	Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a of other signs and symptoms:	combination
	Which may occasionally preclude self-care	100
	Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or, which wax and wane, resulting in periods of incapacitation of at least six weeks total duration	
	per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks	60
	total duration per year	40
	weeks total duration per year	20
	duration per year, or; symptoms controlled by continuous medication	10

#### § 4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19, 1968.

	Rat- ing
For 1 year after date of inactivity, following active tu- berculosis	100
Thereafter: Rate residuals under the specific body	

Following the total rating for the 1 year period after date of inactivity, the schedular evaluation for residuals of nonpulmonary tuberculosis, i.e., ankylosis, surgical removal of a part, etc., will be assigned under the appropriate diagnostic code for the residual preceded by the diagnostic code for tuberculosis of the body part affected. For example, tuberculosis of the hip joint with residual ankylosis would be coded 5001–5250. Where there are existing residuals of pulmonary and nonpulmonary conditions, the evaluations for residual separate functional impairment may be combined.

Where there are existing pulmonary and nonpulmonary conditions, the total rating for the 1 year, after attainment of inactivity, may not be applied to both conditions during the same period. However, the total rating during the 1-year period for the pulmonary or for the nonpulmonary condition will be utilized, combined with evaluation for residuals of the condition not covered by the 1-year total evaluation, so as to allow any additional benefit provided during such period.

[34 FR 5062, Mar. 11, 1969. Redesignated at 59 FR 60902, Nov. 29, 1994]

#### § 4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

Public Law 90-493 repealed section 356 of title 38, United States Code which provided graduated ratings for inactive tuberculosis. The repealed section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90-493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For use in rating cases in which the protective provisions of Pub. L. 90-493 apply, the former evaluations are retained in this section.

	Rat- ing
For 2 years after date of inactivity, following active tuberculosis, which was clinically identified during	
service or subsequently	100
Thereafter, for 4 years, or in any event, to 6 years after date of inactivity	50
Thereafter, for 5 years, or to 11 years after date of inactivity	30
Thereafter, in the absence of a schedular compen-	
sable permanent residual	0

Following the total rating for the 2-year period after date of inactivity, the schedular evaluation for residuals of nonpulmonary tuberculosis, i.e., ankylosis, surgical removal of a part, etc., if in excess of 50 percent or 30 percent will be assigned under the appropriate diagnostic code for the specific residual preceded by the diagnostic code for tuberculosis of the body part affected. For example, tuberculosis of the hipjoint with residual ankylosis would be coded 5001–5250.

The graduated ratings for nonpulmonary tuberculosis will not be combined with residuals of nonpulmonary tuberculosis unless the graduated rating and the rating for residual disability cover separate functional losses, e.g., graduated ratings for tuberculosis of the kidney and residuals of tuberculosis of the spine. Where there are existing pulmonary and nonpulmonary conditions, the graduated evaluation for the pulmonary, or for the nonpulmonary, condition will be utilized, combined with evaluations for residuals of the condition not covered by the graduated evaluation utilized, so as to provide the higher evaluation over such period.

The ending dates of all graduated ratings of nonpulmonary tuberculosis will be controlled by the date of attainment of inactivity.

These ratings are applicable only to veterans with nonpulmonary tuberculosis active on or after October 10, 1949.

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 43 FR 45361, Oct. 2, 1978]

#### THE RESPIRATORY SYSTEM

# § 4.96 Special provisions regarding evaluation of respiratory conditions.

(a) Rating coexisting respiratory conditions. Ratings under diagnostic codes 6600 through 6817 and 6822 through 6847 will not be combined with each other. Where there is lung or pleural involvement, ratings under diagnostic codes 6819 and 6820 will not be combined with each other or with diagnostic codes 6600 through 6817 or 6822 through 6847. A single rating will be assigned under the diagnostic code which reflects the predominant disability with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation. However, in cases protected by the provisions of Pub. L. 90-493, the graduated ratings of 50 and 30 percent for inactive tuberculosis will not be elevated.

(b) Rating "protected" tuberculosis cases. Public Law 90-493 repealed section 356 of title 38, United States Code which had provided graduated ratings for inactive tuberculosis. The repealed

section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90–493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For application in rating cases in which the protective provisions of Pub. L. 90–493 apply the former evaluations pertaining to pulmonary tuberculosis are retained in §4.97.

- (c) Special monthly compensation. When evaluating any claim involving complete organic aphonia, refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, there are other conditions in this section which under certain circumstances also establish entitlement to special monthly compensation.
- (d) Special provisions for the application of evaluation criteria for diagnostic codes 6600, 6603, 6604, 6825–6833, and 6840–6845
- (1) Pulmonary function tests (PFT's) are required to evaluate these conditions except:
- (i) When the results of a maximum exercise capacity test are of record and are 20 ml/kg/min or less. If a maximum exercise capacity test is not of record, evaluate based on alternative criteria.
- (ii) When pulmonary hypertension (documented by an echocardiogram or cardiac catheterization), cor pulmonale, or right ventricular hypertrophy has been diagnosed.
- (iii) When there have been one or more episodes of acute respiratory failure.
- (iv) When outpatient oxygen therapy is required.

- (2) If the DLCO (SB) (Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method) test is not of record, evaluate based on alternative criteria as long as the examiner states why the test would not be useful or valid in a particular case.
- (3) When the PFT's are not consistent with clinical findings, evaluate based on the PFT's unless the examiner states why they are not a valid indication of respiratory functional impairment in a particular case.
- (4) Post-bronchodilator studies are required when PFT's are done for disability evaluation purposes except when the results of pre-bronchodilator pulmonary function tests are normal or when the examiner determines that post-bronchodilator studies should not be done and states why.
- (5) When evaluating based on PFT's, use post-bronchodilator results in applying the evaluation criteria in the rating schedule unless the post-bronchodilator results were poorer than the pre-bronchodilator results. In those cases, use the pre-bronchodilator values for rating purposes.
- (6) When there is a disparity between the results of different PFT's (FEV-1 (Forced Expiratory Volume in one second), FVC (Forced Vital Capacity), etc.), so that the level of evaluation would differ depending on which test result is used, use the test result that the examiner states most accurately reflects the level of disability.
- (7) If the FEV-1 and the FVC are both greater than 100 percent, do not assign a compensable evaluation based on a decreased FEV-1/FVC ratio.

(Authority: 38 U.S.C. 1155)

[34 FR 5062, Mar. 11, 1969, as amended at 61 FR 46727, Sept. 5, 1996; 71 FR 52459, Sept. 6, 2006]

#### §4.97 Schedule of ratings—respiratory system.

		Rating
	DISEASES OF THE NOSE AND THROAT	
6502	Septum, nasal, deviation of: Traumatic only, With 50-percent obstruction of the nasal passage on both sides or complete obstruction on one side	10

		Railing
	Exposing both nasal passages	30
Noto:	Loss of part of one ala, or other obvious disfigurement	10
	Sinusitis, pansinusitis, chronic.	
5511		
	Sinusitis, frontal, chronic.	
5513	Sinusitis, maxillary, chronic.	
514	Sinusitis, sphenoid, chronic.	
	General Rating Formula for Sinusitis (DC's 6510 through 6514):	
	Following radical surgery with chronic osteomyelitis, or; near constant sinusitis characterized by headaches, pain and tenderness of affected sinus, and purulent discharge or crusting after repeated surgeries	50
	Three or more incapacitating episodes per year of sinusitis requiring prolonged (lasting four to six weeks) antibiotic treatment, or; more than six non-incapacitating episodes per year of sinusitis	0.
	characterized by headaches, pain, and purulent discharge or crusting	30
	acterized by headaches, pain, and purulent discharge or crusting  Detected by X-ray only	10
	Note: An incapacitating episode of sinusitis means one that requires bed rest and treatment by a physician.	
ô515	Laryngitis, tuberculous, active or inactive.	
	Rate under §§ 4.88c or 4.89, whichever is appropriate.	
3516	Laryngitis, chronic:	
	Hoarseness, with thickening or nodules of cords, polyps, submucous infiltration, or pre-malignant changes on biopsy	30
	Hoarseness, with inflammation of cords or mucous membrane	10
6518	Laryngectomy, total.	1 100
8510	Rate the residuals of partial laryngectomy as laryngitis (DC 6516), aphonia (DC 6519), or stenosis of larynx (DC 6520).  Aphonia, complete organic:	
5515	Constant inability to communicate by speech	<sup>1</sup> 100
	Constant inability to speak above a whisper	60
	Note: Evaluate incomplete aphonia as laryngitis, chronic (DC 6516).	
6520	Larynx, stenosis of, including residuals of laryngeal trauma (unilateral or bilateral): Forced expiratory volume in one second (FEV–1) less than 40 percent of predicted value, with Flow-Volume	
	Loop compatible with upper airway obstruction, or; permanent tracheostomy  FEV-1 of 40- to 55-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction  FEV-1 of 56- to 70-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction  FEV-1 of 71- to 80-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction	100 60 30 10
	Note: Or evaluate as aphonia (DC 6519).	
3521	Pharynx, injuries to:	
3321	Stricture or obstruction of pharynx or nasopharynx, or; absence of soft palate secondary to trauma, chemical burn, or granulomatous disease, or; paralysis of soft palate with swallowing difficulty (nasal regurgitation)	
	and speech impairment	50
6522	Allergic or vasomotor rhinitis:	0.0
	With polyps Without polyps, but with greater than 50-percent obstruction of nasal passage on both sides or complete ob-	30
5523	struction on one side  Bacterial thinitis:	10
	Rhinoscleroma	50
5524	With permanent hypertrophy of turbinates and with greater than 50-percent obstruction of nasal passage on both sides or complete obstruction on one side	10
0024	Wegener's granulomatosis, lethal midline granuloma	100
	Other types of granulomatous infection	20
	DISEASES OF THE TRACHEA AND BRONCHI	
3600	Bronchitis, chronic:	
	FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy	10
	FEV-1 of 40- to 55-percent predicted, or, FEV-1/FVC of 40 to 55 percent, or, DLCO (SB) of 40- to 55-per-	
	cent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	60
	predicted FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted	30
6601	Bronchiectasis:	

n incapacitating episodes of infection of at least six weeks total duration per year n incapacitating episodes of infection of four to six weeks total duration per year, or; near constant findings f cough with purulent sputum associated with anorexia, weight loss, and frank hemoptysis and requiring
ntibiotic usage almost continuously
rmittent productive cough with acute infection requiring a course of antibiotics at least twice a year
e: An incapacitating episode is one that requires bedrest and treatment by a physician.
na, bronchial:
/-1 less than 40-percent predicted, or; FEV-1/FVC less than 40 percent, or; more than one attack per eek with episodes of respiratory failure, or; requires daily use of systemic (oral or parenteral) high dose orticosteroids or immuno-suppressive medications
/–1 of 40- to 55-percent predicted, or; FEV–1/FVC of 40 to 55 percent, or; at least monthly visits to a phycian for required care of exacerbations, or; intermittent (at least three per year) courses of systemic (oral representation) corticosteroids
/-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; daily inhalational or oral bron- nodilator therapy, or; inhalational anti-inflammatory medication
/-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; intermittent inhalational or oral ronchodilator therapy
<ul> <li>In the absence of clinical findings of asthma at time of examination, a verified history of asthmatic at- icks must be of record.</li> </ul>
pysema, pulmonary:  /-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to orced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon lonoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise apacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor ulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo r cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy.  /-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-perent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)
(-1 of 71 to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted
nic obstructive pulmonary disease:  '-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to orced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon lonoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise apacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor ulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo
r cardiac catheterization), or, episode(s) of acute respiratory failure, or, requires outpatient oxygen therapy.  /-1 of 40- to 55-percent predicted, or, FEV-1/FVC of 40 to 55 percent, or, DLCO (SB) of 40- to 55-per- ent predicted, or, maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) /-1 of 56- to 70-percent predicted, or, FEV-1/FVC of 56 to 70 percent, or, DLCO (SB) 56- to 65-percent redicted /-1 of 71- to 80-percent predicted, or, FEV-1/FVC of 71 to 80 percent, or, DLCO (SB) 66- to 80-percent redicted
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-per- ent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) -/-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent redicted
/–1 of 40- to 55-percent predicted, or; FEV–1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) ————————————————————————————————————
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)/-/-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent redicted
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)/-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent redicted
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) /-/1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent redicted /-/1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 t
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)/-/-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent redicted
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) /-/1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent redicted /-/1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted /-/2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent redicted //2 of 71 to 80 percent,
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)
/-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-perent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) /-/-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent predicted /-/
/—1 of 40- to 55-percent predicted, or; FEV—1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit) ————————————————————————————————————

		Rating
treated bero to the treated bero to the treated bero to the treated bero the treated bero treated by the treated bero treated bero treated by the treated by t	(1): The 100-percent rating under codes 6701 through 6724 is not subject to a requirement of precedent hospital tment. It will be reduced to 50 percent for failure to submit to examination or to follow prescribed treatment upon but to that effect from the medical authorities. When a veteran is placed on the 100-percent rating for inactive tublosis, the medical authorities will be appropriately notified of the fact, and of the necessity, as given in footnote 38 U.S.C. 1156 (and formerly in 38 U.S.C. 356, which has been repealed by Public Law 90–493), to notify the errans Service Center in the event of failure to submit to examination or to follow treatment.  (2): The graduated 50-percent and 30-percent ratings and the permanent 30 percent and 20 percent ratings for tive pulmonary tuberculosis are not to be combined with ratings for other respiratory disabilities. Following acoplasty the rating will be for removal of ribs combined with the rating for collapsed lung. Resection of the ribs dent to thoracoplasty will be rated as removal.	
	Ratings for Pulmonary Tuberculosis Initially Evaluated After August 19, 1968	
6730	Tuberculosis, pulmonary, chronic, active	100
6731	Tuberculosis, pulmonary, chronic, inactive:  Depending on the specific findings, rate residuals as interstitial lung disease, restrictive lung disease, or, when obstructive lung disease is the major residual, as chronic bronchitis (DC 6600). Rate thoracoplasty as removal of ribs under DC 5297.  Note: A mandatory examination will be requested immediately following notification that active tuberculosis evaluated under DC 6730 has become inactive. Any change in evaluation will be carried out under the pro-	
6732	visions of §3.105(e).  Pleurisy, tuberculous, active or inactive:  Rate under §§ 4.88c or 4.89, whichever is appropriate.	
	NONTUBERCULOUS DISEASES	
6817	Pulmonary Vascular Disease:  Primary pulmonary hypertension, or; chronic pulmonary thromboembolism with evidence of pulmonary hypertension, right ventricular hypertrophy, or cor pulmonale, or; pulmonary hypertension secondary to other obstructive disease of pulmonary arteries or veins with evidence of right ventricular hypertrophy or cor pulmonale	100
	without evidence of pulmonary hypertension or right ventricular dysfunction  Symptomatic, following resolution of acute pulmonary embolism  Asymptomatic, following resolution of pulmonary thromboembolism  Note: Evaluate other residuals following pulmonary embolism under the most appropriate diagnostic code, such as chronic bronchitis (DC 6600) or chronic pleural effusion or fibrosis (DC 6844), but do not combine	30
	that evaluation with any of the above evaluations.  6819 Neoplasms, malignant, any specified part of respiratory system exclusive of skin growths	100
6820	Neoplasms, benign, any specified part of respiratory system. Evaluate using an appropriate respiratory analogy.	
	Bacterial Infections of the Lung	
	Actinomycosis.  Nocardiosis.  Chronic lung abscess.  General Rating Formula for Bacterial Infections of the Lung (diagnostic codes 6822 through 6824):  Active infection with systemic symptoms such as fever, night sweats, weight loss, or hemoptysis  Depending on the specific findings, rate residuals as interstitial lung disease, restrictive lung disease, or, when obstructive lung disease is the major residual, as chronic bronchitis (DC 6600).	100
	Interstitial Lung Disease	
6825 6826 6827 6828	Diffuse interstitial fibrosis (interstitial pneumonitis, fibrosing alveolitis). Desquamative interstitial pneumonitis. Pulmonary alveolar proteinosis. Eosinophilic granuloma of lung.	

### 38 CFR Ch. I (7-1-07 Edition)

		Rating
	Radiation-induced pulmonary pneumonitis and fibrosis.	100 60 30 10
	Mycotic Lung Disease	
	Cryptococcosis. Aspergillosis.	100 50 30 0
-	Restrictive Lung Disease	
6840 6841 6842 6843 6844 6845	Spinal cord injurý with respiratory insufficiency. Kyphoscoliosis, pectus excavatum, pectus carinatum. Traumatic chest wall defect, pneumothorax, hernia, etc. Post-surgical residual (lobectomy, pneumonectomy, etc.).	100 60 30 10

		Rating
	Cor pulmonale, or; cardiac involvement with congestive heart failure, or; progressive pulmonary disease with fever, night sweats, and weight loss despite treatment	100
	Pulmonary involvement requiring systemic high dose (therapeutic) corticosteroids for control	60
	Pulmonary involvement with persistent symptoms requiring chronic low dose (maintenance) or intermittent corticosteroids	30
	Chronic hilar adenopathy or stable lung infiltrates without symptoms or physiologic impairment	0
	Or rate active disease or residuals as chronic bronchitis (DC 6600) and extra-pulmonary involvement under specific body system involved.	
6847	Sleep Apnea Syndromes (Obstructive, Central, Mixed):	
	Chronic respiratory failure with carbon dioxide retention or cor pulmonale, or; requires tracheostomy	100
	Requires use of breathing assistance device such as continuous airway pressure (CPAP) machine	50
	Persistent day-time hypersomnolence	30
	Asymptomatic but with documented sleep disorder breathing	0

Note

sec

con Note

an

syn

mat

exa sno

7000 hea

[61 FR 46728, Sept. 5, 1996, as amended at 71 FR 28586, May 17, 2006]

#### THE CARDIOVASCULAR SYSTEM

# § 4.100 Application of the evaluation criteria for diagnostic codes 7000–7007, 7011, and 7015–7020.

- (a) Whether or not cardiac hypertrophy or dilatation (documented by electrocardiogram, echocardiogram, or X-ray) is present and whether or not there is a need for continuous medication must be ascertained in all cases.
- (b) Even if the requirement for a 10% (based on the need for continuous medication) or 30% (based on the presence of cardiac hypertrophy or dilatation) evaluation is met, METs testing is required in all cases except:
- (1) When there is a medical contraindication.
- (2) When the left ventricular ejection fraction has been measured and is 50% or less
- (3) When chronic congestive heart failure is present or there has been more than one episode of congestive heart failure within the past year.
- (4) When a 100% evaluation can be assigned on another basis.
- (c) If left ventricular ejection fraction (LVEF) testing is not of record, evaluate based on the alternative criteria unless the examiner states that the LVEF test is needed in a particular case because the available medical information does not sufficiently reflect the severity of the veteran's cardiovascular disability.

[71 FR 52460, Sept. 6, 2006]

#### §§ 4.101-4.103 [Reserved]

#### §4.104 Schedule of ratings—cardiovascular system.

#### DISEASES OF THE HEART

	Rat- ing
(1): Evaluate cor pulmonale, which is a form of condary heart disease, as part of the pulmonary hedition that causes it.  (2): One MET (metabolic equivalent) is the eny cost of standing quietly at rest and represents oxygen uptake of 3.5 milliliters per kilogram of dy weight per minute. When the level of METs which dyspnea, fatigue, angina, dizziness, or icope develops is required for evaluation, and a orationy determination of METs by exercise test-cannot be done for medical reasons, an estition by a medical examiner of the level of activ-(expressed in METs and supported by specific amples, such as slow stair climbing or shoveling by that results in dyspnea, fatigue, angina, dizess, or syncope may be used.  Valvular heart disease (including rheumatic	
art disease):  During active infection with valvular heart damage and for three months following cessation of therapy for the active infection  Thereafter, with valvular heart disease (documented by findings on physical examination and either echocardiogram, Doppler echocardiogram, or cardiac catheterization) resulting in:	100
Chronic congestive heart failure, or; work-load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100
tion of 30 to 50 percent	60

<sup>&</sup>lt;sup>1</sup> Review for entitlement to special monthly compensation under § 3.350 of this chapter.

### 38 CFR Ch. I (7-1-07 Edition)

#### DISEASES OF THE HEART—Continued

		Rat- ing		Rat- ing
	Workload of greater than 5 METs but not greater than 7 METs results in dyspnea,		Chronic congestive heart failure, or; work-	.5
	fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata-		load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection	
,	tion on electro-cardiogram, echocardio- gram, or X-ray	30	fraction of less than 30 percent	100
•	greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;		of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue,	
	continuous medication requiredndocarditis:  For three months following cessation of	10	angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60
	therapy for active infection with cardiac involvement	100	Workload of greater than 5 METs but not greater than 7 METs results in dyspnea,	00
٦	Thereafter, with endocarditis (documented by findings on physical examination and either echocardiogram, Doppler echo-		fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata- tion on electro-cardiogram, echocardio-	
	cardiogram, or cardiac catheterization) resulting in:		gram, or X-ray Workload of greater than 7 METs but not	30
(	Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or;		greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10
	left ventricular dysfunction with an ejection fraction of less than 30 percent	100	7004 Syphilitic heart disease: Chronic congestive heart failure, or; work-	
ľ	More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater		load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection	
	than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven-		fraction of less than 30 percent  More than one episode of acute congestive	100
١	tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60	heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatique,	
	greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;		angina, dizziness, or syncope, or, left ven- tricular dysfunction with an ejection frac-	00
	evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio- gram, or X-ray	30	tion of 30 to 50 percent	60
١	Norkload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;		fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio-	
	continuous medication required	10	gram, or X-ray Workload of greater than 7 METs but not	30
ŀ	For three months following cessation of therapy for active infection with cardiac involvement	100	greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10
	Thereafter, with documented pericarditis resulting in:		NOTE: Evaluate syphilitic aortic aneurysms under DC 7110 (aortic aneurysm).	
(	Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or;		7005 Arteriosclerotic heart disease (Coronary artery disease):	
,	left ventricular dysfunction with an ejection fraction of less than 30 percent	100	With documented coronary artery disease resulting in: Chronic congestive heart failure, or; work-	
•	heart failure in the past year, or; workload of greater than 3 METs but not greater		load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or;	
	than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection frac-		left ventricular dysfunction with an ejection fraction of less than 30 percent	100
١	tion of 30 to 50 percent	60	heart failure in the past year, or; workload of greater than 3 METs but not greater	
	greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata-		than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac-	
,	tion on electro-cardiogram, echocardio- gram, or X-ray	30	tion of 30 to 50 percent	60
\	greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;		greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata-	
7003 Pe	continuous medication required	10	tion on electrocardiogram, echocardiogram, or X-ray	30

### § 4.104

	Rat- ing		Rat- ing
Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10	Include as part of the overall evaluation for hyperthyroidism under DC 7900. However, when atrial fibrillation is present, hyperthyroidism may be evaluated either under DC 7900 or under DC 7010 (supraventricular arrhythmia), whichever results in a higher evaluation.  7010 Supraventricular arrhythmias: Paroxysmal atrial fibrillation or other supra-	
7006 Myocardial infarction:  During and for three months following myocardial infarction, documented by laboratory tests	100	ventricular tachycardia, with more than four episodes per year documented by ECG or Holter monitor	30
With history of documented myocardial infarction, resulting in: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100	ECG or Holter monitor.  7011 Ventricular arrhythmias (sustained): For indefinite period from date of hospital admission for initial evaluation and medical therapy for a sustained ventricular arrhythmia, or, for indefinite period from date of hospital admission for ventricular aneurysmectomy, or, with an automatic implantable Cardioverter-Defibrillator (AICD) in place	100
than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60	Chronic congestive heart failure, or; work-load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100
tion on electrocardiogram, echocardiogram, or X-ray	30	than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60
Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100	fatigue, angina, dizziness, or syncope, or, evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray	30
heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60	continuous medication required	10
fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio- gram, or X-ray	30	mined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7015 Atrioventricular block:	
fatigue, angina, dizziness, or syncope, or; continuous medication required	10	Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100

### 38 CFR Ch. I (7-1-07 Edition)

#### DISEASES OF THE HEART—Continued

	Rat- ing		Rat- ing
More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60	More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60
continuous medication or a pacemaker required	10	continuous medication required	10
ventricular block associated with a supraventricular arrhythmia or pathological bradycardia should be submitted to the Director, Compensation and Pension Service. Simple delayed P-R conduction time, in the absence of other evidence of cardiac disease, is not a disability.		for implantation or reimplantation	100
<ul> <li>7016 Heart valve replacement (prosthesis):</li> <li>For indefinite period following date of hospital admission for valve replacement</li> <li>Thereafter:</li> <li>Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent</li> <li>More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent</li> <li>Workload of greater than 5 METs but not greater than 7 METs results in dyspnea.</li> </ul>	100	Note: Evaluate implantable Cardioverter-Defibrillators (AICD's) under DC 7011.  7019 Cardiac transplantation: For an indefinite period from date of hospital admission for cardiac transplantation Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fractional results and provided the second secon	100
fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray	30 10	tion of 30 to 50 percent	100
Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent		than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60

### §4.104

Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizzines, or syncope, or evidence of cardiac hypertrophy or distation on electrocardiogram, echocardiogram, chocardiogram, chocardiogr		Rat- ing		Rat- ing
greater than 7 METs results in dyspnea, fatigue, angina, dizziense, or synope, or evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray.  Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizzienses, or synope, or continuous medication required.  Diseases of the Arteries and Weins 7101 Hypertensive vascular disease (hypertension and isolated systolic hypertension):  Diastolic pressure predominantly 130 or more more measure predominantly 200 or more more, or; systolic pressure predominantly 200 or more, or; systolic pressure predominantly 100 or more, or; minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or or systolic pressure predominantly 100 or more who requires continuous medication for control work or expectation or isolated systolic hypertension means that the diastolic blood pressure is predominantly 100 or or systolic byte or systolic byte or systolic byte or systolic byte pressure predominantly 100 or more who requires continuous medication for control with the diastolic blood pressure is predominantly 100 or or systolic hypertension means that the diastolic blood pressure is predominantly 100 or or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 100 or or greater with a diastolic blood pressure of less than 90mm.  Nort: (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  Nort: (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  Nort: (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic hypertensive heart disease.  Nort: (1): Hypertension or isolated systolic hypertension means that t		ing	Claudiantian on well-trailed the CT	ilig
tion on electrocardiogram, echocardiogram, or X-ray Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or, continuous medication required Diseases of the Arteries and Veins Tol1 + Hypertension vascular disease (hypertension and isolated systolic hypertension): Diastolic pressure predominantly 130 or more. Diastolic pressure predominantly 110 or more, or, systolic pressure predominantly 200 or more, or, systolic pressure predominantly 100 or more who requires continuous medication for control or more in the distolic pressure predominantly 100 or more who requires continuous medication for control or more in the distolic pressure predominantly 100 or more who requires continuous medication for control or more in the distolic blood pressure in predominantly 100 or more who requires continuous medication for control or more in the distolic blood pressure in predominantly 100 or more who requires continuous medication for control or more in the distolic blood pressure of less than 90mm. NOTE (2): Evaluate hypertension due to activity in the distolic properties of the distolic properties of the distolic blood pressure of less than 90mm. NOTE (2): Evaluate hypertension due to activity in the distolic properties of the distolic propersure is predominantly 100 mm or greater with a diastolic blood pressure of less than 90mm. NOTE (2): Evaluate hypertension due to activity the solution of the distolic properties of the distolic properties of the distolic propersure is predominantly 10	greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;		on a level grade at 2 miles per hour, and; persistent coldness of the extremity, one	
Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizenses, or synope, or, continuous medication required				60
Workload of greater than 10 METs but not greater than 10 METs results in dyspena, fatigue, angina, dizziness, or syncope, or, continuous medication required.  Diseases of the Arteries and Veins  7101 Hypertensiev execulate disease (hypertension and isolated systolic hypertension): Diastolic pressure predominantly 130 or more.  Diastolic pressure predominantly 110 or more, or, systolic pressure predominantly 200 or more.  Diastolic pressure predominantly 110 or more, or, systolic pressure predominantly 200 or more.  Diastolic pressure predominantly 100 or more, or, systolic pressure predominantly 200 or more in more or, systolic pressure predominantly 200 or more in more, or, systolic pressure predominantly 200 or more in more in the distolic pressure predominantly 200 or more in more in the distolic pressure predominantly 100 or more whore requires continuous medication for control or more in solated systolic hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure or less than 90mm.  Nore (2): Evaluate hypertension due to actic including systolic hypertension on the activation of the condition causing it rather than by a separate evaluation.  Nore (3): Evaluate hypertension separately from hypertension wheat of sease and other types of heart disease.  Nore (3): Evaluate previouslism of surgical correction (including any type of graft insertion).  Procluding exertion  Evaluate residuals of surgical correction according to organ systems affected.  Nore: A rating of 100 percent shall be assigned as of the date of admission for surgical correction including any type of graft insertion, and the provisions of \$3.105(e) of this chapter.  100 Adrica analysis and the subject to the provisions of \$3.105(e) of the subject to the provisions of \$4.05(e) of the subject to the provisions of \$4.05(e) of the subject to the provisions of \$4.05(e) of the subject		30		60
greater than 10 METs results in dyspnea, fatigue, angina, dzizeness, or syncope, or, continuous medication required of the Arteries and Veins  7101 Hypertensive vascular disease (hypertension and isolated systolic hypertension): Diastolic pressure predominantly 130 or more. Diastolic pressure predominantly 110 or more, or, systolic pressure predominantly 200 or more, or, systolic pressure predominantly 200 or more, or, systolic pressure predominantly 110 or more, or, systolic pressure predominantly 100 or more, or, systolic pressure predominantly 160 or more, or, systolic pressure systolic blood pressure states of the section, the term hypertension means that the diastolic blood pressure is predominantly 160 or or greater with a diastolic blood pressure of less than 90mm.  Notte (1): Phypertension or isolated systolic hypertension means that the diastolic blood pressure is predominantly 160m, or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160m, or greater with a diastolic blood pressure of less than 90mm.  Notte (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as pard to the condition causigit it rather than by a separate evaluation.  Notte (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as pard to the condition causigit it rather than by a separate evaluation.  Notte (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic hypertension means that the diastolic blood pressure is predominantly 160m, or greater with a diastolic blood pressure is predominantly 160m, or greater with a diastolic blood pressure is predominantly 160m, or greater with a diastolic blood pressu		30		
fatigue, angina, dizziness, or syncope, or continuous medication required  Disasses of the Arteries and Veins  7101 Hypertersieve vascular disease (hypertension and isolated systolic hypertension):  Diastolic pressure predominantly 130 or more  Diastolic pressure predominantly 110 or more, or, systolic pressure predominantly 200 or more  Diastolic pressure predominantly 100 or more, or, systolic pressure predominantly 100 or more or, or minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control or an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control or an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control was a formation of the condition or control or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 100 ormore organization was a formation or greater with a diastolic blood pressure of less than 90mm.  Notre (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  Notre (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or, if symptomatic, or, for indefinite period from date of hospital admission for surgical correction. Six months following discharge, the appropriate di				
Disasses of the Arteries and Veins 7101 Hypertensive vascular diseases (hypertension): Diastolic pressure predominantly 130 or more Diastolic pressure predominantly 120 or more or more, or systolic pressure predominantly 100 or more, or more, or systolic pressure predominantly 100 or more, or more, or systolic pressure predominantly 100 or more, or more, or systolic pressure predominantly 100 or more who are individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control NOTE (1): Hypertension or isolated systolic hypertension means that the diastolic blood pressure is predominantly 100 or more who requires continuous medication for control NOTE (1): Evaluate hypertension for surgical correction for control was a feet of this section, the term hypertension means that the systolic blood pressure is predominantly 60mm, or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm, or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm, or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensior heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or, if symptomatic, or, or for indefinite period from date of hospital admission for surgical correction. Six months following discharge, the appropriate discharge the present of the date of admission for surgical correction. S	fatigue, angina, dizziness, or syncope, or:			
Disastolic pressure predominantly 120 or more. Diastolic pressure predominantly 120 or more, or, systolic pressure predominantly 120 or more, or, systolic pressure predominantly 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more times on at least three different days. For purposes of this section, the term hypertension means that the distolic blood pressure is predominantly 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more who requires continuous medication for control 100 or more who requires continuous medi	continuous medication required	10		40
7.101 Hypertensive vascular disease (hypertension and isolated systolic pressure predominantly 130 or more			Claudication on walking more than 100	
and isolated systolic phypertension):  Diastolic pressure predominantly 120 or more  Diastolic pressure predominantly 110 or more, or, systolic pressure predominantly 200 or more individual with a history of diastolic pressure predominantly 160 or more, or, systolic pressure predominantly 160 or more, or, minimum evaluation for an individual with a history of diastolic pressure predominantly 160 or more who requires continuous medication for control or more on a least three different days. For purposes of this section, the term hypertension means that the distolic bodo pressure is predominantly 90mm, or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm or greater with a diastolic blood pressure of less than 90mm.  Note (2): Evaluate hypertension due to aortic insufficiency or hyperthyriodism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  Note (3): Evaluate hypertension due to aortic insufficiency or hyperthyriodism, which is usually the isolated systolic bype, as part of the condition causing it rather than by a separate evaluation.  Note (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  If the centimeters or larger in diameter, or, if symptomatic, or, for indefinite period from date of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation.  Note (3): Evaluate hypertension dependent publication of the date of hospital admission for surgical correction. Six months of the date of hospital admission for surgical correction. Six months of the date of admission for surgical correction according to organ systems aff				
Diastolic pressure predominantly 120 or more				20
Diastolic pressure predominantly 120 or more.  Diastolic pressure predominantly 110 or more, or; systolic pressure predominantly 200 or more more, or; systolic pressure predominantly 200 or more, or; systolic pressure predominantly 100 or more, or; systolic pressure predominantly 100 or more, or, systolic pressure predominantly 100 or more with the control of an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control or an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control or an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control or surgical correction must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 160 mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160 mm. or greater, and isolated systolic blood pressure is predominantly 160 mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to acrtic insufficiency or hyperthyroidism, which is usually the isolated systolic type. as part of the condition causing it rather than by a separate evaluation.  Precluding exertion in diameter, or; if symptomatic, or for indefinite period from date of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subjec	Diastolic pressure predominantly 130 or			
Diastolic pressure predominantly 110 or more. Or, systolic pressure predominantly 110 or more, or, systolic pressure predominantly 100 or more who requires continuous medication for control (and individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control (and individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control (and individual with a history of diastolic pressure predominantly 100 or more who requires on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less that be systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less that the diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm or isolated systolic hypertension means that the distolic blood pressure is predominantly 160mm or isolated systolic hypertension means that the distolated systolic hypertension means that the distolated hypertension where the provisions of §3.105(e) of this chapter.  Note (2): Evaluate hypertension due to active hypertension means that the distolated hypertension means that the systolic blood pressure is a frected, evaluate each extremity is affected. Note (3): A rating of the determined by man	more	60		
Diastolic pressure predominantly 100 or more, or; systolic pressure predominantly 200 or more or more, or; systolic pressure predominantly 100 or more, or; systolic pressure predominantly 160 or more, or; systolic pressure predominantly 160 or more, or; systolic pressure predominantly 160 or more, or; minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the distolic blood pressure is predominantly 50mm. or greater, and isolated systolic blood pressure of less than 90mm.  NOTE (2): Exhese evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor, if applicable.  NOTE (3): Exhest each velocity each each extremity of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor, if applicable.  NOTE (3): Exhest each exhest each extremity is affected, evaluate each extremity be a single extremity. If more than one extremity is affected, evaluate each extremity bear applicable.  NOTE (3): Exhest each exhest each exhest each exhertenity is affected, evaluate ach extremity bear appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation causing it rather than by a separate evaluation.  NOTE: If symptomatic, evaluate according to body system affected.  NOTE: If symptomatic, evaluate residuals under the body system affected.  NOTE: If symptomatic, evaluate residuals under the body system affected.  NOTE: If symptomatic and t	Diastolic pressure predominantly 120 or			
Diastolic pressure predominantly 110 or more, or, systolic pressure predominantly 200 or more or misminum evaluation for an individual with a history of diastolic pressure predominantly 160 or more, or, systolic pressure predominantly 160 or more who requires continuous medication for control data of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation as of the date of hospital admission for surgical correction. Six months following discharge, the special correction. Six months following discharge, the appropriate	more	40		
Diastolic pressure predominantly 200 or more more or, systolic pressure predominantly 160 or more, or, minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 160mm, or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to prospersure is predominantly 160mm, which is usually the isolated systolic hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to opportant shall be assigned as of the date of hospital admission for surgical correction according to proportize distance and startery and solitor than a distolic blood pressure is predominantly 160mm. Argical proportized is predominantly 160mm, which is usually the isolated systolic hyperents with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hypertensive heart disease and other types of praft in sertion).  Precluding exertion.  Evaluate residuals				
Diastolic pressure predominantly 100 or more, or; systolic pressure predominantly 160 or more, or; minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 160mm. or greater, and isolated systolic by pertension means that the systolic blood pressure is predominantly 160mm, or greater, and isolated systolic blood pressure of less than 90mm. NoTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE: (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic byto pressure are devaluation.  NOTE: (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic byto pressure are devaluation.  NOTE: (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic byto pressure, and tachyroadia.  NOTE: (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic byto pressure are devaluation.  NOTE: (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic byto pressure, and tachycardia.  NOTE: (3): Evaluate hypertension factor, if symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to organ systems affected.  NOTE: A rating of 100				
more, or; systolic pressure predominantly 160 or more, or; minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control  NOTE (1): Hypertension or isolated systolic hyper- tension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predomi- nantly 90mm. or greater, and isolated systolic hyp- pertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insuffi- ciency or hyperthyroidism, which is usually the iso- laded systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension or pertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (3): Evaluate hypertension or means that the diastolic blood pressure is predominantly 160mm or greater with a diastolic blood pressure of less than 90mm.  NOTE (3): Evaluate hypertension or means that the diastolic blood pressure is predomi- nantly 90mm. or greater, and isolated systolic hyp- pertension means that the systolic blood pressure is predominantly 160mm or greater with a diastolic blood pressure of less than 90mm.  NOTE (3): Evaluate hypertension or  NOTE (3): Evaluate hypertension or supportant or any subsequent examination  NOTE (3): Evaluate hypertension due to aortic insuffi- ciency or hyperthyroidism, which is usually the iso- lated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension due to aortic insuffi- ciency or hyperthyroidism, which is usually the iso- lated systolic type, as part of the condition causing it rather than by a separate evaluation.  Precluding exertion  Evaluate residuals of surgical correction ac-		20	fected, evaluate each extremity separately and	
160 or more, or, minimum evaluation for an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control  NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension of weathing the subject to the provisions of § 3.105(e) of this chapter.  7112 Aneurysm, any small artery: Asymptomatic, evaluate according to body system affected. Following surgery, evaluate residuals under the body system affected.  7113 Arteriovenous fistula, traumatic: With high output heart failure Without cardiac involvement but with edema, stasis dermatitis, and either ulceration or cellulitis: Lower extremity Upper extremity With edema or stasis dermatitis: Lower extremity Upper extremity With edema or stasis dermatitis: Lower extremity Town or a least force of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension for production based upon that or any subsequent examination for surgical correction according to organ systems affected.  NOTE: If symptomatic, evaluate according to body system affected.  7113 Arteriovenous fistula, traumatic: With high output heart failure Without cardiac involvement but with eleman, stasis dermatitis; and either ulceration or cellulitis: Lower extremity Upper extremity Upper extremity  With edema or stasis dermatitis: Lower extremity  100 (1): Arterioscolomic produced in the produce				
an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control  NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (2): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or, if symptomatic, or, for indefinite period from date of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation shall be subject to the provisions of § 3.105(e) of this chapter.  7112 Aneurysm, any small artery:  Asymptomatic. evaluate according to body system affected.  7113 Arteriovenous fistula, traumatic:  Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia.  Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia.  Without cardiac involvement but with edema, stasis dermatitis.  Lower extremity  Upper extremity  With edema or stasis dermatitis:  Lower extremity  Upper extremity  With edema or stasis dermatitis:  Lower extremity  The development of the discovery of the extremity or analke/brachial index of 0.4 or less.  Claudication on walking between 25 and 100 yards on a level grade at 2 miles per hour, and; repitic changes (thin skin, absence of hair, dystrophic nails) or ankle/ brachial index of 0.7 o				
pressure predominantly 100 or more who requires continuous medication for control requires continuous medication for control requires continuous medication for control may be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the disatolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or, if symptomatic, or, for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion).  Precluding exertion  Evaluate residuals of surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination, Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this cha				
NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hypertension means that the disstolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate thypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)  Precluding exertion  Evaluate residuals of surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation and the organization based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7112 Aneurysm, any small artery:  Asymptomatic, evaluate according to body system affected.  7113 Arteriovenous fistula, traumatic:  With bigh output heart failure but with enlarged heart, wide pulse pressure, and tachycardia  Without cardiac involvement but with edema, stasis dermatitis:  Lower extremity.  Vipper extremity.  With edema or stasis dermatitis:  Lower extremity.  Vipper extremity.  100 4 or less.  Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; either deep ischemic ulcers or ankle/brachial index of 0.5 or less  Claudication on walking between 25 a				
NOTE (1): Hypertension or isolated systolic hypertension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension of sparately from hypertensive heart disease and other types of heart disease.  110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination. Any change in evaluation shall be subject to the provisions of § 3.105(e) of this chapter.  7112 Aneurysm, any small artery:  Asymptomatic.  7113 Anteriovenous fistula, traumatic:  With bigh output heart failure.  Without near failure but with enlarged heart, wide pulse pressure, and tachycardia.  With edema or stasis dermatitis:  Lower extremity  Upper extremity  Upper extremity  Vift edema or stasis dermatitis:  Lower extremity  Upper extremity  Total Anteriosolerosis obliterans:  Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; either persistent coldness of the extremity or ankle/brachial index of 0.5 or less		10		
tension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure is pred	·			
to or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 190mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)				
purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7111 Aneurysm, any small artery:  Asymptomatic.  NOTE: If symptomatic, evaluate according to body system affected.  7113 Arteriovenous fistula, traumatic:  Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia  Without cardiac involvement but with edema, stasis dermatitis; and either ulceration or cellulitis:  Lower extremity  Upper extremity  Upper extremity  Upper extremity  Upper extremity  100  7114 Arteriosclerosis obliterias:  Lower extremity  Upper extremity  or ankle/brachial index of 0.4 or less  Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; either persistent colcloses of the extremity or ankle/brachial index of 0.5 or less  Claudication on walking between 25 and 100 yards on a level grade at 2 miles per hour, and; either persistent colcloses of the extremity or				
means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion).  Precluding exertion  Evaluate residuals of surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or greater with a diastolic hoody system affected. Following surgery, evaluate residuals under the body system affected.  7113 Arteriovenous fistula, traumatic:  With high output heart failure but with enlarged heart, wide pulse pressure, and tachycardia  Without cardiac involvement but with edema, stasis dermatitis; and either ulceration or cellulitis:  Lower extremity  Upper extremity  With edema or stasis dermatitis:  Lower extremity  Upper extremity  Vith edema or stasis dermatitis:  Lower extremity  Upper extremity  Or alkel/brachial index of 0.4 or less  Claudication on walking less than 25 yards on a level grade at 2 miles per hour, and; either persistent coldness of the extremity or ankle/brachial index of 0.7 or less  Claudication on walking or ankle/brachial			of this chapter.	
nantly 90mm or greater, and isolated systolic hypertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systom affected.  7113 Arteriovenous fistula, traumatic:  With high output heart failure but with enlarged heart, wide pulse pressure, and tachycardia  Without cardiac involvement but with enlarged heart, wide pulse pressure, and tachycardia  Without cardiac involvement but with enlarged heart, wide pulse pressure, and tachycardia  Upper extremity  With edema or stasis dermatitis.  Lower extremity  With edema or stasis dermatitis.  100 4714 Arteriosclerosis obliterans:  Ischemic limb pain at rest, and; either deep ischemic limb pain at rest			7112 Aneurysm, any small artery:	
pertension means that the systolic blood pressure is predominantly 160mm. Or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to organ systems affected.  NOTE: If symptomatic, evaluate according to body system affected. Following surgery, evaluate residuals under the body system affected.  7113 Arteriovenous fistula, traumatic: With high output heart failure				0
system affected. Following surgery, evaluate residuals under the body system affected.  NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to organ systems affected.  7112 Arteriovenous fistula, traumatic:  Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia  Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia  Without cardiac involvement but with edema, stasis dermatitis:  Lower extremity  With edema or stasis dermatitis:  Lower extremity  Upper extremity  Vithe dema or stasis dermatitis:  Schemic limb pain at rest, and; either deep ischemic limb pain at rest, and; either perisstent coldness of the extremity or ank	pertension means that the systolic blood pressure		NOTE: If symptomatic evaluate according to body	
NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion).  Precluding exertion  Evaluate residuals of surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction form date of hospital admission for surgical correction on date of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.				
NOTE (2): Evaluate hypertension due to aortic insufficiency or hyperthyroidism, which is usually the isolated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)				
lated systolic type, as part of the condition causing it rather than by a separate evaluation.  NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)			, ,	
it rather than by a separate evaluation.  Note: (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)  Precluding exertion  Evaluate residuals of surgical correction according to organ systems affected.  Note: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction according to organ systems affected.  Without cardiac involvement but with enlarged heart, wide pulse pressure, and tachycardiawide pulse pressure, and tachycardia				100
NOTE (3): Evaluate hypertension separately from hypertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or, if symptomatic, or, for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)				100
pertensive heart disease and other types of heart disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)	, ·			60
disease.  7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)				
7110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)				
If five centimeters or larger in diameter, or; if symptomatic, or; for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)				
symptomatic, or, for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)	•		Lower extremity	50
date of hospital admission for surgical correction (including any type of graft insertion)			With adams or stocia dermetities	40
correction (including any type of graft insertion)				30
sertion)			Upper extremity	20
Precluding exertion		100		
Evaluate residuals of surgical correction according to organ systems affected.  NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or, for indefinite period from date of hospital admission for surgical correction		60	Ischemic limb pain at rest, and; either deep	
NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction				
NOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or, for indefinite period from date of hospital admission for surgical correction	cording to organ systems affected.			100
the date of admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or, for indefinite period from date of hospital admission for surgical correction	NOTE: A rating of 100 percent shall be assigned as of			
months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction				
ability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction				60
examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction				00
that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or, for indefinite period from date of hospital admission for surgical correction				
ject to the provisions of §3.105(e) of this chapter.  7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction				
7111 Aneurysm, any large artery:  If symptomatic, or; for indefinite period from date of hospital admission for surgical correction	ject to the provisions of §3.105(e) of this chapter.			
If symptomatic, or; for indefinite period from date of hospital admission for surgical correction	7111 Aneurysm, any large artery:			40
date of hospital admission for surgical correction				
correction				
		100		20
Following surgery: NOTE (1): The ankle/brachial index is the ratio of the			NOTE (1): The ankle/brachial index is the ratio of the	
Ischemic limb pain at rest, and; either deep systolic blood pressure at the ankle (determined by	Ischemic limb pain at rest, and; either deep			
ischemic ulcers or ankle/brachial index of Doppler study) divided by the simultaneous brach-				
0.4 or less	0.4 or less	100		

### 38 CFR Ch. I (7-1-07 Edition)

#### DISEASES OF THE HEART—Continued

#### DISEASES OF THE HEART—Continued

	Rat- ing		Rat- ing
NOTE (2): Evaluate residuals of aortic and large arterial bypass surgery or arterial graft as arteriosclerosis obliterans.  NOTE (3): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable.  7115 Thrombo-angiitis obliterans (Buerger's Disease):  Ischemic limb pain at rest, and; either deep ischemic ulcers or ankle/brachial index of 0.4 or less	100	7119 Erythromelalgia: Characteristic attacks that occur more than once a day, last an average of more than two hours each, respond poorly to treatment, and that restrict most routine daily activities  Characteristic attacks that occur more than once a day, last an average of more than two hours each, and respond poorly to treatment, but that do not restrict most routine daily activities  Characteristic attacks that occur daily or more often but that respond to treatment Characteristic attacks that occur less than daily but at least three times a week and	100
either persistent coldness of the extremity or ankle/brachial index of 0.5 or less  Claudication on walking between 25 and 100 yards on a level grade at 2 miles per hour, and; trophic changes (thin skin, absence of hair, dystrophic nails) or ankle/ brachial index of 0.7 or less  Claudication on walking more than 100 yards, and; diminished peripheral pulses or ankle/brachial index of 0.9 or less	40	that respond to treatment	11
NOTE (1): The ankle/brachial index of 0.9 of less NOTE (1): The ankle/brachial index is the ratio of the systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater.  NOTE (2): These evaluations are for involvement of a	20	7120 Varicose veins:  With the following findings attributed to the effects of varicose veins: Massive board-like edema with constant pain at rest  Persistent edema or subcutaneous induration, stasis pigmentation or eczema, and persistent ulceration	10
single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable.		Persistent edema and stasis pigmentation or eczema, with or without intermittent ulceration	41
7117 Raynaud's syndrome: With two or more digital ulcers plus autoamputation of one or more digits and history of characteristic attacks With two or more digital ulcers and history of characteristic attacks	100	elevation of extremity, with or without be- ginning stasis pigmentation or eczema Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by elevation of extremity or compression ho-	20
Characteristic attacks occurring at least daily  Characteristic attacks occurring four to six times a week  Characteristic attacks occurring one to three times a week	40 20 10	Siery	10
NOTE: For purposes of this section, characteristic at- tacks consist of sequential color changes of the digits of one or more extremities lasting minutes to hours, sometimes with pain and paresthesias, and precipitated by exposure to cold or by emotional upsets. These evaluations are for the disease as a whole, regardless of the number of extremities in-		combine (under §4.25), using the bilateral factor (§4.26), if applicable.  7121 Post-phlebitic syndrome of any etiology: With the following findings attributed to venous disease:  Massive board-like edema with	
volved or whether the nose and ears are involved.  7118 Angioneurotic edema:  Attacks without laryngeal involvement lasting one to seven days or longer and occurring more than eight times a year, or;		constant pain at rest	10
attacks with laryngeal involvement of any duration occurring more than twice a year Attacks without laryngeal involvement last- ing one to seven days and occurring five to eight times a year, or; attacks with la-	40	Persistent edema and stasis pig- mentation or eczema, with or without intermittent ulceration Persistent edema, incompletely re- lieved by elevation of extremity,	4
ryngeal involvement of any duration oc- curring once or twice a year	20	with or without beginning stasis pigmentation or eczema	2

DISEASES OF THE HEART—Continued

Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by elevation of extremity or compression hosiery

Asymptomatic palpable or visible varicose veins

NOTE: These evaluations are for involvement of a single extremity. If more than one extremity is involved, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable.

7122 Cold injury residuals:

With the following in affected parts:

Arthralgia or other pain, numbness, or cold sensitivity plus two or more of the following: tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteo-arthritis)

Arthralgia or other pain, numbness, or cold sensitivity plus tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, or X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteoarthritis)

Arthralgia or other pain, numbness, or cold sensitivity .....

NOTE (1): Separately evaluate amputations of fingers or toes, and complications such as squamous cell carcinoma at the site of a cold injury scar or peripheral neuropathy, under other diagnostic codes. Separately evaluate other disabilities that have been diagnosed as the residual effects of cold injury, such as Raynaud's phenomenon, muscle atrophy, etc., unless they are used to support an evaluation under diagnostic code 7122.

NOTE (2): Evaluate each affected part (e.g., hand, foot, ear, nose) separately and combine the ratings in accordance with §§ 4.25 and 4.26.

7123 Soft tissue sarcoma (of vascular origin) .....

NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.

(Authority: 38 U.S.C. 1155)

[62 FR 65219, Dec. 11, 1997, as amended at 63 FR 37779, July 14, 1998; 71 FR52460, Sept. 6, 2006]

THE DIGESTIVE SYSTEM

#### §4.110 Ulcers.

Rat-

Experience has shown that the term "peptic ulcer" is not sufficiently specific for rating purposes. Manifest differences in ulcers of the stomach or duodenum in comparison with those at an anastomotic stoma are sufficiently recognized as to warrant two separate graduated descriptions. In evaluating the ulcer, care should be taken that the findings adequately identify the particular location.

#### §4.111 Postgastrectomy syndromes.

There are various postgastrectomy symptoms which may occur following anastomotic operations of the stomach. When present, those occurring during or immediately after eating and known as the "dumping syndrome" are characterized by gastrointestinal complaints and generalized symptoms simulating hypoglycemia; those occurring from 1 to 3 hours after eating usually present definite manifestations of hypoglycemia.

#### §4.112 Weight loss.

For purposes of evaluating conditions in §4.114, the term "substantial weight loss" means a loss of greater than 20 percent of the individual's baseline weight, sustained for three months or longer; and the term "minor weight loss" means a weight loss of 10 to 20 percent of the individual's baseline weight, sustained for three months or longer. The term "inability to gain weight" means that there has been substantial weight loss with inability to regain it despite appropriate therapy. "Baseline weight" means the average weight for the two-year-period preceding onset of the disease.

(Authority: 38 U.S.C. 1155) [66 FR 29488, May 31, 2001]

## §4.113 Coexisting abdominal conditions.

There are diseases of the digestive system, particularly within the abdomen, which, while differing in the site of pathology, produce a common disability picture characterized in the main by varying degrees of abdominal

distress or pain, anemia and disturbances in nutrition. Consequently, certain coexisting diseases in this area, as indicated in the instruction under the title "Diseases of the Digestive System," do not lend themselves to distinct and separate disability evaluations without violating the fundamental principle relating to pyramiding as outlined in §4.14.

# \$4.114 Schedule of ratings—digestive system.

Ratings under diagnostic codes 7301 to 7329, inclusive, 7331, 7342, and 7345 to 7348 inclusive will not be combined with each other. A single evaluation will be assigned under the diagnostic code which reflects the predominant disability picture, with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation.

7200 Mouth, injuries of.
Rate as for disfigurement and impairment of
function of mastication.
7201 Lips, injuries of.
Rate as for disfigurement of face.
7202 Tongue, loss of whole or part:
With inability to communicate by speech
One-half or more
7203 Esophagus, stricture of:
Permitting passage of liquids only, with marked
impairment of general health
Severe, permitting liquids only
Moderate
7204 Esophagus, spasm of (cardiospasm).
If not amenable to dilation, rate as for the de-
gree of obstruction (stricture).
7205 Esophagus, diverticulum of, acquired.
Rate as for obstruction (stricture).
7301 Peritoneum, adhesions of:
Severe; definite partial obstruction shown by X-
ray, with frequent and prolonged episodes of
severe colic distension, nausea or vomiting,
following severe peritonitis, ruptured appendix,
perforated ulcer, or operation with drainage
Moderately severe; partial obstruction mani-
fested by delayed motility of barium meal and
less frequent and less prolonged episodes of
pain
Moderate; pulling pain on attempting work or ag-
gravated by movements of the body, or occa-
sional episodes of colic pain, nausea, con-
stipation (perhaps alternating with diarrhea) or
abdominal distension
Mild
Note: Ratings for adhesions will be considered
when there is history of operative or other
traumatic or infectious (intraabdominal) proc-
ess, and at least two of the following: disturb-
ance of motility, actual partial obstruction, re-
flex disturbances, presence of pain.
7304 Ulcer, gastric.
7304 Olcer, gastric. 7305 Ulcer, duodenal:
7303 Oicer, duodenai:

	Rat- ing
Severe; pain only partially relieved by standard ulcer therapy, periodic vomiting, recurrent hematemesis or melena, with manifestations of anemia and weight loss productive of defi-	
nite impairment of health	60
at least four or more times a year	40
manifestations	20
yearly	10
7306 Ulcer, marginal (gastrojejunal): Pronounced; periodic or continuous pain unrelieved by standard ulcer therapy with peri- odic vomiting, recurring melena or hematemesis, and weight loss. Totally inca-	
pacitating	100
nounced and less continuous symptoms with definite impairment of health	60
completely relieved by ulcer therapy, mild and transient episodes of vomiting or melena	40
Moderate; with episodes of recurring symptoms several times a year	20
Mild; with brief episodes of recurring symptoms once or twice yearly	10
gastroscope): Chronic; with severe hemorrhages, or large ul-	60
Chronic; with multiple small eroded or ulcerated	30
areas, and symptoms	
toms	10
7308 Postgastrectomy syndromes: Severe; associated with nausea, sweating, cir- culatory disturbance after meals, diarrhea,	
hypoglycemic symptoms, and weight loss with malnutrition and anemia	60
symptoms after meals but with diarrhea and weight loss	40
with characteristic mild circulatory symptoms or continuous mild manifestations	20
7310 Stomach, injury of, residuals. Rate as peritoneal adhesions.	
7311 Residuals of injury of the liver: Depending on the specific residuals, separately evaluate as adhesions of peritoneum (diag- nostic code 7301), cirrhosis of liver (diagnostic	
code 7312), and chronic liver disease without cirrhosis (diagnostic code 7345).  7312 Cirrhosis of the liver, primary biliary cirrhosis,	

ing

100

30

10

	Rat- ing		Rat- ing
Generalized weakness, substantial weight loss,		Moderate symptoms	10
and persistent jaundice, or; with one of the fol-		Mild or no symptoms	0
lowing refractory to treatment: ascites, hepatic		7325 Enteritis, chronic.	
encephalopathy, hemorrhage from varices or	400	Rate as for irritable colon syndrome.	
portal gastropathy (erosive gastritis)	100	7326 Enterocolitis, chronic.	
History of two or more episodes of ascites, he- patic encephalopathy, or hemorrhage from		Rate as for irritable colon syndrome.	
varices or portal gastropathy (erosive gas-		7327 Diverticulitis.  Rate as for irritable colon syndrome, peritoneal	
tritis), but with periods of remission between		adhesions, or colitis, ulcerative, depending	
attacks	70	upon the predominant disability picture.	
History of one episode of ascites, hepatic		7328 Intestine, small, resection of:	
encephalopathy, or hemorrhage from varices		With marked interference with absorption and	
or portal gastropathy (erosive gastritis)	50	nutrition, manifested by severe impairment of	
Portal hypertension and splenomegaly, with weakness, anorexia, abdominal pain, malaise,		health objectively supported by examination	
and at least minor weight loss	30	findings including material weight loss	60
Symptoms such as weakness, anorexia, abdom-	30	With definite interference with absorption and nutrition, manifested by impairment of health	
inal pain, and malaise	10	objectively supported by examination findings	
NOTE: For evaluation under diagnostic code		including definite weight loss	40
7312, documentation of cirrhosis (by biopsy or		Symptomatic with diarrhea, anemia and inability	.5
imaging) and abnormal liver function tests		to gain weight	20
must be present.		Note: Where residual adhesions constitute the	
7314 Cholecystitis, chronic: Severe; frequent attacks of gall bladder colic	30	predominant disability, rate under diagnostic	
Moderate; gall bladder dyspepsia, confirmed by	30	code 7301.	
X-ray technique, and with infrequent attacks		7329 Intestine, large, resection of:	
(not over two or three a year) of gall bladder		With severe symptoms, objectively supported by examination findings	40
colic, with or without jaundice	10	With moderate symptoms	20
Mild	0	With slight symptoms	10
7315 Cholelithiasis, chronic.		NOTE: Where residual adhesions constitute the	
Rate as for chronic cholecystitis. 7316 Cholangitis, chronic.		predominant disability, rate under diagnostic	
Rate as for chronic cholecystitis.		code 7301.	
7317 Gall bladder, injury of.		7330 Intestine, fistula of, persistent, or after attempt	
Rate as for peritoneal adhesions.		at operative closure:  Copious and frequent, fecal discharge	100
7318 Gall bladder, removal of:		Constant or frequent, fecal discharge	60
With severe symptoms	30	Slight infrequent, fecal discharge	30
With mild symptoms  Nonsymptomatic	10	Healed; rate for peritoneal adhesions.	
Spleen, disease or injury of.		7331 Peritonitis, tuberculous, active or inactive:	
See Hemic and Lymphatic Systems.		Active	100
7319 Irritable colon syndrome (spastic colitis, mu-		Inactive: See §§ 4.88b and 4.89.	
cous colitis, etc.):		7332 Rectum and anus, impairment of sphincter control:	
Severe; diarrhea, or alternating diarrhea and		Complete loss of sphincter control	100
constipation, with more or less constant ab-	30	Extensive leakage and fairly frequent involuntary	100
dominal distress	30	bowel movements	60
ance with abdominal distress	10	Occasional involuntary bowel movements, ne-	
Mild; disturbances of bowel function with occa-	-	cessitating wearing of pad	30
sional episodes of abdominal distress	0	Constant slight, or occasional moderate leakage	10
7321 Amebiasis:		Healed or slight, without leakage	0
Mild gastrointestinal disturbances, lower abdom-		Requiring colostomy	100
inal cramps, nausea, gaseous distention, chronic constipation interrupted by diarrhea	10	Great reduction of lumen, or extensive leakage	50
Asymptomatic	0	Moderate reduction of lumen, or moderate con-	
NOTE: Amebiasis with or without liver abscess is		stant leakage	30
parallel in symptomatology with ulcerative coli-		7334 Rectum, prolapse of:	
tis and should be rated on the scale provided		Severe (or complete), persistent	50
for the latter. Similarly, lung abscess due to		Moderate, persistent or frequently recurring	30
amebiasis will be rated under the respiratory		Mild with constant slight or occasional moderate leakage	10
system schedule, diagnostic code 6809.  7322 Dysentery, bacillary.		7335 Ano, fistula in.	
Rate as for ulcerative colitis		Rate as for impairment of sphincter control.	
7323 Colitis, ulcerative:		7336 Hemorrhoids, external or internal:	
Pronounced; resulting in marked malnutrition,		With persistent bleeding and with secondary	
anemia, and general debility, or with serious		anemia, or with fissures	20
complication as liver abscess	100	Large or thrombotic, irreducible, with excessive	
Severe; with numerous attacks a year and mal-	60	redundant tissue, evidencing frequent recurrences	10
nutrition, the health only fair during remissions Moderately severe; with frequent exacerbations	60 30	Mild or moderate	0
Moderate; with infrequent exacerbations	10	7337 Pruritus ani.	3
7324 Distomiasis, intestinal or hepatic:		Rate for the underlying condition.	
Severe symptoms	30	7338 Hernia, inguinal:	

	Rat- ing		Rat-
Large, postoperative, recurrent, not well supported under ordinary conditions and not readily reducible, when considered inoperable  Small, postoperative recurrent, or unoperated irremediable, not well supported by truss, or not readily reducible.  Postoperative recurrent, readily reducible and well supported by truss or belt.  Not operated, but remediable	60 30 10 0	Daily fatigue, malaise, and anorexia, with minor weight loss and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least four weeks, but less than six weeks, during the past 12-month period.  Daily fatigue, malaise, and anorexia (without weight loss or hepatomegaly), requiring dietary restriction or continuous medication, or; incapacitating episodes (with symptoms such as	40
NOTE: Add 10 percent for bilateral involvement, provided the second hemia is compensable. This means that the more severely disabling hemia is to be evaluated, and 10 percent, only, added for the second hemia, if the latter is of compensable degree.  7339 Hemia, ventral, postoperative:		fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least two weeks, but less than four weeks, during the past 12-month period	20
Massive, persistent, severe diastasis of recti muscles or extensive diffuse destruction or weakening of muscular and fascial support of abdominal wall so as to be inoperable	100	incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, ano- rexia, arthralgia, and right upper quadrant pain) having a total duration of at least one week, but less than two weeks, during the	
Large, not well supported by belt under ordinary conditions	40	past 12-month period	10 0
erative wounds with weakening of abdominal wall and indication for a supporting belt	20 0	same signs and symptoms as the basis for evaluation under DC 7354 and under a diagnostic code for sequelae. (See § 4.14.).	
7340 Hernia, femoral. Rate as for inguinal hernia.	10	NOTE (2): For purposes of evaluating conditions under diagnostic code 7345, "incapacitating episode" means a period of acute signs and	
7342 Visceroptosis, symptomatic, marked	100	symptoms severe enough to require bed rest and treatment by a physician. NOTE (3): Hepatitis B infection must be con- firmed by serologic testing in order to evaluate	
antineoplastic chemotherapy or other thera- peutic procedure. Six months after discontinu- ance of such treatment, the appropriate dis- ability rating shall be determined by mandatory		it under diagnostic code 7345.  7346 Hernia hiatal: Symptoms of pain, vomiting, material weight loss and hematemesis or melena with moderate anemia; or other symptom combinations pro-	
VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no		ductive of severe impairment of health	60
local recurrence or metastasis, rate on residu- als.  7344 Benign neoplasms, exclusive of skin growths:		productive of considerable impairment of health	30
Evaluate under an appropriate diagnostic code, depending on the predominant disability or the specific residuals after treatment.		percent evaluation of less severity	10
7345 Chronic liver disease without cirrhosis (including hepatitis B, chronic active hepatitis, autoimmune hepatitis, hemochromatosis, drug-induced hepatitis, etc., but excluding bile duct disorders and hepatitis C):		dominal pain with few pain free intermissions and with steatorrhea, malabsorption, diarrhea and severe malnutrition	100
Near-constant debilitating symptoms (such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain)	100	tween acute attacks	60
Daily fatigue, malaise, and anorexia, with substantial weight loss (or other indication of malnutrition), and hepatomegaly, or; incapacitating		mission between attacks	30 10
episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) having a total duration of at least six weeks during the past 12-month period, but not occurring		NOTE 1: Abdominal pain in this condition must be confirmed as resulting from pancreatitis by appropriate laboratory and clinical studies. NOTE 2: Following total or partial pancrea-	
constantly	60	tectomy, rate under above, symptoms, min- imum rating 30 percent. 7348 Vagotomy with pyloroplasty or gastro- enterostomy:	

Followed by demonstrably confirmative post- operative complications of stricture or con- tinuing gastric retention	
With symptoms and confirmed diagnosis of alka- line gastritis, or of confirmed persisting diar- rhea	
Recurrent ulcer with incomplete vagotomy	
NOTE: Rate recurrent ulcer following complete vagotomy under diagnostic code 7305, min- imum rating 20 percent; and rate dumping	
syndrome under diagnostic code 7308.	
7351 Liver transplant:  For an indefinite period from the date of hospital	
admission for transplant surgery	
Minimum	
NOTE: A rating of 100 percent shall be assigned as of the date of hospital admission for trans-	
plant surgery and shall continue. One year fol- lowing discharge, the appropriate disability rat-	
ing shall be determined by mandatory VA examination. Any change in evaluation based	
upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of	
this chapter. 7354 Hepatitis C (or non-A, non-B hepatitis):	
7354 Hepatitis C (or non-A, non-B hepatitis): With serologic evidence of hepatitis C infection	
and the following signs and symptoms due to	
hepatitis C infection:	
Near-constant debilitating symptoms (such as	
fatigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain)	
Daily fatigue, malaise, and anorexia, with sub- stantial weight loss (or other indication of mal- nutrition), and hepatomegaly, or; incapacitating episodes (with symptoms such as fatigue,	
malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) hav- ing a total duration of at least six weeks during the past 12-month period, but not occurring	
constantly  Daily fatigue, malaise, and anorexia, with minor	
weight loss and hepatomegaly, or; incapaci- tating episodes (with symptoms such as fa-	
tigue, malaise, nausea, vomiting, anorexia, arthralgia, and right upper quadrant pain) hav-	
ing a total duration of at least four weeks, but	
less than six weeks, during the past 12-month	
period  Daily fatigue, malaise, and anorexia (without	
weight loss or hepatomegaly), requiring dietary restriction or continuous medication, or; inca-	
pacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, anorexia,	
arthralgia, and right upper quadrant pain) hav-	
ing a total duration of at least two weeks, but	
less than four weeks, during the past 12- month period	
Intermittent fatigue, malaise, and anorexia, or;	
incapacitating episodes (with symptoms such as fatigue, malaise, nausea, vomiting, ano- rexia, arthralgia, and right upper quadrant	
pain) having a total duration of at least one week, but less than two weeks, during the	
past 12-month period	

Nonsymptomatic ...

	Rat- ing
NOTE (1): Evaluate sequelae, such as cirrhosis or malignancy of the liver, under an appropriate diagnostic code, but do not use the same signs and symptoms as the basis for evaluation under DC 7354 and under a diagnostic code for sequelae. (See § 4.14.).  NOTE (2): For purposes of evaluating conditions under diagnostic code 7354, "incapacitating episode" means a period of acute signs and symptoms severe enough to require bed rest and treatment by a physician.	

(Authority: 38 U.S.C. 1155)

ing

40

30 20

100

[29 FR 6718, May 22, 1964, as amended at 34 FR 5063, Mar. 11, 1969; 40 FR 42540, Sept. 15, 1975; 41 FR 11301, Mar. 18, 1976; 66 FR 29488, May 31, 2001]

THE GENITOURINARY SYSTEM

#### §4.115 Nephritis.

Albuminuria alone is not nephritis, nor will the presence of transient albumin and casts following acute febrile illness be taken as nephritis. The glomerular type of nephritis is usually preceded by or associated with severe infectious disease; the onset is sudden, and the course marked by red blood cells, salt retention, and edema; it may clear up entirely or progress to a chronic condition. The nephrosclerotic type, originating in hypertension or arteriosclerosis, develops slowly, with minimum laboratory findings, and is associated with natural progress. Separate ratings are not to be assigned for disability from disease of the heart and any form of nephritis, on account of the close interrelationships of cardiovascular disabilities. If, however, absence of a kidney is the sole renal disability, even if removal was required because of nephritis, the absent kidney and any hypertension or heart disease will be separately rated. Also, in the event that chronic renal disease has progressed to the point where regular dialysis is required, any coexisting hypertension or heart disease will be separately rated.

[41 FR 34258, Aug. 13, 1976, as amended at 59 FR 2527, Jan. 18, 1994]

## §4.115a Ratings of the genitourinary system—dysfunctions.

Diseases of the genitourinary system generally result in disabilities related

10

0

#### §4.115b

to renal or voiding dysfunctions, infections, or a combination of these. The following section provides descriptions of various levels of disability in each of these symptom areas. Where diagnostic codes refer the decisionmaker to these specific areas dysfunction, only the predominant area of dysfunction shall be considered for rating purposes. Since the areas of dysfunction described below do not cover all symptoms resulting from genitourinary diseases, specific diagnoses may include a description of symptoms assigned to that diagnosis.

	Rat- ing
Renal dysfunction:	
Requiring regular dialysis, or precluding more than sedentary activity from one of the following: persistent edema and albuminuria; or, BUN more than 80mg%; or, creatinine more than 8mg%; or, markedly decreased function of kidney or other organ systems, estpecially	400
cardiovascular  Persistent edema and albuminuria with BUN 40 to 80mg%; or, creatinine 4 to 8mg%; or, generalized poor health characterized by lethargy, weakness, anorexia, weight loss, or limitation of exertion	100
Constant albuminuria with some edema; or, defi- nite decrease in kidney function; or, hyper- tension at least 40 percent disabling under di-	
agnostic code 7101	60
or slight edema or hypertension at least 10 percent disabling under diagnostic code 7101 Albumin and casts with history of acute nephritis; or, hypertension non-compensable under diag-	30
nostic code 7101	0
Voiding dysfunction: Rate particular condition as urine leakage, fre- quency, or obstructed voiding Continual Urine Leakage, Post Surgical Urinary Diversion, Urinary Incontinence, or Stress In- continence:	
Requiring the use of an appliance or the wearing of absorbent materials which must be changed more than 4 times per day	60
Requiring the wearing of absorbent materials which must be changed 2 to 4 times per day Requiring the wearing of absorbent materials	40
which must be changed less than 2 times per day	20
Daytime voiding interval less than one hour, or; awakening to void five or more times per night Daytime voiding interval between one and two	40
hours, or; awakening to void three to four times per night	20
hours, or; awakening to void two times per night	10
Urinary retention requiring intermittent or contin-	
uous catheterization	30

	Rat- ing
Marked obstructive symptomatology (hesitancy, slow or weak stream, decreased force of stream) with any one or combination of the fol- lowing:	
<ol> <li>Post void residuals greater than 150 cc.</li> <li>Uroflowmetry; markedly diminished peak flow rate (less than 10 cc/sec).</li> </ol>	
<ol><li>Recurrent urinary tract infections sec- ondary to obstruction.</li></ol>	
Stricture disease requiring periodic dilatation every 2 to 3 months	10
Obstructive symptomatology with or without stric- ture disease requiring dilatation 1 to 2 times	
per yearrninary tract infection:	0
Poor renal function: Rate as renal dysfunction.  Recurrent symptomatic infection requiring drainage/frequent hospitalization (greater than two	
times/year), and/or requiring continuous intensive management	30
year and/or requiring intermittent intensive management	10

[59 FR 2527, Jan. 18, 1994; 59 FR 10676, Mar. 7,

#### §4.115b Ratings of the genitourinary system—diagnoses.

		Rat- ing
Note: When evaluating an volving loss or loss of us more creative organs, § 3.350 of this chapter to whether the veteran mattled to special monthly tion. Footnotes in the school cate conditions which establish entitlement monthly compensation; there are other conditions ection which under coumstances also establishent to special monthly tion.	e of one or refer to determine y be enti- compensa- nedule indi- potentially to special however, ons in this tertain cir- ish entitle-	
7500 Kidney, removal of one: Minimum evaluation Or rate as renal dysfunction nephritis, infection, or pa the other.		30
7501 Kidney, abscess of: Rate as urinary tract infecti	on	

100

80

60

30

0

40

20

40

20

10

§4.1	15b
------	-----

	Rat- ing		Rat- ing
7502 Nephritis, chronic:		7517 Bladder, injury of:	
Rate as renal dysfunction.		Rate as voiding dysfunction.	
7504 Pyelonephritis, chronic:		7518 Urethra, stricture of:	
Rate as renal dysfunction or urinary		Rate as voiding dysfunction.	
tract infection, whichever is pre- dominant.		7519 Urethra, fistual of:	
7505 Kidney, tuberculosis of:		Rate as voiding dysfunction.	
Rate in accordance with §§ 4.88b or		Multiple urethroperineal fistulae	100
4.89, whichever is appropriate.		7520 Penis, removal of half or more	30
7507 Nephrosclerosis, arteriolar:		Or rate as voiding dysfunction.	
Rate according to predominant symp-		7521 Penis removal of glans	20
toms as renal dysfunction, hyper-		Or rate as voiding dysfunction.	
tension or heart disease. If rated		7522 Penis, deformity, with loss of erectile	
under the cardiovascular schedule,		power—201.	
however, the percentage rating		7523 Testis, atrophy complete:.	
which would otherwise be assigned		Both—201	
will be elevated to the next higher		One—01	
evaluation.		7524 Testis, removal:.	
7508 Nephrolithiasis:		Both—301	
Rate as hydronephrosis, except for		One—01	
recurrent stone formation requiring		Orie—0	
one or more of the following:  1. diet therapy		Note: In cases of the removal of one	
2. drug therapy		testis as the result of a service-in-	
3. invasive or non-invasive proce-		curred injury or disease, other than	
dures more than two times/year	30	an undescended or congenitally un-	
7509 Hydronephrosis:	00	developed testis, with the absence	
Severe; Rate as renal dysfunction.		or nonfunctioning of the other testis	
Frequent attacks of colic with infection		unrelated to service, an evaluation	
(pyonephrosis), kidney function im-		of 30 percent will be assigned for	
paired	30	the service-connected testicular	
Frequent attacks of colic, requiring cath-		loss. Testis, underscended, or con-	
eter drainage	20	genitally undeveloped is not a rat- able disability.	
Only an occasional attack of colic, not		able disability.	
infected and not requiring catheter	40	7525 Epididymo-orchitis, chronic only:	
drainage	10	Rate as urinary tract infection.	
Rate as hydronephrosis, except for		For tubercular infections: Rate in ac-	
recurrent stone formation requiring		cordance with §§ 4.88b or 4.89,	
one or more of the following:		whichever is appropriate.	
1. diet therapy		7527 Prostate gland injuries, infections, hy-	
2. drug therapy		pertrophy, postoperative residuals:	
3. invasive or non-invasive proce-		Rate as voiding dysfunction or urinary	
dures more than two times/year	30	tract infection, whichever is pre-	
7511 Ureter, stricture of:		dominant.	
Rate as hydronephrosis, except for		7528 Malignant neoplasms of the genito-	
recurrent stone formation requiring		urinary system	100
one or more of the following:			
1. diet therapy			
drug therapy     invasive or non-invasive proce-			
dures more than two times/year	30		
7512 Cystitis, chronic, includes interstitial	50		
and all etiologies, infectious and non-in-			
fectious:			
Rate as voiding dysfunction.			
7515 Bladder, calculus in, with symptoms			
interfering with function:			
Rate as voiding dysfunction			
7516 Bladder, fistula of:			
Rate as voiding dysfunction or urinary			
tract infection, whichever is pre-			
dominant.	400		
Postoperative, suprapubic cystotomy	100		

### 38 CFR Ch. I (7-1-07 Edition)

	Rat- ing		Rat- ing
Note—Following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure, the rating of 100 percent shall continue with a mandatory VA examination at the expiration of six months. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local reoccurrence or metastasis, rate on residuals as voiding dysfunction or renal dysfunction, whichever is predominant.  7529 Benign neoplasms of the genitourinary system: Rate as voiding dysfunction or renal dysfunction, whichever is predominant.  7530 Chronic renal disease requiring regular dialysis: Rate as renal dysfunction.  7531 Kidney transplant: Following transplant surgery	100 30	Or rate as renal dysfunction.  7533 Cystic diseases of the kidneys (polycystic disease, uremic medullary cystic disease, Medullary sponge kidney, and similar conditions):  Rate as renal dysfunction.  7534 Atherosclerotic renal disease (renal artery stenosis or atheroembolic renal disease):  Rate as renal dysfunction.  7535 Toxic nephropathy (antibotics, radiocontrast agents, nonsteroidal anti-inflammatory agents, heavy metals, and similar agents):  Rate as renal dysfunction.  7536 Glomerulonephritis:  Rate as renal dysfunction.  7537 Interstitial nephritis:  Rate as renal dysfunction.  7538 Papillary necrosis:  Rate as renal dysfunction.  7539 Renal amyloid disease:  Rate as renal dysfunction.  7540 Disseminated intravascular coagulation with renal cortical necrosis:  Rate as renal dysfunction.  7541 Renal involvement in diabetes mellitus, sickle cell anemia, systemic lupus erythematosus, vasculitis, or other systemic disease processes.  Rate as renal dysfunction.  7542 Neurogenic bladder:  Rate as voiding dysfunction.  1 Review for entitlement to special monthl pensation under § 3.350 of this chapter.	y com- Iar. 29,

Minimum rating for symptomatic condition

GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

# §4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

		Rating
N	ote 1: Natural menopause, primary amenorrhea, and pregnancy and child-birth are not disabilities for rating purposes. Chronic residuals of medical or surgical complications of pregnancy may be disabilities for rating purposes. ote 2: When evaluating any claim involving loss or loss of use of one or more creative organs or anatomical loss of one or both breasts, refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, almost any condition in this section might, under certain circumstances, establish entitlement to special monthly compensation.	
7610 Vulvovaç		
	jina, disease or injury of.	
	vix, disease or injury of.	
	rus, disease, injury, or adhesions of. opian tube, disease, injury, or adhesions	
of (include	ding pelvic inflammatory disease (PID)).	
7615 Ova	ary, disease, injury, or adhesions of. ating Formula for Disease, Injury, or Ad-	
hesions	of Female Reproductive Organs (diag-	
nostic co	des 7610 through 7615):	
S	ymptoms not controlled by continuous	30
S	treatmentymptoms that require continuous treat-	30
	ment	10
S	ymptoms that do not require continuous	_
7617 Ute	treatmentrus and both ovaries, removal of, com-	0
plete:		
	or three months after removal	1100
	nereafterrus, removal of, including corpus:	<sup>1</sup> 50
	or three months after removal	1100
TI	nereafter	<sup>1</sup> 30
	ary, removal of:	1100
	or three months after removal	100
	Complete removal of both ovaries	1 30
	Removal of one with or without	1.0
7620 Ova	partial removal of the other	<sup>1</sup> 0
	rus, prolapse:	
	omplete, through vagina and introitus	50
	complete	30
	rus, displacement of: lith marked displacement and frequent or	
	continuous menstrual disturbances	30
	ith adhesions and irregular menstruation	10
	gnancy, surgical complications of: lith rectocele or cystocele	50
	ith rectocere of cystocere	10
7624 Fist	ula, rectovaginal:	.0
	aginal fecal leakage at least once a day	
17	requiring wearing of pad	100
V	aginal fecal leakage four or more times per week, but less than daily, requiring	
	wearing of pad	60

•	
	Rating
Vaginal fecal leakage one to three times per week requiring wearing of pad	30
Vaginal fecal leakage less than once a week	10
Without leakage	0
7625 Fistula, urethrovaginal:	ľ
Multiple urethrovaginal fistulae	100
Requiring the use of an appliance or the	
wearing of absorbent materials which must be changed more than four times	
per day	60
Requiring the wearing of absorbent mate- rials which must be changed two to four times per day	40
Requiring the wearing of absorbent mate-	40
rials which must be changed less than two times per day	20
7626 Breast, surgery of:	
Following radical mastectomy:	
Both	180
One	¹50
Following modified radical mastectomy:	
Both	¹60
One	<sup>1</sup> 40
Following simple mastectomy or wide local	
excision with significant alteration of size	
or form:	150
Both	<sup>1</sup> 50
One Following wide local excision without sig-	.30
nificant alteration of size or form:	
Both or one	0
Note: For VA purposes:	
(1) Radical mastectomy means	
removal of the entire breast, un-	
derlying pectoral muscles, and regional lymph nodes up to the	
regional lymph nodes up to the	
coracoclavicular ligament	
(2) Modified radical mastectomy means removal of the entire	
breast and axillary lymph nodes	
(in continuity with the breast).	
Pectoral muscles are left intact	
(3) Simple (or total) mastectomy	
means removal of all of the	
breast tissue, nipple, and a small portion of the overlying	
skin, but lymph nodes and mus-	
cles are left intact	
(4) Wide local excision (including	
partial mastectomy,	
lumpectomy, tylectomy,	
segmentectomy, and	
quadrantectomy) means re-	
moval of a portion of the breast tissue	
7627 Malignant neoplasms of gynecological sys-	
tem or breast	100

#### §4.117

7628 Benign neoplasms of the gynecological system or breast. Rate according to impairment in function of the urinary or gynecological systems, or skin.

#### 7629 Endometriosis:

Lesions involving bowel or bladder confirmed by laparoscopy, pelvic pain or heavy or irregular bleeding not controlled by treatment, and bowel or bladder symptoms

Pelvic pain or heavy or irregular bleeding not controlled by treatment ..

Pelvic pain or heavy or irregular bleeding requiring continuous treatment for control Note: Diagnosis of endometriosis must be substantiated by laparoscopy.

[60 FR 19855, Apr. 21, 1995, as amended at 67 FR 6874, Feb. 14, 2002; 67 FR 37695, May 30,

#### THE HEMIC AND LYMPHATIC SYSTEMS

#### §4.117 Schedule of ratings-hemic and lymphatic systems.

	Rating
7700 Anemia, hypochromic-microcytic and megaloblastic, such as iron-deficiency and pernicious anemia:	
Hemoglobin 5gm/100ml or less, with find- ings such as high output congestive heart failure or dyspnea at rest	100
cardiomegaly, tachycardia (100 to 120 beats per minute) or syncope (three episodes in the last six months)	70
headaches, lightheadedness, or short- ness of breath Hemoglobin 10gm/100ml or less with find-	30
ings such as weakness, easy fatigability or headaches	10
tomatic	0
NOTE: Evaluate complications of pernicious anemia,	Such as

dementia or peripheral neuropathy, separately.

7702 Agranulocytosis, acute:

Requiring bone marrow transplant, or; requiring transfusion of platelets or red cells at least once every six weeks, or; infections recurring at least once every six weeks ...

	Rating
Requiring transfusion of platelets or red cells at least once every three months, or; infections recurring at least once every three months	60
than once every three months	30
Requiring continuous medication for control	10

NOTE: The 100 percent rating for bone marrow transplant shall be assigned as of the date of hospital admission and shall continue with a mandatory VA examination six months following hospital discharge. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.

#### 7703 Leukemia: With active disease or during a treatment 100 phase Otherwise rate as anemia (code 7700) or aplastic anemia (code 7716), whichever would result in the greater benefit.

NOTE: The 100 percent rating shall continue beyond the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no recurrence, rate on residuals.

#### 7704 Polycythemia vera: During periods of treatment with myelosuppressants and for three months following cessation of myelosuppressant therapy 100 Requiring phlebotomy . 40 Stable, with or without continuous medica-10

NOTE: Rate complications such as hypertension, gout, stroke or thrombotic disease separately.

7705 Thrombocytopenia, primary, idiopathic or	im-
mune:	
Platelet count of less than 20,000, with	ac-
tive bleeding, requiring treatment v	vith
medication and transfusions	100
Platelet count between 20,000 and 70,0	100,
not requiring treatment, without bleedi	ng 70
Stable platelet count between 70,000 a	and
100,000, without bleeding	30
Stable platelet count of 100,000 or mo	ore,
without bleeding	0
7706 Splenectomy	
NOTE: Rate complications such as systemic inf encapsulated bacteria separately.	ections with
7707 Spleen, injury of, healed.	
Rate for any residuals.	
7709 Hodgkin's disease:	
With active disease or during a treatm	ent

NOTE: The 100 percent rating shall continue beyond the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recur-

rence or metastasis, rate on residuals.

100

50

30

10

<sup>&</sup>lt;sup>1</sup>Review for entitlement to special monthly compensation under § 3.350 of this chapter.

§4.118

		Rating
7710	Rate under §§4.88c or 4.89 of this part, whichever is appropriate.	
7714	With repeated painful crises, occurring in skin, joints, bones or any major organs caused by hemolysis and sickling of red blood cells, with anemia, thrombosis and infarction, with symptoms precluding	100
	even light manual labor	60
	light manual labor Following repeated hemolytic sickling crises with continuing impairment of health	30
	Asymptomatic, established case in remission, but with identifiable organ impairment	10
tribu Cas the	Sickle cell trait alone, without a history of directable pathological findings, is not a ratable dises of symptomatic sickle cell trait will be forware Director, Compensation and Pension Service, teration under § 3.321(b)(1) of this chapter.	sability. Irded to
7715	Non-Hodgkin's lymphoma: With active disease or during a treatment phase	100

NOTE: The 100 percent rating shall continue beyond the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be dtermined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.

7716 Aplastic anemia: Requiring bone marrow transplant, or; requiring transfusion of platelets or red cells at least once every six weeks, or; infections recurring at least once every six weeks .. 100 Requiring transfusion of platelets or red cells at least once every three months, or; infections recurring at least once every three months .... 60 Requiring transfusion of platelets or red cells at least once per year but less than once every three months, or; infections recurring at least once per year but less than once every three months 30 Requiring continuous medication for control 10

NOTE: The 100 percent rating for bone marrow transplant shall be assigned as of the date of hospital admission and shall continue with a mandatory VA examination six months following hospital discharge. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this

[60 FR 49227, Sept. 22, 1995]

#### THE SKIN

#### §4.118 Schedule of ratings—skin.

	Rat- ing
7800 Disfigurement of the	head face or neck:

	Rat- ing
With visible or palpable tissue loss and either gross distortion or asymmetry of three or more features or paired sets of features (nose, chin, forehead, eyes (including eyelids), ears (auricles), cheeks, lips), or; with six or more characteristics of	
disfigurement With visible or palpable tissue loss and either gross distortion or asymmetry of two features or paired sets of features (nose, chin, forehead, eyes (including eyelids), ears (auricles), cheeks, lips), or; with four	80
or five characteristics of disfigurement With visible or palpable tissue loss and either gross distortion or asymmetry of one feature or paired set of features (nose, chin, forehead, eyes (including eyelids), ears (auricles), cheeks, lips), or; with two	50
or three characteristics of disfigurement With one characteristic of disfigurement Note (1):The 8 characteristics of disfigurement, for purposes of evaluation under § 4.118, are: Scar 5 or more inches (13 or more cm.) in length.	30 10
Scar at least one-quarter inch (0.6 cm.) wide at widest part.  Surface contour of scar elevated or depressed on palpation.  Scar adherent to underlying tissue.  Skin hypo-or hyper-pigmented in an area exceeding six square inches (39 sq. cm.).  Skin texture abnormal (irregular, atrophic, shiny, scaly, etc.) in an area exceeding six square inches (39 sq. cm.).  Underlying soft tissue missing in an area exceeding six square inches (39 sq. cm.).  Skin indurated and inflexible in an area exceeding six square inches (39 sq. cm.).  Note (2): Rate tissue loss of the auricle under DC 6207 (loss of auricle) and anatomical loss of the eye under DC 6061 (anatomical loss of both eyes) or DC 6063 (anatomical loss of one eye), as appropriate.  Note (3): Take into consideration unretouched color photographs when evaluating under these criteria.  7801 Scars, other than head, face, or neck, that are deep or that cause limited motion:	
Area or areas exceeding 144 square inches (929 sq.cm.)	40
(465 sq. cm.)	30
(77 sq. cm.)	20
Note (1): Scars in widely separated areas, as on two or more extremities or on anterior and posterior surfaces of extremities or trunk, will be separately rated and combined in accordance with §4.25 of this part.  Note (2): A deep scar is one associated with underlying soft tissue damage.  7802 Scars, other than head, face, or neck, that are superficial and that do not cause limited motion:	10

Area or areas of 144 square inches (929 sq.

cm.) or greater ......

		Rat-		Ŗat-
	Note (1): Score in widely concreted areas	ing	Note: Evaluate per eutopoque (viceoral)	ing
	<b>Note (1):</b> Scars in widely separated areas, as on two or more extremities or on anterior and posterior surfaces of extremities		Note: Evaluate non-cutaneous (visceral) leishmaniasis under DC 6301 (visceral leishmaniasis).	
	or trunk, will be separately rated and combined in accordance with § 4.25 of this		7809 Discoid lupus erythematosus or subacute cutaneous lupus erythematosus:	
	part.  Note (2): A superficial scar is one not asso-		Rate as disfigurement of the head, face, or neck (DC 7800), scars (DC's 7801, 7802,	
U3 (	ciated with underlying soft tissue damage. Scars, superficial, unstable	10	7803, 7804, or 7805), or dermatitis (DC 7806), depending upon the predominant	
05 (	Note (1): An unstable scar is one where, for	10	disability. Do not combine with ratings	
	any reason, there is frequent loss of covering of skin over the scar.		under DC 6350. 7811 Tuberculosis luposa (lupus vulgaris), active or	
	Note (2): A superficial scar is one not asso-		inactive:	
04 5	ciated with underlying soft tissue damage. Scars, superficial, painful on examination	10	Rate under §§ 4.88c or 4.89, whichever is appropriate.	
• • •	Note (1): A superficial scar is one not asso-		7813 Dermatophytosis (ringworm: of body, tinea	
	ciated with underlying soft tissue damage.  Note (2): In this case, a 10-percent evalua-		corporis; of head, tinea capitis; of feet, tinea pedis; of beard area, tinea barbae; of nails, tinea	
	tion will be assigned for a scar on the tip		unguium; of inguinal area (jock itch), tinea cruris):	
	of a finger or toe even though amputation of the part would not warrant a compen-		Rate as disfigurement of the head, face, or neck (DC 7800), scars (DC's 7801, 7802,	
	sable evaluation.		7803, 7804, or 7805), or dermatitis (DC	
	(See § 4.68 of this part on the amputation rule.)		7806), depending upon the predominant disability.	
	Scars, other; Rate on limitation of function of		7815 Bullous disorders (including pemphigus	
	ed part. Dermatitis or eczema.		vulgaris, pemphigus foliaceous, bullous pemphigoid, dermatitis herpetiformis, epidermolysis	
	More than 40 percent of the entire body or		bullosa acquisita, benign chronic familial	
	more than 40 percent of exposed areas affected, or; constant or near-constant		pemphigus (Hailey-Hailey), and porphyria cutanea tarda):	
	systemic therapy such as corticosteroids		More than 40 percent of the entire body or	
	or other immunosuppressive drugs required during the past 12-month period	60	more than 40 percent of exposed areas affected, or; constant or near-constant	
	20 to 40 percent of the entire body or 20 to		systemic therapy such as corticosteroids	
	40 percent of exposed areas affected, or; systemic therapy such as corticosteroids		or other immunosuppressive drugs required during the past 12-month period	
	or other immunosuppressive drugs re- quired for a total duration of six weeks or		20 to 40 percent of the entire body or 20 to	
	more, but not constantly, during the past		40 percent of exposed areas affected, or; systemic therapy such as corticosteroids	
	12-month period	30	or other immunosuppressive drugs required for a total duration of six weeks or	
	of the entire body, or at least 5 percent,		more, but not constantly, during the past	
	but less than 20 percent, of exposed areas affected, or; intermittent systemic		12-month period	;
	therapy such as corticosteroids or other		of the entire body, or at least 5 percent,	
	immunosuppressive drugs required for a total duration of less than six weeks dur-		but less than 20 percent, of exposed areas affected, or; intermittent systemic	
	ing the past 12-month period	10	therapy such as corticosteroids or other	
	Less than 5 percent of the entire body or less than 5 percent of exposed areas af-		immunosuppressive drugs required for a total duration of less than six weeks dur-	
	fected, and; no more than topical therapy	_	ing the past 12-month period	
	required during the past 12-month period Or rate as disfigurement of the head, face,	0	Less than 5 percent of the entire body or exposed areas affected, and; no more	
	or neck (DC 7800) or scars (DC's 7801,		than topical therapy required during the	
	7802, 7803, 7804, or 7805), depending upon the predominant disability.		past 12-month period	
	American (New World) leishmaniasis ocutaneous, espundia):		or neck (DC 7800) or scars (DC's 7801,	
mucc	Rate as disfigurement of the head, face, or		7802, 7803, 7804, or 7805), depending upon the predominant disability.	
	neck (DC 7800), scars (DC's 7801, 7802, 7803, 7804, or 7805), or dermatitis (DC		7816 Psoriasis:  More than 40 percent of the entire body or	
	7806), depending upon the predominant		more than 40 percent of exposed areas	
	disability.  Note: Evaluate non-cutaneous (visceral)		affected, or; constant or near-constant systemic therapy such as corticosteroids	
	leishmaniasis under DC 6301 (visceral		or other immunosuppressive drugs re-	
08 (	leishmaniasis). Old World leishmaniasis (cutaneous, Oriental		quired during the past 12-month period 20 to 40 percent of the entire body or 20 to	(
sore):	•		40 percent of exposed areas affected, or;	
	Rate as disfigurement of the head, face, or neck (DC 7800), scars (DC's, 7801, 7802,		systemic therapy such as corticosteroids or other immunosuppressive drugs re-	
	7803, 7804, or 7805), or dermatitis (DC		quired for a total duration of six weeks or	
	7806), depending upon the predominant disability.		more, but not constantly, during the past 12-month period	3

	Rat- ing		Rat- ing
At least 5 percent, but less than 20 percent, of the entire body, or at least 5 percent, but less than 20 percent, of exposed areas affected, or; intermittent systemic therapy such as corticosteroids or other immunosuppressive drugs required for a total duration of less than six weeks during the past 12-month period	100 0 100 100 0 10	Note: If a skin malignancy requires therapy that is comparable to that used for systemic malignancies, i.e., systemic chemotherapy, X-ray therapy more extensive than to the skin, or surgery more extensive than to the skin, or surgery more extensive than to the skin, or surgery more extensive than wide local excision, a 100-percent evaluation will be assigned from the date of onset of treatment, and will continue, with a mandatory VA examination six months following the completion of such antineoplastic treatment, and any change in evaluation based upon that or any subsequent examination will be subject to the provisions of § 3.105(e) of this chapter. If there has been no local recurrence or metastasis, evaluation will then be made on residuals. If treatment is confined to the skin, the provisions for a 100-percent evaluation do not apply.  7819 Benign skin neoplasms:  Rate as disfigurement of the head, face, or neck (DC 7800), scars (DC's 7801, 7802, 7803, 7804, or 7805), or impairment of function.  7820 Infections of the skin not listed elsewhere (including bacterial, fungal, viral, treponemal and parasitic diseases):  Rate as disfigurement of the head, face, or neck (DC 7800), scars (DC's 7801, 7802, 7803, 7804, or 7805), or dermatitis (DC 7806), depending upon the predominant disability.  7821 Cutaneous manifestations of collagen-vascular diseases not listed elsewhere (including scleroderma, calcinosis cutis, and dermatomyositis):  More than 40 percent of the entire body or more than 40 percent of exposed areas affected, or; constant or near-constant systemic therapy such as corticosteroids or other immunosuppressive drugs required during the past 12-month period  20 to 40 percent of the entire body or 20 to 40 percent of exposed areas affected, or; intermittent systemic therapy such as corticosteroids or other immunosuppressive drugs required for a total duration of exposed areas affected, or; intermittent systemic therapy such as corticosteroids or other immunosuppressive drugs required for a total du	60 30
		where (including lichen planus, large or small plaque parapsoriasis, pityriasis lichenoides et varioliformis acuta (PLEVA), lymphomatoid papulosus, and pityriasis rubra pilaris (PRP)):	

			1	
	Rat- ing			Rat- ing
More than 40 percent of the entire body or more than 40 percent of exposed areas affected, and; constant or near-constant systemic medications or intensive light			Recurrent debilitating episodes occurring at least four times during the past 12-month period, and; requiring intermittent systemic immunosuppressive therapy for	00
therapy required during the past 12-month period	60		Recurrent episodes occurring one to three times during the past 12-month period,	30
40 percent of exposed areas affected, or; systemic therapy or intensive light therapy required for a total duration of six weeks			and; requiring intermittent systemic immunosuppressive therapy for control  Or rate as disfigurement of the head, face,	10
or more, but not constantly, during the past 12-month period	30	7827	or neck (DC 7800) or scars (DC's 7801, 7802, 7803, 7804, or 7805), depending upon the predominant disability. Erythema multiforme; Toxic epidermal	
but less than 20 percent, of exposed areas affected, or; systemic therapy or intensive light therapy required for a total			lysis:  Recurrent debilitating episodes occurring at least four times during the past 12-month	
duration of less than six weeks during the past 12-month period	10		period despite ongoing immuno- suppressive therapy	60
exposed areas affected, and; no more than topical therapy required during the past 12-month period	0		times during the past 12-month period, and; requiring intermittent systemic im- munosuppressive therapy Recurrent episodes occurring during the	30
Or rate as disfigurement of the head, face, or neck (DC 7800) or scars (DC's 7801, 7802, 7803, 7804, or 7805), depending upon the predominant disability.  7823 Vitiligo:			past 12-month period that respond to treatment with antihistamines or sympathomimetics, or; one to three epi- sodes occurring during the past 12-month	
With exposed areas affected	10 0		period requiring intermittent systemic im- munosuppressive therapy	10
7824 Diseases of keratinization (including icthyoses, Darier's disease, and palmoplantar keratoderma): With either generalized cutaneous involve- ment or systemic manifestations, and;			Or rate as disfigurement of the head, face, or neck (DC 7800) or scars (DC's 7801, 7802, 7803, 7804, or 7805), depending upon the predominant disability.	
constant or near-constant systemic medi- cation, such as immunosuppressive retinoids, required during the past 12-	60	7828	Acne:  Deep acne (deep inflamed nodules and pusfilled cysts) affecting 40 percent or more of the face and neck	30
month period	60		Deep acne (deep inflamed nodules and pus- filled cysts) affecting less than 40 percent of the face and neck, or; deep acne other	10
immunosuppressive retinoids, required for a total duration of six weeks or more, but not constantly, during the past 12-month	20		than on the face and neck Superficial acne (comedones, papules, pustules, superficial cysts) of any extent Or rate as disfigurement of the head, face,	0
period	30	7829	or neck (DC 7800) or scars (DC's 7801, 7802, 7803, 7804, or 7805), depending upon the predominant disability. Chloracne:	
less than six weeks during the past 12- month period	10		Deep acne (deep inflamed nodules and pus- filled cysts) affecting 40 percent or more of the face and neck	30
ing the past 12-month period	0		Deep acne (deep inflamed nodules and pus- filled cysts) affecting less than 40 percent of the face and neck, or; deep acne other than on the face and neck	10
least four times during the past 12-month period despite continuous immuno-suppressive therapy	60		Superficial acne (comedones, papules, pustules, superficial cysts) of any extent Or rate as disfigurement of the head, face,	0
Recurrent debilitating episodes occurring at least four times during the past 12-month period, and; requiring intermittent sys- temic immunosuppressive therapy for		7830	or neck (DC 7800) or scars (DC's 7801, 7802, 7803, 7804, or 7805), depending upon the predominant disability.  Scarring alopecia:	
control	30		Affecting more than 40 percent of the scalp Affecting 20 to 40 percent of the scalp Affecting less than 20 percent of the scalp	20 10 0
and; responding to treatment with antihistamines or sympathomimetics	10		Alopecia areata:  With loss of all body hair  With loss of hair limited to scalp and face	10 0
Recurrent debilitating episodes occurring at least four times during the past 12-month period despite continuous immuno-		7832	Hyperhidrosis:  Unable to handle paper or tools because of moisture, and unresponsive to therapy	30

		Rat- ing
7833	Malignant melanoma: Rate as scars (DC's 7801, 7802, 7803, 7804, or 7805), disfigurement of the head, face, or neck (DC 7800), or impairment of function (under the appropriate body system).  Note: If a skin malignancy requires therapy that is comparable to that used for systemic malignancies, i.e., systemic chemotherapy, X-ray therapy more extensive than to the skin, or surgery more extensive than to the skin, or surgery more extensive than wide local excision, a 100-percent evaluation will be assigned from the date of onset of treatment, and will continue, with a mandatory VA examination six months following the completion of such antineoplastic treatment, and any change in evaluation based upon that or any subsequent examination will be subject to the provisions of § 3.105(e). If there has been no local recurrence or metastasis, evaluation will then be made on residuals. If treatment is confined to the skin, the provisions for a 100-percent evaluation do not apply.	

(Authority: 38 U.S.C. 1155)

[67 FR 49596, July 31, 2002; 67 FR 58448, 58449, Sept. 16, 2002]

### THE ENDOCRINE SYSTEM

# $\$\,4.119$ Schedule of ratings—endocrine system.

	Rat- ing
7900 Hyperthyroidism	
Thyroid enlargement, tachycardia (more than 100 beats per minute), eye involvement, muscular weakness, loss of weight, and sympathetic nervous system, cardiovascular, or	
astrointestinal symptoms	100
Emotional instability, tachycardia, fatigability, and increased pulse pressure or blood pressure	60
Tachycardia, tremor, and increased pulse pres- sure or blood pressure	30
Tachycardia, which may be intermittent, and trem-	
or, or; continuous medication required for con- trol	10
NOTE (1): If disease of the heart is the predominant finding, evaluate as hyperthyroid heart disease (DC 7008) if doing so would result in a higher evaluation than using the criteria above.  NOTE (2): If ophthalmopathy is the sole finding, evaluate as field vision, impairment of (DC 6080); diplopia (DC 6090); or impairment of central visual acuity (DC 6061–6079).	
7901 Thyroid gland, toxic adenoma of Thyroid enlargement, tachycardia (more than 100 beats per minute), eye involvement, muscular weakness, loss of weight, and sympathetic nervous system, cardiovascular, or qastro-	
intestinal symptoms	100
Emotional instability, tachycardia, fatigability, and increased pulse pressure or blood pressure	60
Tachycardia, tremor, and increased pulse pres- sure or blood pressure	30

	9
Tachycardia, which may be intermittent, and trem- or, or; continuous medication required for con-	
trol	10
NOTE (1): If disease of the heart is the predomi-	
nant finding, evaluate as hyperthyroid heart dis-	
ease (DC 7008) if doing so would result in a	
higher evaluation than using the criteria above.	
NOTE (2): If ophthalmopathy is the sole finding, evaluate as field vision, impairment of (DC	
6080); diplopia (DC 6090); or impairment of	
central visual acuity (DC 6061-6079).	
7902 Thyroid gland, nontoxic adenoma of	
With disfigurement of the head or neck	20
Without disfigurement of the head or neck	0
NOTE: If there are symptoms due to pressure on	
adjacent organs such as the trachea, larynx, or	
esophagus, evaluate under the diagnostic code	
for disability of that organ, if doing so would re-	
sult in a higher evaluation than using this diag-	
nostic code.	
7903 Hypothyroidism	
Cold intolerance, muscular weakness, cardio-	
vascular involvement, mental disturbance (dementia, slowing of thought, depression),	
bradycardia (less than 60 beats per minute),	
and sleepiness	100
Muscular weakness, mental disturbance, and	100
weight gain	60
Fatigability, constipation, and mental sluggishness	30
Fatigability, or; continuous medication required for	1
control	10
7904 Hyperparathyroidism	
Generalized decalcification of bones, kidney	
stones, gastrointestinal symptoms (nausea,	
vomiting, anorexia, constipation, weight loss, or	
peptic ulcer), and weakness	100
Gastrointestinal symptoms and weakness	60
Continuous medication required for control	10
Note: Following surgery or treatment, evaluate as	
digestive, skeletal, renal, or cardiovascular re-	
siduals or as endocrine dysfunction.	
7905 Hypoparathyroidism	
Marked neuromuscular excitability (such as con-	
vulsions, muscular spasms (tetany), or laryn-	
geal stridor) plus either cataract or evidence of	
increased intracranial pressure (such as papilledema)	100
Marked neuropusaular systemility	100
Marked neuromuscular excitability, or;	
paresthesias (of arms, legs, or circumoral area) plus either cataract or evidence of increased	
intracranial pressure	60
Continuous medication required for control	10
7907 Cushing's syndrome	10
As active, progressive disease including loss of	
muscle strength, areas of osteoporosis, hyper-	
tension, weakness, and enlargement of pituitary	
or adrenal gland	100
Loss of muscle strength and enlargement of pitui-	
tary or adrenal gland	60
With striae, obesity, moon face, glucose intoler-	
ance, and vascular fragility	30
NOTE: With recovery or control, evaluate as re-	
siduals of adrenal insufficiency or cardio-	
siduals of adrenal insufficiency or cardio- vascular, psychiatric, skin, or skeletal complica-	
tions under appropriate diagnostic code.	
7908 Acromegaly	
Evidence of increased intracranial pressure (such	
as visual field defect), arthropathy, glucose in- tolerance, and either hypertension or cardio-	
megaly	100
Arthropathy, glucose intolerance, and hyper-	00
tension	60

#### §4.120

	Rat- ing		Rat- ing
Enlargement of acral parts or overgrowth of long bones, and enlarged sella turcica	30 100 60 40 20 60 40	NOTE (1): Evaluate compensable complications of diabetes separately unless they are part of the criteria used to support a 100 percent evaluation. Noncompensable complications are considered part of the diabetic process under diagnostic code 7913.  NOTE (2): When diabetes mellitus has been conclusively diagnosed, do not request a glucose tolerance test solely for rating purposes.  7914 Neoplasm, malignant, any specified part of the endocrine system	100
required for control	20	als.  7915 Neoplasm, benign, any specified part of the endocrine system rate as residuals of endocrine dysfunction.  7916 Hyperpituitarism (prolactin secreting pituitary dysfunction)  7917 Hyperaldosteronism (benign or malignant)  7918 Pheochromocytoma (benign or malignant)  NOTE: Evaluate diagnostic codes 7916, 7917, and 7918 as malignant or benign neoplasm as appropriate.  7919 C-cell hyperplasia of the thyroid  NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.	100
Evaluate according to major manifestations.  7913 Diabetes mellitus		[61 FR 20446, May 7, 1996]	

7913 Diabetes mellitus

Requiring more than one daily injection of insulin, restricted diet, and regulation of activities (avoidance of strenuous occupational and recreational activities) with episodes of ketoacidosis or hypoglycemic reactions requiring at least three hospitalizations per year or weekly visits to a diabetic care provider, plus either progressive loss of weight and strength or complications that would be compensable if separately evaluated ...

Requiring insulin, restricted diet, and regulation of activities with episodes of ketoacidosis or hypoglycemic reactions requiring one or two hospitalizations per year or twice a month visits to a diabetic care provider, plus complications that would not be compensable if separately evaluated .

Requiring insulin, restricted diet, and regulation of

Requiring insulin and restricted diet, or; oral hypoglycemic agent and restricted diet ..... Manageable by restricted diet only .....

[61 FR 20446, May 7, 1996]

NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

#### § 4.120 Evaluations by comparison.

Disability in this field is ordinarily to be rated in proportion to the impairment of motor, sensory or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, injury to the skull, etc. In rating disability from the conditions in the preceding sentence refer to the appropriate schedule. In rating peripheral nerve injuries and their residuals, attention should be

100

60

40

20

given to the site and character of the injury, the relative impairment in motor function, trophic changes, or sensory disturbances.

#### §4.121 Identification of epilepsy.

When there is doubt as to the true nature of epileptiform attacks, neurological observation in a hospital adequate to make such a study is necessary. To warrant a rating for epilepsy, the seizures must be witnessed or verified at some time by a physician. As to frequency, competent, consistent lay testimony emphasizing convulsive and immediate post-convulsive characteristics may be accepted. The frequency of seizures should be ascertained under the ordinary conditions of life (while not hospitalized).

#### §4.122 Psychomotor epilepsy.

The term psychomotor epilepsy refers to a condition that is characterized by seizures and not uncommonly by a chronic psychiatric disturbance as well

(a) Psychomotor seizures consist of episodic alterations in conscious control that may be associated with automatic states, generalized convulsions, random motor movements (chewing, lip smacking, fumbling), hallucinatory phenomena (involving taste, smell, sound, vision), perceptual illusions (deja vu, feelings of loneliness, strangeness, macropsia, micropsia, dreamy states), alterations in thinking (not open to reason), alterations in memory, abnormalities of mood or affect (fear, alarm, terror, anger, dread, wellbeing), and autonomic disturbances (sweating, pallor, flushing of the face, visceral phenomena such as nausea, vomiting, defecation, a rising feeling of warmth in the abdomen). Automatic states or automatisms are characterized by episodes of irrational, irrelevant, disjointed, unconventional, asocial, purposeless though seemingly coordinated and purposeful, confused or inappropriate activity of one to several minutes (or, infrequently, hours) duration with subsequent amnesia for the seizure. Examples: A person of high social standing remained seated, muttered angrily, and rubbed the arms of his chair while the National Anthem was being played; an apparently normal person suddenly disrobed in public; a man traded an expensive automobile for an antiquated automobile in poor mechanical condition and after regaining conscious control, discovered that he had signed an agreement to pay an additional sum of money in the trade. The seizure manifestations of psychomotor epilepsy vary from patient to patient and in the same patient from seizure to seizure.

(b) A chronic mental disorder is not uncommon as an interseizure manifestation of psychomotor epilepsy and may include psychiatric disturbances extending from minimal anxiety to severe personality disorder (as distinguished from developmental) or almost complete personality disintegration (psychosis). The manifestations of a chronic mental disorder associated with psychomotor epilepsy, like those of the seizures, are protean in character.

#### §4.123 Neuritis, cranial or peripheral.

Neuritis, cranial or peripheral, characterized by loss of reflexes, muscle atrophy, sensory disturbances, and constant pain, at times excruciating, is to be rated on the scale provided for injury of the nerve involved, with a maximum equal to severe, incomplete, paralysis. See nerve involved for diagnostic code number and rating. The maximum rating which may be assigned for neuritis not characterized by organic changes referred to in this section will be that for moderate, or with sciatic nerve involvement, for moderately severe, incomplete paralysis.

## § 4.124 Neuralgia, cranial or peripheral.

Neuralgia, cranial or peripheral, characterized usually by a dull and intermittent pain, of typical distribution so as to identify the nerve, is to be rated on the same scale, with a maximum equal to moderate incomplete paralysis. See nerve involved for diagnostic code number and rating. Tic douloureux, or trifacial neuralgia, may be rated up to complete paralysis of the affected nerve.

Rat-

### §4.124a

#### §4.124a Schedule of ratings—neurological conditions and convulsive disorders.

[With the exceptions noted, disability from the following diseases and their residuals may be rated from 10 percent to 100 percent in proportion to the impairment of motor, sensory, or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, etc., referring to the appropriate bodily system of the schedule. With partial loss of use of one or more extremities from neurological lesions, rate by comparison with the mild, moderate, severe, or complete paralysis of peripheral nerves]

## ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM

	Rat- ing
8000 Encephalitis, epidemic, chronic:	
As active febrile disease	100
Rate residuals, minimum	10
Brain, new growth of:	
8002 Malignant	100
Note: The rating in code 8002 will be continued	
for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized,	
the rating will be made on neurological residu-	
als according to symptomatology.	
Minimum rating	30
8003 Benign, minimum	60
Rate residuals, minimum	10
8004 Paralysis agitans:	
Minimum rating	30
8005 Bulbar palsy	100
8007 Brain, vessels, embolism of.	
8008 Brain, vessels, thrombosis of.	
8009 Brain, vessels, hemorrhage from:	
Rate the vascular conditions under Codes 8007	
through 8009, for 6 months	100
Rate residuals, thereafter, minimum	10
8010 Myelitis:	
Minimum rating	10
8011 Poliomyelitis, anterior:	
As active febrile disease	100
Rate residuals, minimum	10
8012 Hematomyelia:	
For 6 months	100
Rate residuals, minimum	10
8013 Syphilis, cerebrospinal.	
8014 Syphilis, meningovascular.	
8015 Tabes dorsalis.	
NOTE: Rate upon the severity of convulsions, pa-	
ralysis, visual impairment or psychotic involve-	
ment, etc.	
8017 Amyotrophic lateral sclerosis:	
Minimum rating	30
8018 Multiple sclerosis:	
Minimum rating	30
8019 Meningitis, cerebrospinal, epidemic:	
As active febrile disease	100
Rate residuals, minimum	10
8020 Brain, abscess of:	40-
As active disease	100

## ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

	ing
Rate residuals, minimum	10
Spinal cord, new growths of:.	
8021 Malignant	100
NOTE: The rating in code 8021 will be continued	
for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized, the rating will be made on neurological residu-	
als according to symptomatology.	
Minimum rating	30
	60
0	10
Rate residuals, minimum	10
	20
Minimum rating	30
8024 Syringomyelia:	20
Minimum rating	30
8025 Myasthenia gravis:	
Minimum rating	30
NOTE: It is required for the minimum ratings for	
residuals under diagnostic codes 8000–8025, that there be ascertainable residuals. Deter-	
minations as to the presence of residuals not	
capable of objective verification, i.e., head-	
aches, dizziness, fatigability, must be ap-	
proached on the basis of the diagnosis re-	
corded; subjective residuals will be accepted	
when consistent with the disease and not	
more likely attributable to other disease or no	
disease. It is of exceptional importance that	
when ratings in excess of the prescribed min-	
imum ratings are assigned, the diagnostic codes utilized as bases of evaluation be cited,	
in addition to the codes identifying the diag-	
noses.	
8045 Brain disease due to trauma:	
Purely neurological disabilities, such as hemi-	
plegia, epileptiform seizures, facial nerve pa-	
ralysis, etc., following trauma to the brain, will	
be rated under the diagnostic codes specifi-	
cally dealing with such disabilities, with citation	
of a hyphenated diagnostic code (e.g., 8045-	
8207).	
Purely subjective complaints such as headache,	
dizziness, insomnia, etc., recognized as symp-	
tomatic of brain trauma, will be rated 10 per-	
cent and no more under diagnostic code 9304.	
This 10 percent rating will not be combined with any other rating for a disability due to	
brain trauma. Ratings in excess of 10 percent	
for brain disease due to trauma under diag-	
nostic code 9304 are not assignable in the ab-	
sence of a diagnosis of multi-infarct dementia	
associated with brain trauma.	
8046 Cerebral arteriosclerosis:	

## §4.124a

## **Department of Veterans Affairs**

# ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

	Rat- ing
Purely neurological disabilities, such as hemiplegia, cranial nerve paralysis, etc., due to cerebral arteriosclerosis will be rated under the diagnostic codes dealing with such specific disabilities, with citation of a hyphenated diagnostic code (e.g., 8046–8207).  Purely subjective complaints such as headache, dizziness, tinnitus, insomnia and irritability, recognized as symptomatic of a properly diagnosed cerebral arteriosclerosis, will be rated 10 percent and no more under diagnostic code 9305. This 10 percent rating will not be combined with any other rating for a disability due to cerebral or generalized arteriosclerosis. Ratings in excess of 10 percent for cerebral arteriosclerosis under diagnostic code 9305 are not assignable in the absence of a diagnosis of multi-infarct dementia with cerebral arteriosclerosis.  NOTE: The ratings under code 8046 apply only when the diagnosis of cerebral arteriosclerosis is substantiated by the entire clinical picture and not solely on findings of retinal arteriosclerosis	

### MISCELLANEOUS DISEASES

8100 Migraine:	
With very frequent completely prostrating and prolonged attacks productive of severe economic inadaptability	50
With characteristic prostrating attacks occurring on an average once a month over last several	
months	30
one in 2 months over last several months	10
With less frequent attacks	0
8103 Tic, convulsive:	
Severe	30
Moderate	10
Mild	0
NOTE: Depending upon frequency, severity, muscle groups involved.	
8104 Paramyoclonus multiplex (convulsive state,	
myoclonic type):	00
Rate as tic; convulsive; severe cases	60
Pronounced, progressive grave types	100
Severe	80
Moderately severe	50
Moderate	30
Mild	10
NOTE: Consider rheumatic etiology and com-	
plications.	
8106 Chorea, Huntington's.	
Rate as Sydenham's chorea. This, though a familial disease, has its onset in late adult life, and is considered a ratable disability.	
8107 Athetosis, acquired.	
Rate as chorea.	
8108 Narcolepsy.	
Rate as for epilepsy, petit mal.	

### DISEASES OF THE CRANIAL NERVES

	Rat- ing
Disability from lesions of peripheral portions of first, second, third, fourth, sixth, and eighth nerves will be rated under the Organs of Special Sense. The ratings for the cranial nerves are for unilateral involvement; when bilateral, combine but without the bilateral factor.	
Fifth (trigeminal) cranial nerve	
8205 Paralysis of:	
Complete	50
Incomplete, severe Incomplete, moderate	30
Note: Dependent upon relative degree of sen-	"
sory manifestation or motor loss.	
8305 Neuritis.	
8405 Neuralgia.	
NOTE: Tic douloureux may be rated in accord-	
ance with severity, up to complete paralysis.	
Seventh (facial) cranial nerve	
8207 Paralysis of:	30
Complete Incomplete, severe	20
Incomplete, moderate	10
Note: Dependent upon relative loss of innerva-	"
tion of facial muscles.	
8307 Neuritis.	
8407 Neuralgia.	
Ninth (glossopharyngeal) cranial nerve.	
8209 Paralysis of:	
Complete	30
Incomplete, severe Incomplete, moderate	10
Note: Dependent upon relative loss of ordinary	1
sensation in mucous membrane of the phar-	
ynx, fauces, and tonsils.	
8309 Neuritis.	
8409 Neuralgia.	
Tenth (pneumogastric, vagus) cranial nerve. 8210 Paralysis of:	
Complete	50
Incomplete, severe	30
Incomplete, moderate	10
NOTE: Dependent upon extent of sensory and	
motor loss to organs of voice, respiration,	
pharynx, stomach and heart.	
8310 Neuritis.	
8410 Neuralgia. Eleventh (spinal accessory, external branch) cra-	
nial nerve.	
8211 Paralysis of:	
Complete	30
Incomplete, severe	20
Incomplete, moderate	10
NOTE: Dependent upon loss of motor function of	
sternomastoid and trapezius muscles.	
8311 Neuritis. 8411 Neuralgia.	
Twelfth (hypoglossal) cranial nerve.	
8212 Paralysis of:	
Complete	50
Incomplete, severe	30
Incomplete, moderate	10
NOTE: Dependent upon loss of motor function of	
tongue.	
8312 Neuritis. 8412 Neuralgia.	

## §4.124a

### DISEASES OF THE PERIPHERAL NERVES

# DISEASES OF THE PERIPHERAL NERVES—Continued

	Rat	ina	Continued		
Schedule of ratings	Major	Minor	Schedule of ratings	Rat	ing
			ochedule of fattings	Major	Minor
The term "incomplete paralysis," with this and other peripheral nerve injuries, indi- cates a degree of lost or impaired func-			8613 Neuritis. 8713 Neuralgia.		
tion substantially less than the type pic- ture for complete paralysis given with			The musculospiral nerve (radial nerve)		
each nerve, whether due to varied level			8514 Paralysis of:		
of the nerve lesion or to partial regenera-			Complete; drop of hand and fingers,		
tion. When the involvement is wholly			wrist and fingers perpetually flexed,		
sensory, the rating should be for the mild, or at most, the moderate degree.			the thumb adducted falling within the line of the outer border of the index		
The ratings for the peripheral nerves are			finger; can not extend hand at wrist,		
for unilateral involvement; when bilateral,			extend proximal phalanges of fingers,		
combine with application of the bilateral factor.			extend thumb, or make lateral move- ment of wrist; supination of hand, ex-		
Upper radicular group (fifth and sixth			tension and flexion of elbow weak-		
cervicals)			ened, the loss of synergic motion of		
8510 Paralysis of:			extensors impairs the hand grip seri-		
Complete; all shoulder and elbow move-			ously; total paralysis of the triceps oc- curs only as the greatest rarity	70	60
ments lost or severely affected, hand			Incomplete:	,,,	00
and wrist movements not affected	70	60	Severe	50	40
Incomplete:	50	40	Moderate	30	20
Severe Moderate	50 40	40 30	Mild 8614 Neuritis.	20	20
Mild	20	20	8714 Neuralgia.		
8610 Neuritis.			NOTE: Lesions involving only "dissociat	ion of e	xtensor
8710 Neuralgia.			communis digitorum" and "paralysis be	low the e	xtensor
Middle radicular group			communis digitorum," will not exceed the ing under code 8514.	ne moder	ate rat-
8511 Paralysis of:			The median nerve		
Complete; adduction, abduction and ro-			8515 Paralysis of:		
tation of arm, flexion of elbow, and ex- tension of wrist lost or severely af-			Complete; the hand inclined to the ulnar		
fected	70	60	side, the index and middle fingers		
Incomplete:			more extended than normally, considerable atrophy of the muscles of the		
Severe	50	40	thenar eminence, the thumb in the		
Moderate	40 20	30 20	plane of the hand (ape hand);		
Mild	20	20	pronation incomplete and defective, absence of flexion of index finger and		
8711 Neuralgia.			feeble flexion of middle finger, cannot		
			make a fist, index and middle fingers		
Lower radicular group			remain extended; cannot flex distal phalanx of thumb, defective opposition		
8512 Paralysis of: Complete; all intrinsic muscles of hand,			and abduction of the thumb, at right		
and some or all of flexors of wrist and			angles to palm; flexion of wrist weak-		
fingers, paralyzed (substantial loss of			ened; pain with trophic disturbances	70	60
use of hand)	70	60	Incomplete: Severe	50	40
Incomplete: Severe	50	40	Moderate	30	20
Moderate	40	30	Mild	10	10
Mild	20	20	8615 Neuritis.		
8612 Neuritis.			8715 Neuralgia.		
8712 Neuralgia.			The ulnar nerve		
All radicular groups			8516 Paralysis of:		
8513 Paralysis of:			Complete; the "griffin claw" deformity, due to flexor contraction of ring and lit-		
Complete	90	80	tle fingers, atrophy very marked in dor-		
Incomplete:			sal interspace and thenar and		
Severe	70	60	hypothenar eminences; loss of extension of ring and little fingers cannot		
Moderate	40	30	sion of ring and little fingers cannot spread the fingers (or reverse), cannot		
Mild	20	20	adduct the thumb; flexion of wrist		
			weakened	60	50
			Incomplete: Severe	40	30
			Moderate	30	20
			Mild	10	10

## DISEASES OF THE PERIPHERAL NERVES—Continued

Schodula of ratings	Rating	
Schedule of ratings	Major	Minor
8616 Neuritis.		
8716 Neuralgia.		
Musculocutaneous nerve		
8517 Paralysis of:		
Complete; weakness but not loss of flex- ion of elbow and supination of forearm Incomplete:	30	20
Severe	20	20
Moderate	10	10
Mild	0	0
8617 Neuritis.		
8717 Neuralgia.		
Circumflex nerve		
8518 Paralysis of: Complete; abduction of arm is impos- sible, outward rotation is weakened; muscles supplied are deltoid and teres		
minor Incomplete:	50	40
Severe	30	20
Moderate	10	10
Mild 8618 Neuritis. 8718 Neuralgia.	0	0
Long thoracic nerve		
8519 Paralysis of: Complete; inability to raise arm above shoulder level, winged scapula de-		
formity	30	20
Severe	20	20
ModerateMild	10 0	10
NOTE: Not to be combined with lost motion		

NOTE: Not to be combined with lost motion above shoulder level.

8619 Neuritis. 8719 Neuralgia.

NOTE: Combined nerve injuries should be rated by reference to the major involvement, or if sufficient in extent, consider radicular group ratings.

	Rating
Sciatic nerve	
8520 Paralysis of:	
Complete; the foot dangles and drops, no active movement possible of mus- cles below the knee, flexion of knee	
weakened or (very rarely) lost	80
Incomplete:	
Severe, with marked muscular atrophy	60
Moderately severe	40
Moderate	20
Mild	10

3,	4. 124U
	Rating
8620 Neuritis. 8720 Neuralgia.	
External popliteal nerve (common peroneal)	
8521 Paralysis of:  Complete; foot drop and slight droop of first phalanges of all toes, cannot dorsiflex the foot, extension (dorsal flexion) of proximal phalanges of toes lost; abduction of foot lost, adduction weakened; anesthesia covers entire dorsum of foot and toes	40
Incomplete: Severe Moderate	30 20
Mild	10
Musculocutaneous nerve (superficial peroneal)	
8522 Paralysis of: Complete; eversion of foot weakened	30
Incomplete: Severe	20 10 0
8622 Neuritis. 8722 Neuralgia.	
Anterior tibial nerve (deep peroneal)	
8523 Paralysis of: Complete; dorsal flexion of foot lost	30
Incomplete: Severe Moderate Mild	20 10 0
8623 Neuritis. 8723 Neuralgia.	
Internal popliteal nerve (tibial)	
8524 Paralysis of: Complete; plantar flexion lost, frank adduction of foot impossible, flexion and separation of toes abolished; no muscle in sole can move; in lesions of the nerve high in popliteal fossa, plantar flexion of foot is lost	40
Severe	30 20 10
Posterior tibial nerve	
8525 Paralysis of:  Complete; paralysis of all muscles of sole of foot, frequently with painful paralysis of a causalgic nature; toes cannot be flexed; adduction is weakened; plantar flexion is impaired	30
Incomplete: Severe	20
OCACIC	20

#### §4.124a

### 38 CFR Ch. I (7-1-07 Edition)

THE EPILEPSIES—Continued

	Rating
Moderate	10
Mild	"
8725 Neuralgia.	
Anterior crural nerve (femoral)	
8526 Paralysis of:	
Complete; paralysis of quadriceps extensor muscles	40
Incomplete:	40
Severe	30
Moderate	20
Mild	10
8626 Neuritis. 8726 Neuralgia.	
•	
Internal saphenous nerve	
8527 Paralysis of: Severe to complete	10
Mild to moderate	
8627 Neuritis.	
8727 Neuralgia.	
Obturator nerve	
8528 Paralysis of:	
Severe to complete	10
Mild or moderate	0
8628 Neuritis. 8728 Neuralgia.	
External cutaneous nerve of thigh 8529 Paralysis of:	
Severe to complete	10
Mild or moderate	C
8629 Neuritis.	
8729 Neuralgia.	
Ilio-inguinal nerve	
8530 Paralysis of:	
Severe to complete	10
Mild or moderate8630 Neuritis.	C
8730 Neuralgia.	
8540 Soft-tissue sarcoma (of neurogenic	
origin)	100
NOTE: The 100 percent rating will be co	ntinued

NOTE: The 100 percent rating will be continued for 6 months following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. At this point, if there has been no local recurrence or metastases, the rating will be made on residuals.

#### THE EPILEPSIES

	Rat- ing
A thorough study of all material in §§ 4.121 and 4.122 of the preface and under the ratings for epilepsy is necessary prior to any rating action.  8910 Epilepsy, grand mal. Rate under the general rating formula for major seizures.  8911 Epilepsy, petit mal.	

	Rat- ing
Rate under the general rating formula for minor seizures.	
NOTE (1): A major seizure is characterized by the generalized tonic-clonic convulsion with unconsciousness.	
NOTE (2): A minor seizure consists of a brief interruption in consciousness or conscious control associated with staring or rhythmic blinking of the eyes or nodding of the head ("pure" petit mal), or sudden jerking movements of the arms, trunk, or head (myoclonic type) or sudden loss of postural control (akinetic type).	
General Rating Formula for Major and Minor Epileptic Seizures:	
Averaging at least 1 major seizure per month over the last year	100
months over the last year; or more than 10 minor seizures weekly	80
months over the last year; or 9–10 minor seizures per week	60
or 2 in the last year; or averaging at least 5 to 8 minor seizures weekly	40
or at least 2 minor seizures in the last 6 months	20
A confirmed diagnosis of epilepsy with a history of seizures	10
seizures, rate the predominating type. NOTE (3): There will be no distinction between diurnal and nocturnal major seizures. 8912 Epilepsy, Jacksonian and focal motor or sen- sory.	
<ul> <li>8913 Epilepsy, diencephalic.</li> <li>Rate as minor seizures, except in the presence of major and minor seizures, rate the predominating type.</li> <li>8914 Epilepsy, psychomotor.</li> </ul>	
Major seizures:  Psychomotor seizures will be rated as major seizures under the general rating formula when characterized by automatic states and/or generalized convulsions with unconsciousness.	
Minor seizures:  Psychomotor seizures will be rated as minor seizures under the general rating formula when characterized by brief transient episodes of random motor movements, hallucinations, perceptual illusions, abnormalities of thinking, memory or mood, or autonomic disturbances.	

Mental Disorders in Epilepsies: A nonpsychotic organic brain syndrome will be rated separately under the appropriate diagnostic code (e.g., 9304 or 9326). In the absence of a diagnosis of non-psychotic organic psychiatric disturbance (psychotic, psychoneurotic or personality disorder) if diagnosed and shown to be secondary to or directly associated with epilepsy will be rated separately. The psychotic or psychroneurotic disorder will be rated under the appropriate diagnostic code. The personality disorder will be rated as a dementia (e.g., diagnostic code 9304 or 9326).

Epilepsy and Unemployability: (1) Rating specialists must bear in mind that the epileptic, although his or her seizures are controlled, may find employment and rehabilitation difficult of attainment due to employer reluctance to the hiring of the

cylieptic.

(2) Where a case is encountered with a definite history of unemployment, full and complete development should be undertaken to ascertain whether the epilepsy is the determining factor in his or her inability to obtain employment.

(3) The assent of the claimant should first be obtained for

(3) The assent of the california more many permission to conduct this economic and social survey. The purpose of this survey is to secure all the relevant facts and data necessary to permit of a true judgment as to the reason for his or her unemployment and should include information on the california many permits.

(a) Education;
(b) Occupations prior and subsequent to service;
(c) Places of employment and reasons for termination;
(d) Wages received;
(e) Number of seizures.
(4) Upon completion of this survey and current examination, the case should have rating board consideration. Where in the judgment of the rating board the veteran's unemployability is due to epilepsy and jurisdiction is not vested in that body by reason of schedular evaluations, the case should be submitted to the Director, Compensation and Pension Service.

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 40 FR 42540, Sept. 15, 1975; 41 FR 11302, Mar. 18, 1976; 43 FR 45362, Oct. 2, 1978; 54 FR 4282, Jan. 30, 1989; 54 FR 49755, Dec. 1, 1989; 55 FR 154, Jan. 3, 1990; 56 FR 51653, Oct. 15, 1991; 57 FR 24364, June 9, 1992; 70 FR 75399, Dec. 20, 2005]

#### MENTAL DISORDERS

### $\S 4.125$ Diagnosis of mental disorders.

(a) If the diagnosis of a mental disorder does not conform to DSM-IV or is not supported by the findings on the examination report, the rating agency shall return the report to the examiner to substantiate the diagnosis.

(b) If the diagnosis of a mental disorder is changed, the rating agency shall determine whether the new diagnosis represents progression of the prior diagnosis, correction of an error in the prior diagnosis, or development of a new and separate condition. If it is not clear from the available records what the change of diagnosis represents, the rating agency shall return the report to the examiner for a determination.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

#### §4.126 Evaluation of disability from mental disorders.

(a) When evaluating a mental disorder, the rating agency shall consider the frequency, severity, and duration of psychiatric symptoms, the length of remissions, and the veteran's capacity for adjustment during periods of remission. The rating agency shall assign an

evaluation based on all the evidence of record that bears on occupational and social impairment rather than solely on the examiner's assessment of the level of disability at the moment of the examination.

(b) When evaluating the level of disability from a mental disorder, the rating agency will consider the extent of social impairment, but shall not assign an evaluation solely on the basis of social impairment.

(c) Delirium, dementia, and amnestic and other cognitive disorders shall be evaluated under the general rating formula for mental disorders; neurologic deficits or other impairments stemming from the same etiology (e.g., a head injury) shall be evaluated separately and combined with the evaluation for delirium, dementia, or amnestic or other cognitive disorder (see § 4.25).

(d) When a single disability has been diagnosed both as a physical condition and as a mental disorder, the rating agency shall evaluate it using a diagnostic code which represents the dominant (more disabling) aspect of the condition (see § 4.14).

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

#### §4.127 Mental retardation and personality disorders.

Mental retardation and personality disorders are not diseases or injuries for compensation purposes, and, except as provided in §3.310(a) of this chapter, disability resulting from them may not be service-connected. However, disability resulting from a mental disorder that is superimposed upon mental retardation or a personality disorder may be service-connected.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

#### § 4.128 Convalescence ratings lowing extended hospitalization.

If a mental disorder has been assigned a total evaluation due to a continuous period of hospitalization lasting six months or more, the rating agency shall continue the total evaluation indefinitely and schedule a mandatory examination six months after the veteran is discharged or released to

#### §4.129

nonbed care. A change in evaluation based on that or any subsequent examination shall be subject to the provisions of  $\S 3.105(e)$  of this chapter.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

## §4.129 Mental disorders due to traumatic stress.

When a mental disorder that develops in service as a result of a highly stressful event is severe enough to bring about the veteran's release from active military service, the rating agency shall assign an evaluation of not less than 50 percent and schedule an examination within the six month period following the veteran's discharge to de-

termine whether a change in evaluation is warranted.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

## § 4.130 Schedule of ratings—mental disorders.

The nomenclature employed in this portion of the rating schedule is based upon the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, of the American Psychiatric Association (DSM-IV). Rating agencies must be thoroughly familiar with this manual to properly implement the directives in §4.125 through §4.129 and to apply the general rating formula for mental disorders in §4.130. The schedule for rating for mental disorders is set forth as follows:

		Rating
	Schizophrenia and Other Psychotic Disorders	
9201	Schizophrenia, disorganized type	
9202	Schizophrenia, catatonic type	
9203		
9204		
9205		
9208		
9210	.,	
9211	Schizoaffective disorder	
	Delirium, Dementia, and Amnestic and Other Cognitive Disorders	
9300	Delirium	
9301		
9304		
	Vascular dementia	
	Dementia of unknown etiology	
	Dementia of the Alzheimer's type	
	Dementia due to other neurologic or general medical conditions (endocrine disorders, metabolic disorders, k's disease, brain tumors, etc.) or that are substance-induced (drugs, alcohol, poisons)	
9327		
9321	Organic mental disorder, other (including personality change due to a general medical condition)	
	Anxiety Disorders	
9400		
9403		
	Obsessive compulsive disorder	
	Other and unspecified neurosis	
9411		
	Panic disorder and/or agoraphobia	
9413	Anxiety disorder, not otherwise specified	
	Dissociative Disorders	
9416	Dissociative amnesia; dissociative fugue; dissociative identity disorder (multiple personality disorder)	
9417	Depersonalization disorder	
	Somatoform Disorders	
9421	Somatization disorder	
9422		
9423	Undifferentiated somatoform disorder	
9424	Conversion disorder	
9425	Hypochondriasis	

## **Department of Veterans Affairs**

§4.130

		Rating
	Mood Disorders	
9433 9434	Bipolar disorder Dysthymic disorder Major depressive disorder	
9435		
	Chronic Adjustment Disorder	
9440	Chronic adjustment disorder  General Rating Formula for Mental Disorders:  Total occupational and social impairment, due to such symptoms as: gross impairment in thought processes or communication; persistent delusions or hallucinations; grossly inappropriate behavior; persistent danger of hurting self or others; intermittent inability to perform activities of daily living (including maintenance of minimal personal hygiene); disorientation to time or place; memory loss for names of close relatives, own occupation, or own name  Occupational and social impairment, with deficiencies in most areas, such as work, school, family relations, judgment, thinking, or mood, due to such symptoms as: suicidal ideation; obsessional rituals which interfere with routine activities; speech intermittently illogical, obscure, or irrelevant: near-continuous panic or depression affecting the ability to function independently.	100
	appropriately and effectively; impaired impulse control (such as unprovoked irritability with periods of violence); spatial disorientation; neglect of personal appearance and hygiene; difficulty in adapting to stressful circumstances (including work or a worklike setting); inability to establish and maintain effective relationships	70
	and long-term memory (e.g., retention of only highly learned material, forgetting to complete tasks); impaired judgment; impaired abstract thinking; disturbances of motivation and mood; difficulty in establishing and maintaining effective work and social relationships	50
	pressed mood, anxiety, suspiciousness, panic attacks (weekly or less often), chronic sleep impairment, mild memory loss (such as forgetting names, directions, recent events)	30 10
	interfere with occupational and social functioning or to require continuous medication	0
	Eating Disorders	
9520 9521	Anorexia nervosa Bulimia nervosa Rating Formula for Eating Disorders: Self-induced weight loss to less than 80 percent of expected minimum weight, with incapacitating episodes of at least six weeks total duration per year, and requiring hospitalization	
	more than twice a year for parenteral nutrition or tube feeding	100 60
	Self-induced weight loss to less than 85 percent of expected minimum weight with incapacitating episodes of more than two but less than six weeks total duration per year	30
	ing disorder and incapacitating episodes of up to two weeks total duration per year	10

NOTE: An incapacitating episode is a period during which bed rest and treatment by a physician are required.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

## §4.149

### DENTAL AND ORAL CONDITIONS

### §4.149 [Reserved]

## $\$\,4.150~$ Schedule of ratings—dental and oral conditions.

		Rat- ing
9900	Maxilla or mandible, chronic osteomyelitis or	
oste	eoradionecrosis of:	
	Rate as osteomyelitis, chronic under diag-	
	nostic code 5000.	
9901	Mandible, loss of, complete, between angles	100
9902	Mandible, loss of approximately one-half:	
	Involving temporomandibular articulation	50
	Not involving temporomandibular articulation	30
9903	Mandible, nonunion of:	
	Severe	30
	Moderate	10
	NOTE—Dependent upon degree of motion and relative loss of masticatory function.	
9904	Mandible, malunion of:	
9904	Severe displacement	20
	Moderate displacement	10
	Slight displacement	10
	Note—Dependent upon degree of motion	ľ
	and relative loss of masticatory function.	
9905		
tion		
	Inter-incisal range:	
	0 to 10 mm	40
	11 to 20 mm	30
	21 to 30 mm	20
	31 to 40 mm	10
	Range of lateral excursion:	
	0 to 4 mm	10
	Note—Ratings for limited inter-incisal move-	
	ment shall not be combined with ratings	
	for limited lateral excursion.	
9906	Ramus, loss of whole or part of:	
	Involving loss of temporomandibular articu-	
	lation	
	Bilateral	50
	Unilateral  Not involving loss of temporomandibular ar-	30
	ticulation	
	Bilateral	30
	Unilateral	20
9907		20
	nce of, not involving loss of continuity:	
otai	Bilateral	20
	Unilateral	10
9908	Condyloid process, loss of, one or both sides	30
9909	Coronoid process, loss of:	30
	Bilateral	20

		Rat
	Unilateral	
9911	1 Hard palate, loss of half or more:	
	Not replaceable by prosthesis	:
	Replaceable by prosthesis	
9912		
0012	Not replaceable by prosthesis	
	Replaceable by prosthesis	
9913		
bo	ody of maxilla or mandible without loss of con-	
	Where the lost masticatory surface cannot	
	be restored by suitable prosthesis:	
	Loss of all teeth	
	Loss of all upper teeth	
	Loss of all lower teeth	
	All upper and lower posterior teeth	
	missing	
	All upper and lower anterior teeth	
	missing	
	All upper anterior teeth missing	
	All lower anterior teeth missing	
	All upper and lower teeth on one	
	side missing	
	Where the loss of masticatory surface can	
	be restored by suitable prosthesis	
	Note—These ratings apply only to bone	
	loss through trauma or disease such as	
	osteomyelitis, and not to the loss of the	
	alveolar process as a result of periodontal	
	disease, since such loss is not considered	
	disabling.	
9914	S .	
3311	Not replaceable by prosthesis	1
	Replaceable by prostnesis	
9915		
2216	Loss of 25 to 50 percent:	
	Not replaceable by prosthesis	
	Replaceable by prosthesis	
	Loss of less than 25 percent:	
	Not replaceable by prosthesis	
0044	Replaceable by prosthesis	
9916		
	Severe displacement	
	Moderate displacement	
	Slight displacement	

[59 FR 2530, Jan. 18, 1994]

APPENDIX A TO PART 4—TABLE OF AMENDMENTS AND EFFECTIVE DATES SINCE 1946

Sec.	Diagnostic code No.	
4.71a	5000	Evaluation February 1, 1962.
	5001	Evaluation March 11, 1969.
	5002	Evaluation March 1, 1963.
	5003	Added July 6, 1950.
	5012	Criterion March 10, 1976.
	5024	Criterion March 1, 1963.
	5025	Added May 7, 1996.
	5051	Added September 22, 1978.
	5052	Added September 22, 1978.
	5053	Added September 22, 1978.
	5054	Added September 22, 1978.
	5055	Added September 22, 1978.
	5056	Added September 22, 1978.
	5100-5103	Removed March 10, 1976.
	5104	Criterion March 10, 1976.

Sec.	Diagnostic code No.	
	5105	Criterion March 10, 1976.
	5164	Evaluation June 9, 1952.
	5166	Criterion September 22, 1978.
	5172	Added July 6, 1950.
	5173	Added June 9, 1952.
	5174 5211	Added September 9, 1975; removed September 22, 1978.  Criterion September 22, 1978.
	5212	Criterion September 22, 1978.
	5214	Criterion September 22, 1978.
	5216	Preceding paragraph criterion September 22, 1978.
	5217	Criterion August 26, 2002.
	5218	Criterion August 26, 2002.
	5219 5220	Criterion September 22, 1978; criterion August 26, 2002.
	5223	Preceding paragraph criterion September 22, 1978; criterion August 26, 2002. Criterion August 26, 2002.
	5224	Criterion August 26, 2002.
	5225	Criterion August 26, 2002.
	5226	Criterion August 26, 2002.
	5227	Criterion September 22, 1978; criterion August 26, 2002.
	5228	Added August 26, 2002.
	5229 5230	Added August 26, 2002.
	5235-5243	Added August 26, 2002.   Replaces 5285–5295 September 26, 2003.
	5243	Criterion September 26, 2003.
	5255	Criterion July 6, 1950.
	5257	Evaluation July 6, 1950.
	5264	Added September 9, 1975; removed September 22, 1978.
	5275	Criterion March 10, 1976; criterion September 22, 1978.
	5285–5292 5293	Revised to 5235–5243 September 26, 2003.  Criterion March 10, 1976; criterion September 23, 2002; revised and moved
	3293	5235–5243 September 26, 2003.
	5294	Evaluation March 10, 1976; revised and moved to 5235–5243 September 2 2003.
	5295	Evaluation March 10, 1976; revised and moved to 5235–5243 September 2 2003.
	5296	Criterion March 10, 1976.
	5297	Criterion August 23, 1948; criterion February 1, 1962.
1.70	5298	Added August 23, 1948.
1.73	5317	Introduction NOTE criterion July 3, 1997.
	5324	Criterion September 22, 1978. Added February 1, 1962.
	5325	Criterion July 3, 1997.
	5327	Added March 10, 1976; criterion October 15, 1991; criterion July 3, 1997.
	5328	Added NOTE March 10, 1976.
	5329	Added NOTE July 3, 1997.
4.84a		Table V criterion July 1, 1994.
	6010	Criterion March 11, 1969.
	6019 6029	Criterion September 22, 1978.  NOTE August 23, 1948; criterion September 22, 1978.
	6035	Added September 9, 1975.
	6050-6062	Removed March 10, 1976.
	6061	Added March 10, 1976.
	6062	Added March 10, 1976.
	6063–6079	Criterion September 22, 1978.
	6064 6071	Criterion March 10, 1976.
	6076	Criterion March 10, 1976.  Evaluation August 23, 1948.
	6080	Criterion September 22, 1978.
	6081	Criterion March 10, 1976.
	6090	Criterion September 22, 1978; criterion September 12, 1988.
.84b	6260	Added October 1, 1961; criterion October 1, 1961; evaluation March 10, 197
		removed December 18, 1987; re-designated § 4.87a December 18, 1987.
.87		Tables VI and VII replaced by new Tables VI, VIA, and VII December 18, 198
070	6200 6200	6200–6260 revised and re-designated § 4.87 June 10, 1999.
.87a	6200–6260 6275–6276	Moved to § 4.87 June 10, 1999.  Moved from § 4.87b June 10, 1999.
	6277–6297	March 23, 1956 removed, December 17, 1987; Table II revised Table V Mar
	0211-0291	10, 1976; Table II revised to Table VII September 22, 1978; text from §4.8
		Schedule of ratings-ear re-designated from §4.87 December 17, 1987.
	6286	Removed December 17, 1987.
	6291	Criterion March 10, 1976; removed December 17, 1987.
	6297	Criterion March 10, 1976; removed December 17, 1987.

4.88a4.88b	6300	March 11, 1969; re-designated §4.88b November 29, 1994; §4.88a added to read "Chronic fatigue syndrome"; criterion November 29, 1994.  Added March 11, 1969; re-designated §4.88c November 29, 1994; §4.88a re-
4.88b		
4.886		Added March 11 1969; re-designated 6.4 88c November 29 1994; 6.4 88a re-
	6300	designated to § 4.88b November 29, 1994.
		Criterion August 30, 1996.
	6302	Criterion September 22, 1978; criterion August 30, 1996.
	6304 6305	Evaluation August 30, 1996. Criterion March 1, 1989; evaluation August 30, 1996.
	6306	Evaluation August 30, 1996.
	6307	Criterion August 30, 1996.
	6308 6309	Criterion August 30, 1996. Added March 1, 1963; criterion March 1, 1989; criterion August 30, 1996.
	6314	Evaluation March 1, 1989; evaluation August 30, 1996.
	6315	Criterion August 30, 1996.
	6316	Evaluation March 1, 1989; evaluation August 30, 1996.
	6317 6318	Criterion August 30, 1996. Added March 1, 1989; criterion August 30, 1996.
	6319	Added August 30, 1996.
	6320	Added August 30, 1996.
	6350	Evaluation March 1, 1963; evaluation March 10, 1976; evaluation August 30, 1996.
	6351	Added March 1, 1989; evaluation March 24, 1992; criterion August 30, 1996.
	6352	Added March 1, 1989; removed March 24, 1992.
	6353 6354	Added March 1, 1989; removed March 24, 1992.
4.88c	0334	Added November 29, 1994; criterion August 30, 1996.  Re-designated from § 4.88b November 29, 1994.
4.89		Ratings for nonpulmonary TB December 1, 1949; criterion March 11, 1969.
4.97	6502	Criterion October 7, 1996.
	6504 6510–6514	Criterion October 7, 1996. Criterion October 7, 1996.
	6515	Criterion March 11, 1969.
	6516	Criterion October 7, 1996.
	6517 6518	Removed October 7, 1996. Criterion October 7, 1996.
	6519	Criterion October 7, 1996.
	6520	Criterion October 7, 1996.
	6521	Added October 7, 1996.
	6522 6523	Added October 7, 1996. Added October 7, 1996.
	6524	Added October 7, 1996.
	6600	Evaluation September 9, 1975; criterion October 7, 1996.
	6601	Criterion October 7, 1996.
	6602 6603	Criterion September 9, 1975; criterion October 7, 1996.  Added September 9, 1975; criterion October 7, 1996.
	6604	Added October 7, 1996.
	6701	Evaluation October 7, 1996.
	6702 6703	Evaluation October 7, 1996. Evaluation October 7, 1996.
	6704	Subparagraph (1) following December 1, 1949; criterion March 11, 1969; criterion
		September 22, 1978.
	6705 6707–6710	Removed March 11, 1969. Added March 11, 1969; removed September 22, 1978.
	6721	Criterion July 6, 1950; criterion September 22, 1978.
	6724	Second note following December 1, 1949; criterion March 11, 1969; evaluation
	6705 6700	October 7, 1996.
	6725–6728 6730	Added March 11, 1969; removed September 22, 1978. Added September 22, 1978; criterion October 7, 1996.
	6731	Evaluation September 22, 1978; criterion October 7, 1996.
	6732	Criterion March 11, 1969.
	6800 6801	Criterion September 9, 1975; removed October 7, 1996.
	6801 6802	Removed October 7, 1996. Criterion September 9, 1975; removed October 7, 1996.
	6810-6813	Removed October 7, 1996.
	6814	Criterion March 10, 1976; removed October 7, 1996.
	6815 6816	Removed October 7, 1996. Removed October 7, 1996.
	6817	Evaluation October 7, 1996.
	6818	Removed October 7, 1996.
	6819	Criterion March 10, 1976; criterion October 7, 1996.
	6821 6822–6847	Evaluation August 23, 1948. Added October 7, 1996.
4.104	7000	Evaluation July 6, 1950; evaluation September 22, 1978; evaluation January 12,

Sec.	Diagnostic code No.	
	7001	Evaluation January 12, 1998.
	7002	Evaluation January 12, 1998.
	7003	Evaluation January 12, 1998.
	7004 7005	Criterion September 22, 1978; evaluation January 12, 1998.    Evaluation September 9, 1975; evaluation September 22, 1978; evaluation Janu
		ary 12, 1998.
	7006	Evaluation January 12, 1998.
	7007	Evaluation September 22, 1978; evaluation January 12, 1998.
	7008 7010	Evaluation January 12, 1998. Evaluation January 12, 1998.
	7011	Evaluation January 12, 1998.
	7013	Removed January 12, 1998.
	7014	Removed January 12, 1998.
	7015	Evaluation September 9, 1975; criterion January 12, 1998.
	7016 7017	Added September 9, 1975; evaluation January 12, 1998. Added September 22, 1978; evaluation January 12, 1998.
	7017	Added January 12, 1998.
	7019	Added January 12, 1998.
	7020	Added January 12, 1998.
	7100	Evaluation July 6, 1950.
	7101	Criterion September 1, 1960; criterion September 9, 1975; criterion January 1: 1998.
	7110	Evaluation September 9, 1975; evaluation January 12, 1998.
	7111 7112	Criterion September 9, 1975; evaluation January 12, 1998. Evaluation January 12, 1998.
	7112	Evaluation January 12, 1998.
	7114	Added June 9, 1952; evaluation January 12, 1998.
	7115	Added June 9, 1952; evaluation January 12, 1998.
	7116	Added June 9, 1952; evaluation March 10, 1976; removed January 12, 1998.
	7117	Added June 9, 1952; evaluation January 12, 1998.
	7118 7119	Criterion January 12, 1998. Evaluation January 12, 1998.
	7120	Note following July 6, 1950; evaluation January 12, 1998.
	7121	Criterion July 6, 1950; evaluation March 10, 1976; evaluation January 12, 1998.
	7122	Last sentence of Note following July 6, 1950; evaluation January 12, 1998; cr terion August 13, 1998.
	7123	Added October 15, 1991; criterion January 12, 1998.
4.114		Introduction paragraph revised March 10, 1976.
	7304 7305	Evaluation November 1, 1962.  Evaluation November 1, 1962.
	7308	Evaluation April 8, 1959.
	7311	Criterion July 2, 2001.
	7312	Evaluation March 10, 1976; evaluation July 2, 2001.
	7313	Evaluation March 10, 1976; removed July 2, 2001.
	7319	Evaluation November 1, 1962.
	7321 7328	Evaluation July 6, 1950; criterion March 10, 1976.  Evaluation November 1, 1962.
	7329	Evaluation November 1, 1962.
	7330	Evaluation November 1, 1962.
	7331	Criterion March 11, 1969.
	7332	Evaluation November 1, 1962.
	7334	Evaluation July 6, 1950; evaluation November 1, 1962.
	7339 7341	Criterion March 10, 1976. Removed March 10, 1976.
	7343	Criterion March 10, 1976; criterion July 2, 2001.
	7344	Criterion July 2, 2001.
	7345	Evaluation August 23, 1948; evaluation February 17, 1955; evaluation July 2001.
	7346	Evaluation February 1, 1962.
	7347	Added September 9, 1975.
	7348	Added March 10, 1976.
	7351 7354	Added July 2, 2001. Added July 2, 2001.
1.115a		Re-designated and revised as §4.115b; new §4.115a "Ratings of the genite urinary system-dysfunctions" added February 17, 1994.
I.115b	7500	Note July 6, 1950; evaluation February 17, 1994, criterion September 8, 1994.
	7501	Evaluation February 17, 1994.
	7502	Evaluation February 17, 1994.
	7503	Removed February 17, 1994.
	7504	Criterion February 17, 1994.
		Customer Storch 44 4000, evelvation Fahrman, 47 4004
	7505 7507	Criterion March 11, 1969; evaluation February 17, 1994. Criterion February 17, 1994.

Sec.	Diagnostic code No.	
	7509	Criterion February 17, 1994.
	7510	Evaluation February 17, 1994.
	7511	Evaluation February 17, 1994.
	7512	Evaluation February 17, 1994.
	7513 7514	Removed February 17, 1994. Criterion March 11, 1969; removed February 17, 1994.
	7515	Criterion February 17, 1994.
	7516	Criterion February 17, 1994.
	7517	Criterion February 17, 1994.
	7518	Evaluation February 17, 1994.
	7519	Evaluation March 10, 1976; evaluation February 17, 1994.
	7520 7521	Criterion February 17, 1994. Criterion February 17, 1994.
	7522	Criterion September 8, 1994.
	7523	Criterion September 8, 1994.
	7524	Note July 6, 1950; evaluation February 17, 1994; evaluation September 8, 1994
	7525	Criterion March 11, 1969; evaluation February 17, 1994.
	7526	Removed February 17, 1994.
	7527	Criterion February 17, 1994.
	7528 7529	Criterion March 10, 1976; criterion February 17, 1994. Criterion February 17, 1994.
	7530	Added September 9, 1975; evaluation February 17, 1994.
	7531	Added September 9, 1975; criterion February 17, 1994.
	7532-7542	Added February 17, 1994.
.116		§4.116 removed and §4.116a re-designated §4.116 "Schedule of ratings-gyn
	7040	cological conditions and disorders of the breasts" May 22, 1995.
	7610	Criterion May 22, 1995.
	7611 7612	Criterion May 22, 1995.  Criterion May 22, 1995.
	7613	Criterion May 22, 1995.
	7614	Criterion May 22, 1995.
	7615	Criterion May 22, 1995.
	7617	Criterion May 22, 1995.
	7618	Criterion May 22, 1995.
	7619	Criterion May 22, 1995.
	7620	Criterion May 22, 1995.
	7621	Criterion May 22, 1995.  Evaluation May 22, 1995.
	7622 7623	Evaluation May 22, 1995.
	7624	Criterion August 9, 1976; evaluation May 22, 1995.
	7625	Criterion August 9, 1976; evaluation May 22, 1995.
	7626	Criterion May 22, 1995; criterion March 18, 2002.
	7627	Criterion March 10, 1976; criterion May 22, 1995.
	7628	Added May 22, 1995.
117	7629	Added May 22, 1995.
.117	7700 7701	Evaluation October 23, 1995. Removed October 23, 1995.
	7702	Evaluation October 23, 1995.
	7703	Evaluation August 23, 1948; criterion October 23, 1995.
	7704	Evaluation October 23, 1995.
	7705	Evaluation October 23, 1995.
	7706	Evaluation October 23, 1995.
	7707	Criterion October 23, 1995.  Evaluation March 10, 1976; criterion October 23, 1995.
	7709 7710	Criterion October 23, 1995.
	7711	Criterion October 23, 1995.
	7712	Criterion October 23, 1995.
	7713	Removed October 23, 1995.
	7714	Added September 9, 1975; criterion October 23, 1995.
	7715	Added October 26, 1990.
110	7716	Added October 23, 1995.
118	7800 7801	Evaluation August 30, 2002. Criterion July 6, 1950; criterion August 30, 2002.
	7802	Criterion September 22, 1978; criterion August 30, 2002.
	7803	Criterion August 30, 2002.
	7804	Criterion July 6, 1950; criterion September 22, 1978; criterion August 30, 2002.
	7806	Criterion September 9, 1975; evaluation August 30, 2002.
	7807	Criterion August 30, 2002.
	7808	Criterion August 30, 2002.
	7809	Criterion August 30, 2002.
	7810	Removed August 30, 2002.
	7811	Criterion March 11, 1969; evaluation August 30, 2002.  Removed August 30, 2002.

Sec.	Diagnostic code No.	
	7813	Criterion August 30, 2002.
	7814	Removed August 30, 2002.
	7815	Evaluation August 30, 2002.
	7816	Evaluation August 30, 2002.
	7817 7818	Evaluation August 30, 2002. Criterion August 30, 2002.
	7819	Criterion August 30, 2002.
	7820-7833	Added August 30, 2002.
4.119	7900	Criterion August 13, 1981; evaluation June 9, 1996.
	7901	Criterion August 13, 1981; evaluation June 9, 1996.
	7902 7903	Evaluation August 13, 1981; criterion June 9, 1996. Criterion August 13, 1981; evaluation June 9, 1996.
	7904	Criterion August 13, 1981; evaluation June 9, 1996.
	7905	Evaluation; August 13, 1981; evaluation June 9, 1996.
	7907	Evaluation August 13, 1981; evaluation June 9, 1996.
	7908	Criterion August 13, 1981; criterion June 9, 1996.
	7909	Evaluation August 13, 1981; criterion June 9, 1996.
	7910 7911	Removed June 9, 1996. Evaluation March 11, 1969; evaluation August 13, 1981; criterion June 9, 1996.
	7913	Criterion September 9, 1975; criterion August 13, 1981; criterion June 6, 1996.
	7914	Criterion March 10, 1976; criterion August 13, 1981; criterion June 9, 1996.
	7916	Added June 9, 1996.
	7917	Added June 9, 1996.
	7918	Added June 9, 1996.
1 1240	7919 8002	Added June 9, 1996. Criterion September 22, 1978.
4.124a	8021	Criterion September 22, 1976.  Criterion September 22, 1978; criterion October 1, 1961; criterion March 10,
	002.	1976; criterion March 1, 1989.
	8046	Added October 1, 1961; criterion March 10, 1976; criterion March 1, 1989.
	8100	Evaluation June 9, 1953.
	8540	Added October 15, 1991.
	8910	Added October 1, 1961.
	8911 8912	Added October 1, 1961; evaluation September 9, 1975. Added October 1, 1961.
	8913	Added October 1, 1961.
	8914	Added October 1, 1961; criterion September 9, 1975; criterion March 10, 1976.
	8910-8914	Evaluations September 9, 1975.
4.125—4.132		All Diagnostic Codes under Mental Disorders October 1, 1961; except as to eval-
1 120		uation for Diagnostic Codes 9500 through 9511 September 9, 1975.
4.130	9200	Re-designated from § 4.132 November 7, 1996. Removed February 3, 1988.
	9201	Criterion February 3, 1988.
	9202	Criterion February 3, 1988.
	9203	Criterion February 3, 1988.
	9204	Criterion February 3, 1988.
	9205 9206	Criterion February 3, 1988; criterion November 7, 1996. Criterion February 3, 1988; removed November 7, 1996.
	9207	Criterion February 3, 1988; removed November 7, 1996.
	9208	Criterion February 3, 1988; removed November 7, 1996.
	9209	Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	2010	1996.
	9210 9211	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.  Added November 7, 1996.
	9300	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.
	9301	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.
	9302	Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
		1996.
	9303	Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9304	1996. Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.
	9305	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.
	9306	Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9307	1996. Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9308	1996. Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9309	1996. Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9310	1996. Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.
	9311	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9312	Added March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.

## Pt. 4, App. B

Sec.	Diagnostic code No.	
	9313	Added March 10, 1976; removed February 3, 1988.
	9314	Added March 10, 1976; removed February 3, 1988.
	9315	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9316–9321	Added March 10, 1976; removed February 3, 1988.
	9322	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9323	Added March 10, 1976; removed February 3, 1988.
	9324	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9325	Added March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9326	Added March 10, 1976; removed February 3, 1988; added November 7, 1996.
	9327	Added November 7, 1996.
	9400–9411	Evaluations February 3, 1988.
	9400 9401	Criterion March 10, 1976; criterion February 3, 1988.  Criterion March 10, 1976; criterion February 3, 1988; removed November 7,
	9401	1996.
	9402	Criterion March 10, 1976; criterion February 3, 1988; removed November 7, 1996.
	9403	Criterion March 10, 1976; criterion February 3, 1988; criterion November 7, 1996.
	9410	Added March 10, 1976; criterion February 3, 1988.
	9411	Added February 3, 1988.
	9412	Added November 7, 1996.
	9413 9416	Added November 7, 1996. Added November 7, 1996.
	9417	Added November 7, 1996.
	9421	Added November 7, 1996.
	9422	Added November 7, 1996.
	9423	Added November 7, 1996.
	9424	Added November 7, 1996.
	9425	Added November 7, 1996.
	9431	Added November 7, 1996.
	9432	Added November 7, 1996.
	9433	Added November 7, 1996.
	9434 9435	Added November 7, 1996. Added November 7, 1996.
	9440	Added November 7, 1996.
	9500	Criterion March 10, 1976; criterion February 3, 1988.
	9501	Criterion March 10, 1976; criterion February 3, 1988.
	9502	Criterion March 10, 1976; criterion February 3, 1988.
	9503	Removed March 10, 1976.
	9504	Criterion September 9, 1975; removed March 10, 1976.
	9505	Added March 10, 1976; criterion February 3, 1988.
	9506	Added March 10, 1976; criterion February 3, 1988.
	9507 9508	Added March 10, 1976; criterion February 3, 1988. Added March 10, 1976; criterion February 3, 1988.
	9509	Added March 10, 1976; criterion February 3, 1988.
	9510	Added March 10, 1976; criterion February 3, 1988.
	9511	Added March 10, 1976; criterion February 3, 1988.
	9520	Added November 7, 1996.
	9521	Added November 7, 1996.
4.132		Re-designated as § 4.130 November 7, 1996.
4.150	9900	Criterion September 22, 1978; criterion February 17, 1994.
	9901	Criterion February 17, 1994.
	9902	Criterion February 17, 1994.
	9903 9905	Criterion February 17, 1994. Criterion September 22, 1978; evaluation February 17, 1994.
	9910	Removed February 17, 1994.
	9913	Criterion February 17, 1994.
	9914	Added February 17, 1994.
	9915	Added February 17, 1994.
	9916	Added February 17, 1994.

[72 FR 12983, Mar. 20, 2007; 72 FR 16728, Apr. 5, 2007]

# APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES

Diagnostic Code No.		
THE MUSCULOSKELETAL SYSTEM Acute, Subacute, or Chronic Diseases		
5000 5001	Osteomyelitis, acute, subacute, or chronic. Bones and Joints, tuberculosis.	

Diagnostic Code No.	
5002	Arthritis, rheumatoid (atrophic).
5003	Arthritis, degenerative (hypertrophic or osteoarthritis).
5004	Arthritis, gonorrheal.
5005	Arthritis, pneumococcic.
5006	Arthritis, typhoid.
5007	Arthritis, syphilitic.
5008	Arthritis, streptococcic.
5009	Arthritis, other types (specify).
5010 5011	Arthritis, due to trauma.  Bones, caisson disease.
5012	Bones, new growths, malignant.
5013	Osteoporosis, with joint manifestations.
5014	Osteomalacia.
5015	Bones, new growths, benign.
5016	Osteitis deformans.
5017	Gout.
5018	Hydrarthrosis, intermittent.
5019 5020	Bursitis. Synovitis.
5021	Myositis.
5022	Periostitis.
5023	Myositis ossificans.
5024	Tenosynovitis.
5025	Fibromyalgia.
	Prosthetic Implants
5051	Shoulder replacement (prosthesis).
5052	Elbow replacement (prostness).
5053	Wrist replacement (prosthesis).
5054	Hip replacement (prosthesis).
5055	Knee replacement (prosthesis).
5056	Ankle replacement (prosthesis).
	Combination of Disabilities
5104	Anatomical loss of one hand and loss of use of one foot.
5105	Anatomical loss of one foot and loss of use of one hand.
5106	Anatomical loss of both hands.
5107	Anatomical loss of both feet.
5108	Anatomical loss of one hand and one foot.
5109	Loss of use of both hands.
5110 5111	Loss of use of one hand and are feet
3111	Loss of use of one hand and one foot.
	Amputations: Upper Extremity
Arm amputation of: 5120	Disarticulation.
5121	Above insertion of deltoid.
5122	Below insertion of deltoid.
Forearm amputation of:	<u>L</u>
5123	Above insertion of pronator teres.
5124	Below insertion of pronator teres.
- · - · · · · · · · · · · · · · · · · ·	
5125	Hand, loss of use of.
	Hand, loss of use of.  Multiple Finger Amputations
5126	Multiple Finger Amputations
5125	Multiple Finger Amputations  Five digits of one hand.
5126	Multiple Finger Amputations
5126	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, long and little. Thumb, index, ring and little.
5126	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, long and little. Thumb, index, ring and little. Thumb, long, ring and little.
5126	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, long and little. Thumb, index, ring and little.
5126	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, long and little. Thumb, index, ring and little. Thumb, long, ring and little.
5125	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, long and little. Thumb, index, ring and little. Thumb, long, ring and little. Index, long, ring and little. Index, long, ring and little.
5126	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, long and little. Thumb, long, ring and little. Thumb, long, ring and little. Index, long, ring and little. Thumb, index and long. Thumb, index and ring.
5126	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, ring and little. Thumb, long, ring and little. Thumb, long, ring and little. Index, long, ring and little. Index, long, ring and little.  Thumb, index and long. Thumb, index and ring. Thumb, index and little.
5125	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, long and little. Thumb, index, ring and little. Thumb, long, ring and little. Index, long, ring and little.  Thumb, index and long. Thumb, index and ring. Thumb, index and little. Thumb, index and ring. Thumb, long and ring.
5126	Multiple Finger Amputations  Five digits of one hand.  Thumb, index, long and ring. Thumb, index, ring and little. Thumb, long, ring and little. Thumb, long, ring and little. Index, long, ring and little. Index, long, ring and little.  Thumb, index and long. Thumb, index and ring. Thumb, index and little.

## 38 CFR Ch. I (7-1-07 Edition)

	·	
Diagnostic Code No.		
5138 5139	Index, long and ring.	
5140	Index, long and little. Index, ring and little.	
5141	Long, ring and little.	
	<u> </u>	
Two digits of one hand:	Through and index	
5142 5143	Thumb and index. Thumb and long.	
5144	Thumb and ring.	
5145	Thumb and little.	
5146	Index and long.	
5147 5148	Index and ring. Index and little.	
5149	Long and ring.	
5150	Long and little.	
5151	Ring and little.	
Ois als finances		
Single finger: 5152	Thumb.	
5153	Index finger.	
5154	Long finger.	
5155	Ring finger.	
5156	Little finger.	
	Amputations: Lower Extremity	
<del></del>	<u> </u>	
Thigh amputation of:	Discreticulation	
5160 5161	Disarticulation. Upper third.	
5162	Middle or lower thirds.	
Leg amputation of:	Mark detection of the	
5163 5164	With defective stump.  Not improvable by prosthesis controlled by natural knee action.	
5165	At a lower level, permitting prosthesis.	
5166	Forefoot, proximal to metatarsal bones.	
5167	Foot, loss of use of.	
5170	Toes, all, without metatarsal loss.	
5171 5172	Toe, great. Toes, other than great, with removal of metatarsal head.	
5173	Toes, three or more, without metatarsal involvement.	
	Shoulder and Arm	
	Shoulder and Arm	
5200	Scapulohumeral articulation, ankylosis.	
5201	Arm, limitation of motion.	
5202 5203	Humerus, other impairment.  Clavicle or scapula, impairment.	
	Olaviole of Scapula, Impairment.	
	Elbow and Forearm	
5205	Elbow, ankylosis.	
5206	Forearm, limitation of flexion.	
5207	Forearm, limitation of extension.	
5208	Forearm, flexion limited.	
5209	Elbow, other impairment.	
5210 5211	Radius and ulna, nonunion. Ulna, impairment.	
5212	Radius, impairment.	
5213	Supination and pronation, impairment.	
Wrist		
5214 5215	Wrist, ankylosis. Wrist, limitation of motion.	
JZ IJ	virist, illintation of motion.	
Limitation of Motion		
Multiple Digits: Unfavorable Anky	vlosis:	
5216	Five digits of one hand.	
5217	Four digits of one hand.	
5218	Three digits of one hand.	
5219	Two digits of one hand.	
Multiple Digits: Favorable Ankylosis:		

## **Department of Veterans Affairs**

Diagnostic Code No.	
·	
5220	Five digits of one hand.
5221	Four digits of one hand.
5222	Three digits of one hand.
5223	Two digits of one hand.
A 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
Ankylosis of Individual Digits:	Thurst
5224	Thumb.
5225	Index finger.
5226	Long finger.
5227	Ring or little finger.
Limitation of Motion of Individual	Digits:
5228	Thumb.
5229	Index or long finger.
5230	Ring or little finger.
-	
	Spine
5235	Vertebral fracture or dislocation.
5236	Sacroiliac injury and weakness.
5237	Lumbosacral or cervical strain.
5238 5239	Spinal stenosis.
5240	Spondylolisthesis or segmental instability.  Ankylosing spondylitis.
5241	Spinal fusion.
5242	Degenerative arthritis.
5243	Intervertebral disc syndrome.
	Hip and Thigh
5050	
5250	Hip, ankylosis.
5251 5252	Thigh, limitation of extension.
	Thigh, limitation of flexion.
5253 5254	Thigh, impairment. Hip, flail joint.
5255	Femur, impairment.
	Temat, impairment.
	Knee and Leg
5256	Knee, ankylosis.
5257	Knee, ankylosis. Knee, other impairment.
5257 5258	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated.
5257	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal.
5257	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion.
5257 5258 5259 5260 5261	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension.
5257	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment.
5257 5258 5259 5260 5261	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension.
5257	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment.
5257 5258 5259 5260 5261 5262 5263	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle
5257 5258 5259 5260 5261 5262 5263	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis.
5257 5258 5259 5260 5261 5262 5263 5270 5270	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion.
5257 5258 5259 5260 5261 5262 5263 5270 5270 5272	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion.
5257 5258 5259 5260 5261 5262 5263 5270 5270 5272	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5272 5273 5274	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5272 5273 5274	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273 5274	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5272 5273 5274	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273 5274 5275 5276 5276 5276	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273 5274 5275 5276 5276 5277 5278	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273 5274 5274 5275 5276 5276 5277 5276 5277 5278 5278	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease).
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273 5274 5275 5276 5276 5277 5278 5279 5279 5279 5279	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux valgus.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5273 5274 5275 5276 5277 5278 5279 5280 5280	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux valgus.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5274 5275 5275 5278 5278 5278 5279 5280 5281 52528	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux rigidus. Hallux rigidus. Hammer toe.
5257       5258       5259       5260       5261       5262       5263       5270       5271       5272       5273       5274       5275       5276       5277       5278       5280       5281       5282       5283	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of fexension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux valgus. Hallux rigidus. Hammer toe. Tarsal or metatarsal bones.
5257 5258 5259 5260 5261 5262 5263 5270 5271 5272 5274 5275 5275 5278 5278 5278 5279 5280 5281 52528	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux rigidus. Hallux rigidus. Hammer toe.
5257       5258       5259       5260       5261       5262       5263       5270       5271       5272       5273       5274       5275       5276       5277       5278       5280       5281       5282       5283	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux valgus. Hallux rigidus. Hammer toe. Tarsal or metatarsal bones.
5257       5258       5259       5260       5261       5262       5263       5270       5271       5272       5273       5274       5275       5276       5277       5278       5279       5280       5281       5282       5283       5284	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux valgus. Hammer toe. Tarsal or metatarsal bones. Foot injuries, other.  The Skull
5257       5258       5259       5260       5261       5262       5263       5270       5271       5272       5273       5274       5275       5276       5277       5278       5280       5281       5282       5283	Knee, ankylosis. Knee, other impairment. Cartilage, semilunar, dislocated. Cartilage, semilunar, removal. Leg, limitation of flexion. Leg, limitation of extension. Tibia and fibula, impairment. Genu recurvatum.  Ankle  Ankle, ankylosis. Ankle, limited motion. Subastragalar or tarsal joint, ankylosis. Os calcis or astragalus, malunion. Astragalectomy.  Shortening of the Lower Extremity  Bones, of the lower extremity  The Foot  Flatfoot, acquired. Weak foot, bilateral. Claw foot (pes cavus), acquired. Metatarsalgia, anterior (Morton's disease). Hallux valgus. Hallux rigidus. Hammer toe. Tarsal or metatarsal bones. Foot injuries, other.

Dispusatio Octo No.			
Diagnostic Code No.			
The Ribs			
5297	Removal of.		
	The Coccyx		
5298	Removal of.		
	MUSCLE INJURIES Shoulder Girdle and Arm		
5301	Group I Function: Upward rotation of scapula. Group II Function: Depression of arm.		
5303	Group III Function: Depression of arm.		
5304	Group IV Function: Stabilization of shoulder.		
5305 5306	Group V Function: Elbow supination. Group VI Function: Extension of elbow.		
	Forearm and Hand		
5307	Group VII Function: Flexion of wrist and fingers.		
5308	Group VIII Function: Extension of wrist, fingers, thumb.		
5309	Group IX Function: Forearm muscles.		
	Foot and Leg		
5310	Group X Function: Movement of forefoot and toes.		
5311	Group XI Function: Propulsion of foot. Group XII Function: Dorsiflexion.		
	Pelvic Girdle and Thigh		
5313	Group XIII Function: Extension of hip and flexion of knee.		
5314	Group XIV Function: Extension of knee. Group XV Function: Adduction of hip.		
5316	Group XVI Function: Flexion of hip.		
5317	Group XVII Function: Extension of hip.		
5318	Group XVIII Function: Outward rotation of thigh.		
	Torso and Neck		
5319	Group XIX Function: Abdominal wall and lower thorax.		
5320 5321	Group XX Function: Postural support of body.  Group XXI Function: Respiration.		
5322	Group XXII Function: Respiration:  Group XXII Function: Rotary and forward movements, head.		
5323	Group XXIII Function: Movements of head.		
	Miscellaneous		
5324	Diaphragm, rupture.		
5325	Muscle injury, facial muscles.		
5326 5327	Muscle hernia.  Muscle, neoplasm of, malignant.		
5328	Muscle, neoplasm of, benign.		
5329	Sarcoma, soft tissue.		
	THE EYE Diseases of the Eye		
6000	Uveitis.		
6001	Keratitis.		
6002	Scleritis.		
6003 6004	Iritis.		
6005	Cyclitis. Choroiditis.		
6006	Retinitis.		
6007	Hemorrhage, intra-ocular, recent.		
6008 6009	Retina, detachment.  Eye, injury of, unhealed.		
6010	Eye, injury or, unnealed. Eye, tuberculosis.		
6011	Retina, localized scars.		
6012	Glaucoma, congestive or inflammatory.		
6013	Glaucoma, simple, primary, noncongestive.		
6014 6015	New growths, malignant, eyeball.  New growths, benign, eyeball and adnexa.		

6016	Nystagmus, central.
6017	Conjunctivitis, trachomatous, chronic.
5018	Conjunctivitis, other, chronic.
6019	Ptosis unilateral or bilateral.
6020	Ectropion.
6021	Entropion.
6022	Lagophthalmos.
6023	Eyebrows, loss.
6024	Eyelashes, loss.
6025	Epiphora.
8026	Neuritis, optic.
6027	Cataract, traumatic.
6028	Cataract, senile, and others.
6029	Aphakia.
6030	
	Accommodation, paralysis.
5031	Dacryocystitis.
5032	Eyelids, loss of portion.
6033	Lens, crystalline, dislocation.
6034	Pterygium.
6035	Keratoconus.
	Impairment of Central Visual Acuity
6061	Anatomical loss both eyes.
6062	Blindness, both eyes, only light perception.
Anatomical loss of 1 eye:	
6063	Other eye 5/200 (1.5/60).
5064	
	Other eye 10/200 (3/60).
5064	Other eye 15/200 (4.5/60).
6064	Other eye 20/200 (6/60).
6065	Other eye 20/100 (6/30).
6065	Other eye 20/70 (6/21).
6065	Other eye 20/50 (6/15).
6066	Other eye 20/40 (6/12).
Blindness in 1 eye, only light perd	ception:
6067	Other eye 5/200 (1.5/60).
6068	Other eye 10/200 (3/60).
6068	Other eye 15/200 (4.5/60).
6068	Other eye 20/200 (6/60).
6069	Other eye 20/100 (6/30).
6069	Other eye 20/70 (6/21).
6069	Other eye 20/50 (6/15).
6070	Other eye 20/40 (6/12).
Vision in 1 eye 5/200 (1.5/60):	
6071	Other eye 5/200 (1.5/60).
6072	Other eye 10/200 (3/60).
5072	Other eye 15/200 (4.5/60).
6072	Other eye 20/200 (6/60).
6073	Other eye 20/100 (6/30).
6073	Other eye 20/70 (6/21).
6073	Other eye 20/50 (6/15).
6074	
	Other eye 20/40 (6/12).
/ision in 1 eye 10/200 (3/60):	
6075	Other eve 10/200 (2/60)
	Other eye 10/200 (3/60).
6075	Other eye 15/200 (4.5/60).
6075	Other eye 20/200 (6/60).
6076	Other eye 20/100 (6/30).
6076	Other eye 20/70 (6/21).
6076	Other eye 20/50 (6/15).
6077	Other eye 20/40 (6/12).
Vision in 4 ave 45/000 (4.5/00)	
Vision in 1 eye 15/200 (4.5/60):	
6075	Other eye 15/200 (4.5/60).
6075	Other eye 20/200 (6/60).
6076	Other eye 20/100 (6/30).
	Other eye 20/70 (6/21).
6076	
	Other eve 20/50 (6/15).
6076 6076 6077	Other eye 20/50 (6/15). Other eye 20/40 (6/12).

Diagnostic Code No.	
0076	Other are 20/400 (0/20)
6076 6076	Other eye 20/100 (6/30).
6076	Other eye 20/70 (6/21). Other eye 20/50 (6/15).
6077	Other eye 20/40 (6/12).
	0.10. 0/0 20.10 (0.12).
Vision in 1 eye 20/100 (6/30):	
6078	Other eye 20/100 (6/30).
6078	Other eye 20/70 (6/21).
6078	Other eye 20/50 (6/15).
6079	Other eye 20/40 (6/12).
Vision in 1 eye 20/70 (6/21):	
6078	Other eye 20/70 (6/21).
6078	Other eye 20/50 (6/15).
6079	Other eye 20/40 (6/12).
)/: :	
Vision in 1 eye 20/50 (6/15): 6078	Other ave 20/E0 (6/1E)
6079	Other eye 20/50 (6/15).  Other eye 20/40 (6/12).
0073	Other 6y6 20/40 (0/12).
Impairment of Field Vision:	
6080	Field vision, impairment.
6081	Scotoma.
Impairment of Musels Functions	1
Impairment of Muscle Function: 6090	Diplopia.
6091	Symblepharon.
6092	Diplopia, limited muscle function.
	THE EAR
6200	Chronic suppurative otitis media.
6201	Chronic nonsuppurative otitis media.
6202	Otosclerosis.
6204	Peripheral vestibular disorders.
6205	Meniere's syndrome.
6207	Loss of auricle.
6208	Malignant neoplasm.
6209	Benign neoplasm.
6210	Chronic otitis externa.
6211 6260	Tympanic membrane. Tinnitus, recurrent.
	Tillinas, recurrent.
	OTHER SENSE ORGANS
6275	Small, complete loca
6276	Smell, complete loss.  Taste, complete loss.
0270	Taste, complete loss.
INFECTIOU	S DISEASES, IMMUNE DISORDERS AND NUTRITIONAL DEFICIENCIES
2000	Ohalawa Asiatia
6300 6301	Cholera, Asiatic. Visceral Leishmaniasis.
6302	Leprosy (Hansen's Disease).
6304	Malaria.
6305	Lymphatic Filariasis.
6306	Bartonellosis.
6307	Plague.
6308	Relapsing fever.
6309	Rheumatic fever. Syphilis.
6311	Tuberculosis, miliary.
6313	Avitaminosis.
6314	Beriberi.
6315	Pellagra.
6316	Brucellosis.
6317	Typhus, scrub.
6318	Melioidosis.
6319	Lyme disease.
6350	Parasitic diseases. Lupus erythematosus.
6351	HIV-Related Illness.
6354	Chronic Fatigue Syndrome (CFS).
	1

Diagnostic Code No.	
	THE RESPIRATORY SYSTEM  Nose and Throat
6502	Septum, nasal, deviation.
6504	Nose, loss of part of, or scars.
6510	Sinusitis, pansinusitis, chronic.
6511	Sinusitis, ethmoid, chronic.
6512	Sinusitis, frontal, chronic.
6513 6514	Sinusitis, maxillary, chronic. Sinusitis, sphenoid, chronic.
6515	Laryngitis, tuberculous.
6516	Laryngitis, chronic.
6518	Laryngectomy, total.
6519	Aphonia, complete organic.
6520	Larynx, stenosis of.
6521 6522	Pharynx, injuries to. Allergic or vasomotor rhinitis.
6523	Bacterial rhinitis.
6524	Granulomatous rhinitis.
	Trachea and Bronchi
6600	Bronchitis, chronic.
6601	Bronchiectasis.
6602	Asthma, bronchial.
6603	Emphysema, pulmonary.
6604	Chronic obstructive pulmonary disease.
	Lungs and Pleura Tuberculosis
Ratings for Pulmonary Tuberculo 6701	sis (Chronic) Entitled on August 19, 1968:   Active, far advanced.
6702	Active, moderately advanced.
6703	Active, minimal.
6704 6721	Active, advancement unspecified.
6722	Inactive, far advanced. Inactive, moderately advanced.
6723	Inactive, minimal.
6724	Inactive, advancement unspecified.
	sis Initially Evaluated After August 19, 1968:
6730	Chronic, active.
6731 6732	Chronic, inactive.  Pleurisy, active or inactive.
	Nontuberculous Diseases
6817	Pulmonary Vascular Disease.
6819 6820	Neoplasms, malignant. Neoplasms, benign.
0020	
6822	Bacterial Infections of the Lung
6823	Actinomycosis.  Nocardiosis.
6824	Chronic lung abscess.
	Interstitial Lung Disease
6825	Fibrosis of lung, diffuse interstitial.
6826	Desquamative interstitial pneumonitis.
6827 6828	Pulmonary alveolar proteinosis.
6829	Eosinophilic granuloma.  Drug-induced, pneumonitis & fibrosis.
6830	Radiation-induced, pneumonitis & fibrosis.
6831	Hypersensitivity pneumonitis.
6832	Pneumoconiosis.
6833	Asbestosis.
	Mycotic Lung Disease
6834	Histoplasmosis.
6835	Coccidioidomycosis.
6836	Blastomycosis.
6837	l Cryptococcosis.

## 38 CFR Ch. I (7-1-07 Edition)

Diagnostic Code No.	
Diagnostic Code 140.	
6838	Aspergillosis.
6839	Mucormycosis.
	Restrictive Lung Disease
	Restrictive Lung Disease
6840	Diaphragm paralysis or paresis.
6841	Spinal cord injury with respiratory insufficiency.
6842	Kyphoscoliosis, pectus excavatum/carinatum.
6843 6844	Traumatic chest wall defect. Post-surgical residual.
6845	Pleural effusion or fibrosis.
6846	Sarcoidosis.
6847	Sleep Apnea Syndromes.
	THE CARRIOVACCHI AR SYSTEM
	THE CARDIOVASCULAR SYSTEM Diseases of the Heart
7000	Valvular heart disease.
7001	Endocarditis.
7002 7003	Pericarditis. Pericardial adhesions.
7004	Syphilitic heart disease.
7005	Arteriosclerotic heart disease.
7006	Myocardial infarction.
7007	Hypertensive heart disease.
7008 7010	Hyperthyroid heart disease. Supraventricular arrhythmias.
7011	Ventricular arrhythmias.
7015	Atrioventricular block.
7016	Heart valve replacement.
7017	Coronary bypass surgery.
7018 7019	Implantable cardiac pacemakers. Cardiac transplantation.
7020	Cardiomyopathy.
	Diseases of the Arteries and Veins
7101	Hypertensive vascular disease.
7110	Aortic aneurysm.
7111	Aneurysm, large artery.
7112	Aneurysm, small artery.
7113	Arteriovenous fistula, traumatic.
7114 7115	Arteriosclerosis obliterans.  Thrombo-angiitis obliterans (Buerger's Disease).
7117	Raynaud's syndrome.
7118	Angioneurotic edema.
7119	Erythromelalgia.
7120	Varicose veins.
7121 7122	Post-phlebitic syndrome. Cold injury residuals.
7123	Soft tissue sarcoma.
	THE DIGESTIVE SYSTEM
7200	Mouth, injuries.
7201	Lips, injuries.
7202	Tongue, loss.
7203	Esophagus, stricture.
7204	Esophagus, spasm.
7205 7301	Esophagus, diverticulum. Peritoneum, adhesions.
7304	Ulcer, gastric.
7305	Ulcer, duodenal.
7306	Ulcer, marginal.
7307	Gastritis, hypertrophic.
7308 7309	Postgastrectomy syndromes. Stomach, stenosis.
7310	Stomach, injury of, residuals.
7311	Liver, injury of, residuals.
7312	Liver, cirrhosis.
7314	Cholecystitis, chronic.
7315	Cholelithiasis, chronic.
7316 7317	Cholangitis, chronic. Gall bladder, injury.
7318	
-	

Diagnostic Code No.	
7319	Colon, irritable syndrome.
7321	Amebiasis.
7322	Dysentery, bacillary.
7323	Colitis, ulcerative.
7324	Distomiasis, intestinal or hepatic.
7325	Enteritis, chronic.
7326	Enterocolitis, chronic.
7327	Diverticulitis.
7328	Intestine, small, resection.
7329	Intestine, large, resection.
7330	Intestine, fistula.
7331	Peritonitis.
7332	Rectum & anus, impairment.
7333	Rectum & anus, stricture.
7334	Rectum, prolapse.
7335	Ano, fistula in.
7336	Hemorrhoids.
7337	Pruritus ani.
7338	Hernia, inguinal.
7339	Hernia, ventral, postoperative.
7340	Hernia, femoral.
7342	Visceroptosis.
7343	Neoplasms, malignant.
7344	Neoplasms, benign.
7345	Liver disease, chronic, without cirrhosis.
7346	Hernia, hiatal.
7347	Pancreatitis.
7348	Vagotomy.
7351	Liver transplant.
7354	Hepatitis C.

### THE GENITOURINARY SYSTEM

7500	Kidney, removal.
7501	Kidney, abscess.
7502	Nephritis, chronic.
7504	Pyelonephritis, chronic.
7505	Kidney, tuberculosis.
7507	Nephrosclerosis, arteriolar.
7508	Nephrolithiasis.
7509	Hydronephrosis.
7510	Ureterolithiasis.
7511	Ureter, stricture.
7512	Cystitis, chronic.
7515	Bladder, calculus.
7516	Bladder, fistula.
7517	Bladder, injury.
7518	Urethra, stricture.
7519	Urethra, fistula.
7520	Penis, removal of half or more.
7521	Penis, removal of glans.
7522	Penis, deformity, with loss of erectile power.
7523	Testis, atrophy, complete.
7524	Testis, removal.
7525	Epididymo-orchitis, chronic only.
7527	Prostate gland.
7528	Malignant neoplasms.
7529	Benign neoplasms.
7530	Renal disease, chronic.
7531	Kidney transplant.
7532	Renal tubular disorders.
7533	Kidneys, cystic diseases.
7534	Atherosclerotic renal disease.
7535	Toxic nephropathy.
7536	Glomerulonephritis.
7537	Interstitial nephritis.
7538	Papillary necrosis.
7539	Renal amyloid disease.
7540	Disseminated intravascular coagulation.
7541	Renal involvement in systemic diseases.
7542	Neurogenic bladder.

### GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

7610 ...... Vulva, disease or injury.

## 38 CFR Ch. I (7-1-07 Edition)

Vagina, disease or injury.
Pregnancy, surgical complications.
Endometriosis.
THE HEMIC AND LYMPHATIC SYSTEMS
Adenitis, tuberculous.
Aplastic anemia.
THE SKIN
Disfigurement of, head, face or neck.
3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
Vasculitis, primary cutaneous. Erythema multiforme.
Scarring alopecia.
Malignant melanoma.
THE ENDOODING OVOTEM
THE ENDOCRINE SYSTEM
Hyperthyroidism.

Diagnostic Code No.		
7904	Hyperparathyroidism.	
7905	Hypoparathyroidism.	
7907	Cushing's syndrome.	
7908	Acromegaly.	
7909	Diabetes insipidus.	
7911	Addison's disease.	
7912	Pluriglandular syndrome.	
7913	Diabetes mellitus.	
7914	Malignant neoplasm.	
7915	Benign neoplasm.	
7916	Hyperpituitarism.	
7917	Hyperaldosteronism. Pheochromocytoma.	
7918 7919		
7919	C-cell hyperplasia, thyroid.	
NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS Organic Diseases of the Central Nervous System		
8000	Encephalitis, epidemic, chronic.	
	Brain, New Growth of	
8002	Malignant.	
8003	Benign.	
8004	Paralysis agitans.	
8005	Bulbar palsy.	
8007	Brain, vessels, embolism.	
8008	Brain, vessels, thrombosis.	
8009	Brain, vessels, hemorrhage.	
8010	Myelitis.	
8011	Poliomyelitis, anterior.	
8012	Hematomyelia.	
8013	Syphilis, cerebrospinal.	
8014	Syphilis, meningovascular.	
8015	Tabes dorsalis.	
8017	Amyotrophic lateral sclerosis.	
8018	Multiple sclerosis.	
8019	Meningitis, cerebrospinal, epidemic.	
8020	Brain, abscess.	
	Spinal Cord, New Growths	
8021	Malignant.	
8022	Benign.	
8023	Progressive muscular atrophy.	
8024	Syringomyelia.	
8025	Myasthenia gravis.	
8045	Brain disease due to trauma.	
8046	Cerebral arteriosclerosis.	
-		
	Miscellaneous Diseases	
8100	Migraine	
8103	Tic, convulsive.	
8104	Paramyoclonus multiplex.	
8105	Chorea, Sydenham's.	
8106	Chorea, Huntington's.	
8107	Athetosis, acquired.	
8108	Narcolepsy.	
The Cranial Nerves		
8205	Fifth (trigeminal), paralysis.	
8207	Seventh (facial), paralysis.	
8209	Ninth (glossopharyngeal), paralysis.	
8210	Tenth (pneumogastric, vagus), paralysis.	
8211	Eleventh (spinal accessory, external branch), paralysis.	
8212	Twelfth (hypoglossal), paralysis.	
8305	Neuritis, fifth cranial nerve.	
8307	Neuritis, seventh cranial nerve.	
8309	Neuritis, ninth cranial nerve.	
8310	Neuritis, tenth cranial nerve.	
8311		
8312	Neuritis, eleventh cranial nerve.	
8312 8405		

Diagnostic Code No.	
8407	Neuralgia, seventh cranial nerve.
8409	Neuralgia, ninth cranial nerve.
8410	Neuralgia, tenth cranial nerve.
8411	Neuralgia, eleventh cranial nerve.
8412	Neuralgia, twelfth cranial nerve.
	Peripheral Nerves
8510	Upper radicular group, paralysis.
8511	Middle radicular group, paralysis.
8512	Lower radicular group, paralysis.
8513	All radicular groups, paralysis.
8514	Musculospiral nerve (radial), paralysis.
8515 8516	Median nerve, paralysis.  Ulnar nerve, paralysis.
8517	Musculocutaneous nerve, paralysis.
8518	Circumflex nerve, paralysis.
8519	Long thoracic nerve, paralysis.
8520	Sciatic nerve, paralysis.
8521	External popliteal nerve (common peroneal), paralysis.
8522	Musculocutaneous nerve (superficial peroneal), paralysis.
8523	Anterior tibial nerve (deep peroneal), paralysis.
8524 8525	Internal popliteal nerve (tibial), paralysis.  Posterior tibial nerve, paralysis.
8526	Anterior crural nerve (femoral), paralysis.
8527	Internal saphenous nerve, paralysis.
8528	Obturator nerve, paralysis.
8529	External cutaneous nerve of thigh, paralysis.
8530	Ilio-inguinal nerve, paralysis.
8540 8610	Soft-tissue sarcoma (Neurogenic origin).
8611	Neuritis, upper radicular group.  Neuritis, middle radicular group.
8612	Neuritis, lower radicular group.
8613	Neuritis, all radicular group.
8614	Neuritis, musculospiral (radial) nerve.
8615	Neuritis, median nerve.
8616	Neuritis, ulnar nerve.
8617 8618	Neuritis, musculocutaneous nerve.  Neuritis, circumflex nerve.
8619	Neuritis, long thoracic nerve.
8620	Neuritis, sciatic nerve.
8621	Neuritis, external popliteal (common peroneal) nerve.
8622	Neuritis, musculocutaneous (superficial peroneal) nerve.
8623	Neuritis, anterior tibial (deep peroneal) nerve.
8624 8625	Neuritis, internal popliteal (tibial) nerve.
8626	Neuritis, posterior tibial nerve.  Neuritis, anterior crural (femoral) nerve.
8627	Neuritis, internal saphenous nerve.
8628	Neuritis, obturator nerve.
8629	Neuritis, external cutaneous nerve of thigh.
8630	Neuritis, ilio-inguinal nerve.
8710	Neuralgia, upper radicular group.
8711 8712	Neuralgia, middle radicular group.  Neuralgia, lower radicular group.
8713	Neuralgia, ill radicular groups.
8714	Neuralgia, musculospiral nerve (radial).
8715	Neuralgia, median nerve.
8716	Neuralgia, ulnar nerve.
8717 8718	Neuralgia, musculocutaneous nerve.  Neuralgia, circumflex nerve.
8719	Neuralgia, long thoracic nerve.
8720	Neuralgia, sciatic nerve.
8721	Neuralgia, external popliteal nerve (common peroneal).
8722	Neuralgia, musculocutaneous nerve (superficial peroneal).
8723	Neuralgia, anterior tibial nerve (deep peroneal).
8724 8725	Neuralgia, internal popliteal nerve (tibial).
8726	Neuralgia, posterior tibial nerve.  Neuralgia, anterior crural nerve (femoral).
8727	Neuralgia, internal saphenous nerve.
8728	Neuralgia, obturator nerve.
8729	Neuralgia, external cutaneous nerve of thigh.
8730	Neuralgia, ilio-inguinal nerve.

Diagnostic Code No.	
	The Epilepsies
	The Ephiepsies
8910	
8911 8912	
8913	
8914	Psychomotor.
	Mental Disorders
9201	. Schizophrenia, disorganized type.
9202	Schizophrenia, catatonic type.
9203 9204	
9205	.
9208	Delusional disorder.
9210 9211	1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
9211	
	Delirium, Dementia, Amnestic and Other Cognitive Disorders
9300	
9301 9304	
9305	. Vascular dementia.
9310	
9312 9326	
9327	
	Anxiety Disorders
9400	Generalized anxiety disorder.
9403	
9404	
9410 9411	
9412	
9413	Anxiety disorder, not otherwise specified.
	Dissociative Disorder
9416	
9417	'
	Somatoform Disorders
9421	
9422 9423	
9424	
9425	Hypochondriasis.
	Mood Disorders
9431	1 - 2
9432	
9433 9434	
9435	
	Chronic Adjustment Disorder
9440	<u> </u>
	Eating Disorders
9520	<u> </u>
9521	
	DENTAL AND ORAL CONDITIONS
9900	
9901 9902	
9902	.   Mandible, loss of approximately one-half.

## 38 CFR Ch. I (7-1-07 Edition)

# Pt. 4, App. C

Diagnostic Code No.	
9903	Mandible, nonunion.
9904	Mandible, malunion.
9905	Temporomandibular articulation, limited motion.
9906	Ramus, loss of whole or part.
9907	Ramus, loss of less than one-half.
9908	Condyloid process.
9909	Coronoid process.
9911	Hard palate, loss of half or more.
9912	Hard palate, loss of less than half.
9913	Teeth, loss of.
9914	Maxilla, loss of more than half.
9915	Maxilla, loss of half or less.
9916	Maxilla, malunion or nonunion of.

#### [72 FR 12990, Mar. 20, 2007]

# APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

	Diagnostic code No.
Abscess:	
Brain	802
Kidney	750
Lung	682
Acne	782
Acromegaly	790
Actinomycosis	682
Addison's disease	79
Agranulocytosis	77
Alopecia areata	78
Amebiasis	73:
Amputation:	70.
Arm:	
Disarticulation	51:
Above insertion of deltoid	51
Below insertion of deltoid	51
Digits, five of one hand	51
Digits, four of one hand:	
Thumb, index, long and ring	51
Thumb, index, long and little	51
Thumb, index, ring and little	51
Thumb, long, ring and little	51
Index, long, ring and little	51
Digits, three of one hand:	
Thumb, index and long	51
Thumb, index and ring	51
Thumb, index and little	51
Thumb, long and ring	51
Thumb, long and little	51
Thumb, ring and little	51
Index, long and ring	51
Index, long and little	51
Index, ring and little	51
Long, ring and little	51
Digits, two of one hand:	31
Thumb and index	51
	51
Thumb and long	
Thumb and ring	51
Thumb and little	51
Index and long	51
Index and ring	51
Index and little	51
Long and ring	51
Long and little	51
Ring and little	51
Single finger:	
Thumb	51
Index finger	51
Long finger	51
Ring finger	51
Little finger	51
Forearm:	0.

		Diagnostic code No.
A	Above insertion of pronator teres	512
	Below insertion of pronator teres	512
Leg:	·	
	Vith defective stump	516
	Not improvable by prosthesis controlled by natural knee action	516 516
	At a lower level, permitting prosthesis	516 516
	oes, all, without metatarsal loss	517
	oe, great	517
	oes, other than great, with removal of metatarsal head	517
	oes, three or more, without metatarsal involvement	517
Thigh:		
	Disarticulation	516
	Jpper third	516
	Aiddle or lower thirds	516
	hic lateral sclerosis	801
Anatomica	arioss or: Both eyes	606
	with visual acuity of other eye:	000
One eye,	5/200 (1.5/60)	606
	10/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	606
	20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	606
	20/40 (6/12)	606
Е	Soth feet	510
	Soth hands	510
	One hand and one foot	510
	One foot and loss of use of one hand	510
(	One hand and loss of use of one foot	510
		7700
Aneurysm		
	Aortic	7110
	arge artery	711
	Small arteryortic edema	711: 711:
Ankylosis:		7110
	nkle	5270
Digits, ind		327
Digito, iiia	Thumb	5224
	Index finger	522
	Long finger	5226
	Ring or little finger	522
E	lbow	520
H	Hand	
Favorable		
	ive digits of one hand	5220
	our digits of one hand	522
	Three digits of one hand	5222
	wo digits of one hand	522
Unfavorab	ive digits of one hand	5216
	our digits of one hand	521
	Three digits of one hand	5218
	wo digits of one hand	5219
		5250
		525
Scapulohi	umeral articulation	520
Subastrag	palar or tarsal joint	527
		521
	g spondylitis	524
		602
	organic	651
	nemia	771
Arrhythmi		704
	Supraventricular	701
	/entricularerosis obliterans	701 <sup>-</sup> 711-
	erotic heart disease	700
	ious fistula	711
Arthritis:	local and the second	/ 11
	Degenerative (hypertrophic or osteoarthritis)	500
	Due to trauma	501
	Gonorrheal	500
(		

	Diagnosti code No
Pneumococcic	5
Rheumatoid (atrophic)	5
Streptococcic	5
Syphilitic	5
Typhoid	5
bestosis	6
pergillosis	6
thma, bronchial	6
tragalectomy	5
nerosclerotic renal disease	7
netosis	8
ioventricular block	7
itaminosis	. 6
rtonellosis	6
riberi	ě
adder:	
Calculus in	7
Fistula in	7
Injury of	7
Neurogenic	7
astomycosis	6
ndness: see also Vision and Anatomical Loss	_
Both eyes, only light perception	6
One eye, only light perception and other eye:	_
5/200 (1.5/60)	6
10/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	6
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	6
20/40 (6/12)	6
nes:	
Caisson disease	5
New growths, benign	5
New growths, malignant	5
Shortening of the lower extremity	5
ain:	_
Abscess	8
Disease due to trauma	8
east surgery	7
onchiectasis	6
onchitis	
ucellosis	6
erger's disease	7
lbar palsy	8
Illous disorders	7
rsitis	5
ardiac:	
Pacemakers, implantable	7
Transplantation	7
rdiomyopathy	7
cell hyperplasia, thyroid	7
taract:	
Senile and others	6
Traumatic	è
rebral arteriosclerosis	8
ervical strain	Ę
	7
rvix disease or injury	,
orea:	,
Huntington's	3
Sydenham's	3
loracne	7
olangitis, chronic	7
olecystitis, chronic	7
olelithiasis, chronic	7
olera, Asiatic	6
noroiditis	6
ronic Fatigue Syndrome (CFS)	e
ronic lung abscess	6
	6
pronic obstructive pulmonary disease	6
	7
occidioidomycosis	,
occidioidomycosis	
occidioidomycosis	
oronic obstructive pulmonary disease occidioidomycosis old injury residuals olitis, ulcerative onjunctivitis: or Trachomatous	7

## **Department of Veterans Affairs**

	Diagnosti code No.
Coronary bypass surgery	70
Cryptococcosis	68
Cushing's syndrome	79
Cutaneous manifestations of collagen-vascular diseases	78
Cyclitis	60
Cystitis, chronic	75
Dacryocystitis	60
Permatitis or eczema	78
Dermatophytosis	68
Diabetes:	
Insipidus	79 79
Diaphragm:	
	6
	5:
Diplopia, limited muscle function, eye	6
	79
Buerger's	7
Chronic obstructive pulmonary disease	6
Hodgkin's	7
Diplopia	6
Diplopia, limited muscle function, eye Disease:  Addison's Buerger's Chronic obstructive pulmonary disease Hodgkin's Leprosy (Hansen's) Lyme Morton's Parasitic Disfigurement of, head, face or neck Dislocated: Cartilage, semilunar Lens, crystalline Disseminated intravascular coagulation Distomiasis, intestinal or hepatic Diverticulitis	6
	5
	6
	7
Cartilage, semilunar	5
Lens, crystalline	6
Disseminated intravascular coagulation	7
	7
Diverticulitis	7
Dysentery, bacillary	7
Ectropion	6
mbolism, brain	8
mphysema, pulmonary	6
Encephalitis, epidemic, chronic	8
Endocarditis	7
Endometriosis	7
Enterocolitis, chronic	7
Interception	6
Eosinophilic granuloma of lung	6
Epididymo-orchitis	7
pilepsies:	
Diencephalic	8
Grand mal	8
Jacksonian and focal motor or sensory	8
Petit mal	8
Psychomotor	8
piphora	6
rythema multiforme	7
rythromelalgia	7
sophagus:	_
Diverticulum	7
Spasm	7
Stricture	7
xifoliative dermatitis	7
ever:	·
Relapsing	6
Rheumatic	6
ibrosis of lung, diffuse interstitial	6
ibromyalgia	5
istula in ano	7
istula:	·
Rectovaginal	7
Urethrovaginal	70
Flatfoot, acquired	5
	7
Sastritis, hypertrophic	

		Diagnostic code No.
Glaucom	a:	
	Congestive or inflammatory	60
	Simple, primary, noncongestive	60
Glomerul	onephritis	75
		50
Hallux:		
	Rigidus	52
	Valgus	52
	toe	52
	ve replacement	70
	nyelia	80
Hemorrh		0/
	Brain	80
	Intra-ocular	60
	oids	73 73
перацііs Hernia:	C	73
	Femoral	73
		73
	Hiatal	73
	Inguinal	53
	Ventral	73
Hip:	Degenerative arthritis	E
	ů	52 52
	Flail joint	
	mosis	68
	ted Illness	63
-	s disease	77
	rosis, intermittent	50
	phrosis	75
	osteronism	79
	rosis	78
	rathyroidism	79
	uitarism	79
	nsitivity	68
Hyperten		70
	Heart disease	70 71
	Vascular disease	70
	roid heart disease	79
	roidism	
	athyroidism	79
	oidism	79
Impairme		-
	Humerus	52
	Clavicle or scapula	52
	Elbow	52
	Thigh	52
	Femur	52
	Knee, other	52
	Field vision	60
	Tibia and fibula	52
	Rectum & anus	73
	Ulna	52
	ble cardiac pacemakers	70
	s of the skin	78
Injury:	Platfor	7/
	Bladder	75
	Eye, unhealed	60
	Foot	52
	Gall bladder	73
	Lips	72
	Liver, residuals	73
	Mouth	72
Muscle:	Fasial	
	Facial	53
	Group I Function: Upward rotation of scapula	53
	Group II Function: Depression of arm	50
	Group III Function: Elevation and abduction of arm	53
	Group IV Function: Stabilization of shoulder	50
	Group V Function: Elbow supination	50
	Group VI Function: Extension of elbow	53
	Group VII Function: Flexion of wrist and fingers	53
	Group VIII Function: Extension of wrist, fingers, thumb	5

G		Diagnosti code No.
	roup IX Function: Forearm muscles	50
G	roup X Function: Movement of forefoot and toes	53
G	roup XI Function: Propulsion of foot	53
	roup XII Function: Dorsiflexion	53
	roup XIII Function: Extension of hip and flexion of knee	53
	roup XIV Function: Extension of knee	50
	roup XV Function: Adduction of hiproup XVI Function: Flexion of hip	50 50
G	roup XVII Function: Extension of hip	50
	roup XVIII Function: Outward rotation of thigh	50
	roup XIX Function: Abdominal wall and lower thorax	5
	roup XX Function: Postural support of body	50
G	roup XXI Function: Respiration	53
G	roup XXII Function: Rotary and forward movements, head	53
G	roup XXIII Function: Movements of head	53
harynx		6
		52
	pinal cord	68
	tomach, residuals of	73
		60
	nephritis	7:
	ral disc syndrome	52
	istula of	7:
	lon syndrome	7:
	ion, diseases of	7
		6
	us	6
(idney:	hanna	7
	bscessystic diseases	7: 7:
	emoval	7:
	ransplant	7:
	uberculosis	7:
	iosis, pectus excavatum / carinatum	68
	lmos	60
	omy	6
aryngitis:		
	uberculous	6
С	hronic	6
arynx, ste	enosis of	6
eishmania	asis:	
Α	merican (New World)	7
0	ld World	7
eprosy (H	Hansen's Disease)	6
		7
	of extension:	
imitation i	orearm	5
		J
F	eg	
Fe Le		5
Fe Le R	eg	5 5 5
Fr Lo R S Ti	eg adius	5 5 5
Fo Lo R S Ti imitation	eg adius	5 5 5 5
For London For Indicated For I	eg adius upination and pronation upination and flexion: of extension and flexion:	5 5 5 5
For London For Imitation of Imi	eg adius	5 5 5 5 5
Final	eg adius	5 5 5 5 5
Finitation (Finitation)	eg adius	5 5 5 5 5 5
File London  R S Till S Till S S Till S S Till S T S T S T T T S T S T S T S T S T S	eg adius	5 5 5 5 5 5
Finitation of the control of the con	eg adius	5 5 5 5 5 5 5 5
Final Follows From the Community of the	eg adius upination and pronation bigh of extension and flexion: orearm of flexion: orearm orearm eg eg en bigh of motion: of motion: or flexion: orearm eg eg en bigh of motion: orearm eg en bigh en	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
File Long R R S TI I I I I I I I I I I I I I I I I I	eg adius	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
File London File File File File File File File File	eg adius	5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5
File Long R R R R R R R R R R R R R R R R R R R	eg adius	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
Fig. 1. Fig. 1	eg adius	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
File Long R R S T I I I I I I I I I I I I I I I I I I	eg adius	5 5 5 5 5 5 5 5 5 9 5
File Luck R R S TI	eg adius	5 5 5 5 5 5 5 5 5 9 5
File Let R R S T I I I I I I I I I I I I I I I I I I	eg adius upination and pronation high of extension and flexion: orearm of flexion: orearm eg	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
File Line Line Line Line Line Line Line Lin	eg adius	5 5 5 5 5 5 5 5 5 5 5 5 7 7 7
File Let R R R S TI I I I I I I I I I I I I I I I I I	eg adius	5.5 5.5 5.5 5.5 5.5 5.5 5.5 5.5 5.7 7.7
File Line Line Line Report Services Proceedings of the Line Report Services Proceeding	eg adius	5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.
File Line Research Re	eg adius	5.5 5.5 5.5 5.5 5.5 5.5 5.5 5.5 5.7 7.7
File Lide R S S T T I I I I I I I I I I I I I I I I	eg adius	5 5 5 5 5 5 5 5 5 5 5 7 7 7 7

	Diagnosti code No.
Eyebrows	60
Eyelashes	60
Eyelids	60
fandible:	
One-half	99
Completelaxilla:	98
More than half	99
Less than half	99
ose, part of, or scars	65
alate, hard:	
Half or more  Less than half	99
amus:	33
Whole or part	99
Less than one-half	99
kull, part of	52
nell, sense of	62
ste, sense of	62
eth, loss ofngue, loss of whole or part	99 72
ss of use of:	1.
Both feet	5
Both hands	5
Foot	5
Hand	5
One hand and one foot	5
mbosacral strain	5
ous: Erythematosus	6
Erythematosus, discoid	7
ne disease	6
mphatic filariasis	6
alaria	6
alignant melanoma	78
alunion:  Mandible	99
Os calcis or astragalus	5:
axilla, malunion or nonunion	9:
elioidosis	6:
eniere's syndrome	6:
ningitis, cerebrospinal, epidemicental disorders:	8
Anxiety disorders:	
Generalized anxiety disorder	9
Obsessive compulsive disorder	9
Other and unspecified neurosis	9
Panic disorder	9
Post-traumatic stress disorder	9
Specific (simple) phobia	9
Chronic adjustment disorder	9
Delirium, dementia, amnestic and other cognitive disorders	
	9
Alzheimers	9
Alzheimers	9
Alzheimers  Delirium  Head trauma	۵
Alzheimers Delirium Head trauma Infection	
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions	9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions	9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia	9 9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders:	9 9 9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Amnesia, fugue, identity disorders	9 9 9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Anmesia, fugue, identity disorders Depersonalization disorder	9 9 9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Amnesia, fugue, identity disorders Depersonalization disorder Eating Disorder:	9 9 9 9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Annesia, fugue, identity disorders Depersonalization disorder Eating Disorder: Annesia hugue, identity disorders Annesia, fugue, identity disorders	9 9 9 9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Anmesia, fugue, identity disorders Depersonalization disorder Eating Disorder: Anorexia nervosa Bulimia nervosa	9 9 9 9 9
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Amnesia, fugue, identity disorders Depersonalization disorder  Eating Disorder: Anorexia nervosa Bulimia nervosa Mood Disorders:	9: 9: 9: 9: 9: 9:
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Amnesia, fugue, identity disorders Depersonalization disorder Eating Disorder: Anorexia nervosa Bulimia nervosa	9: 9: 9: 9: 9: 9: 9:
Alzheimers Delirium Head trauma Infection Organic mental disorder Other medical conditions Unknown etiology Vascular dementia Dissociative disorders: Amnesia, fugue, identity disorders Depersonalization disorder  Eating Disorder: Anorexia nervosa Bulimia nervosa Mood Disorders: Bipolar disorder	9: 9: 9: 9: 9: 9:

		Diagnosti code No
Schizophrer	ia and other psychotic disorders:	
	tatonic type	9.
De	usional disorder	9
Dis	organized type	9
	/chotic disorder	9
Pa	ranoid type	9
Re	sidual type	9
Sc	nizoaffective disorder	9
Un	differentiated type	9
Somatoform	:	
Co	nversion disorder	9
Hy	oochondriasis	9
	n disorder	9
	matization disorder	9
	differentiated somatoform disorder	9
Metatarsalgia		5
		8
Morton's disease		5
Mucormycosis		6
Multiple sclerosis		8
Myasthenia gravis		8
		8
		7
Myositis ossificans		5
		5
Narcolepsy		8
Neoplasms:		
Benign:		
Dig	estive system	7
Ea		6
En	docrine	7
Ge	nitourinary	7
Gy	necological or breast	7
Mu	scle	5
Re	spiratory	6
Ski	n	7
Malignant:		
Dig	estive system	7
Ea	,	6
En	docrine	7
_	nitourinary	7
	necological or breast	7
	scle	5
_	spiratory	6
	n	7
		7
		7
	riolar	7
Neuralgia:		•
Cranial Ner	ves	
	h (trigeminal)	8
	venth (facial)	8
	th (glossopharyngeal)	8
	nth (pneumogastric, vagus)	8
	venth (spinal accessory, external branch)	8
	elfth (hypoglossal)	8
Peripheral N		O
	per radicular group	8
	Idle radicular group	8
	ver radicular group	8
	radicular groups	8
	sculospiral (radial)	8
	dian	8
	ar	8
	sculocutaneous	8
	cumflex	8
	ng thoracic	8
	atic	8
	ernal popliteal (common peroneal)	8
	sculocutaneous (superficial peroneal)	8
	erior tibial (deep peroneal)	8
	ernal popliteal (tibial)	8
Inte		

		Diagnost code No
	Anterior crural (femoral)	8
	Internal saphenous	8
	Obturator	8
	External cutaneous nerve of thigh	8
euritis:	llio-inguinal	8
	nerves	
	Fifth (trigeminal)	8
	Seventh (facial)	8
	Ninth (glossopharyngeal) Tenth (pneumogastric, vagus)	8
	Eleventh (spinal accessory, external branch)	8
	Twelfth (hypoglossal)	8
		6
Periphe	eral Nerves	
	Upper radicular group	8
	Middle radicular group Lower radicular group	8
	All radicular groups	8
	Musculospiral (radial)	8
	Median	8
	Ulnar	8
	Musculocutaneous	8
	Circumflex	8
	Long thoracic	8
	External popliteal (common peroneal)	8
	Musculocutaneous (superficial peroneal)	8
	Anterior tibial (deep peroneal)	8
	Internal popliteal (tibial)	8
	Posterior tibial	8
	Anterior crural (femoral)	8
	Internal saphenous  Obturator	8
	External cutaneous nerve of thigh	8
urogenic blad w growths: Benign	lder	7
Dorligh	Bones	5
	Brain	8
	Eyeball and adnexa	6
Maliana	Spinal cord	8
Maligna	Bones	5
	Brain	8
	Eyeball	e
	Spinal cord	8
		6
n-Hodgkin's i nunion:	ymphoma	7
	le	g
	and ulna	5
	tral	6
	ins	5
		5
		5
	axilla or mandible	5
itis media:	ith joint manifestations	5
	3	6
	opurative	ě
	ative	6
Nonsup Suppur		6
Nonsup Suppur osclerosis raries, atrophy	y of both	,
Nonsup Suppur osclerosis raries, atrophy rary:	of both	
Nonsup Suppur osclerosis raries, atrophy rary: Diseaso		7
Nonsup Suppur cosclerosis varies, atrophy vary: Disease Remov alsy, bulbar	e or injuryal	7 7 7 8
Nonsup Suppur osclerosis varies, atrophy vary: Disease Remov alsy, bulbar	e or injury	7 7 8 7
Nonsup Suppur osclerosis varies, atrophy vary: Disease Remov alsy, bulbar ancreatitis apillary necros	e or injuryal	7 7 8

		Diagnostic code No.
Accommodatio	n	60
		80
Paralysis, nerve:		
Cranial nerves		
	trigeminal)	82
	nth (facial)	82
	(glossopharyngeal)	82
	(pneumogastric, vagus)	82
	nth (spinal accessory, external branch)	82 82
Peripheral Ner		02
	r radicular group	85
	e radicular group	85
	r radicular group	85
All rac	dicular groups	85
Muscu	ulospiral (radial)	85
	ın	85
		85
	ulocutaneous	85
	nflexthoragin	85
	thoracic	85 85
	c	85 85
	ulocutaneous (superficial peroneal)	85
	or tibial nerve (deep peroneal)	85
	al popliteal (tibial)	85
	rior tibial nerve	85
	or crural nerve (femoral)	85
	al saphenous	85
Obtura	ator	85
Exterr	nal cutaneous nerve of thigh	85
	guinal	85
	X	81
		63
		63
Penis		63
Penis Deformity, with	loss of erectile power	75
Penis Deformity, with Removal of gla	loss of erectile powerans	75 75
Penis Deformity, with Removal of gla Removal of ha	l loss of erectile power	75 75 75
Penis Deformity, with Removal of gla Removal of hal Pericardial adhesions	I loss of erectile power	75 75 75 70
Denis Deformity, with Removal of gla Removal of hal Pericardial adhesions	I loss of erectile power	75 75 75 70 70
Deformity, with Removal of gla Removal of har Pericardial adhesions Pericarditis	l loss of erectile power	7! 7! 7! 70 70 50
Penis Deformity, with Removal of gla Removal of hai Pericardial adhesions Pericarditis	I loss of erectile power	7: 7: 7: 7: 7: 5: 6:
Deformity, with Removal of gla Removal of ha Pericardial adhesions Pericarditis Periopheral vestibular dis Perioneum, adhesions	I loss of erectile power ans ans are a second of the secon	75 75 75 70 70 50 62 73
Penis  Deformity, with Removal of gla Removal of ha Pericardial adhesions Pericarditis Peripheral vestibular dis Peritoneum, adhesions Peritonitis	I loss of erectile power ans I for more I fo	75 75 75
Penis Deformity, with Removal of gla Removal of ha Pericardial adhesions Pericarditis Pericostitis Peripheral vestibular dis Peritonitis Personantial Peritoneum, adhesions Personantial Personantial Personantial Personantia	I loss of erectile power	7: 7: 7: 7: 7: 5: 6: 7: 7:
Penis Deformity, with Removal of gla Removal of ha Pericardial adhesions Pericarditis Periostitis Periostitis Periostitis Periotheral vestibular dis Peritonetm, adhesions Peritonitis Pes cavus (Claw foot) a Pheochromocytoma Plague	l loss of erectile power	7: 7: 7: 7: 7: 5: 6: 7: 7: 7:
Penis Deformity, with Removal of gla Removal of ha Pericardial adhesions Pericostitis Peripheral vestibular dis Peritoneum, adhesions Peritonitis Pers cavus (Claw foot) a Pheochromocytoma Plague	I loss of erectile power	7: 7: 7: 7: 7: 5: 6: 7: 5: 7: 6: 6: 6:
Penis Deformity, with Removal of gla Removal of hal Removal of hal reicardial adhesions dericarditis retricostitis retributis retrib	I loss of erectile power ans If or more  Sorders  acquired	7: 7: 7: 7: 5: 6: 7: 7: 6: 6:
Penis  Deformity, with Removal of gla Removal of hal Pericarditia adhesions Pericarditis Periostitis Periostitis Periostitis Periotitis Periotitis Periotitis Periotitis Periotitis Periotitis Periotitis Periotitis Pes cavus (Claw foot) a Pleocal refusion or fibros Pelurial effusion or fibros Pelurial andular syndrome Peneumoconiosis	I loss of erectile power	7: 7: 7: 7: 5: 6: 7: 7: 6: 6:
Penis  Deformity, with Removal of gla Removal of gla Removal of halfericardial adhesions dericarditis reripheral vestibular disteritoneum, adhesions deritonitis des cavus (Claw foot) acheochromocytoma delague defusion or fibrosituriglandular syndrome ineumoconiosis denumonitis & fibrosis:	I loss of erectile power	7: 7: 7: 7: 7: 5: 6: 6: 7: 7: 6: 6: 7:
Deformity, with Removal of gla Removal of hal rericardial adhesions dericarditis retributes with retributes re	I loss of erectile power ans If or more  Sorders  Society  Society	7: 7: 7: 7: 5: 6: 7: 6: 6: 6:
enis  Deformity, with Removal of gla Removal of hal Rerioarditis Rerioarditis Rerioarditis Res cavus (Claw foot) a Recorbormocytoma Regue Reval effusion or fibros Reval of hall	I loss of erectile power	7. 7. 7. 7. 5. 6. 7. 5. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6.
Penis Deformity, with Removal of gla Removal of gla Removal of halfericardial adhesions Pericarditis	I loss of erectile power	7: 7: 7: 7: 7: 5: 6: 6: 7: 6: 6: 6: 6: 6: 6:
enis Deformity, with Removal of gla Removal of gla Removal of hat ericardial adhesions ericarditis ericonstitis eripheral vestibular diseritoneum, adhesions eritonitis es cavus (Claw foot) a heochromocytoma lague leural effusion or fibros luriglandular syndrome neumoconiosis neumonitis & fibrosis: Drug-induced Radiation-inducoliomyelitis, anterior olycythemia vera	I loss of erectile power ans ans are also are al	7, 7, 7, 7, 5, 6, 7, 7, 6, 6, 6, 6, 6, 6, 6, 7, 7, 6, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7,
enis  Deformity, with Removal of gla Removal of gla Removal of hal ericardial adhesions ericarditis eriostitis eriostitis eriostitis eriostitis eriothem, adhesions eritoneum, adhesions eritonitis es cavus (Claw foot) a heochromocytoma lague lague lague lauriglandular syndrome neumoconiosis neumonitis & fibrosis: Drug-induced Radiation-induc oliomyelitis, anterior olycythemia vera ostgastrectomy syndrogome	I loss of erectile power ans	7. 7. 7. 7. 5. 6. 7. 5. 6. 6. 6. 6. 6. 6. 8. 7. 7.
enis  Deformity, with Removal of gla Removal of gla Removal of hale ricardital adhesions are ricarditis eriostitis eriostitis eriopheral vestibular diseritoneum, adhesions eritonitis es cavus (Claw foot) a heochromocytoma lague leural effusion or fibros luriglandular syndrome neumoconiosis neumonitis & fibrosis: Drug-induced Radiation-induo oliomyelitis, anterior olycythemia vera ost-phlebitic syndrome ost-phlebitic syndrome ost-phlebitic syndrome ost-phlebitic syndrome	I loss of erectile power	7. 7. 7. 7. 5. 6. 7. 7. 6. 6. 6. 6. 8. 7. 7.
enis  Deformity, with Removal of gla Removal of gla Removal of ha ericardial adhesions ericarditis eritoneum, adhesions eritoneum, adhesions eritonitis ses cavus (Claw foot) a heochromocytoma lague elural effusion or fibros luriglandular syndrome neumoconiosis neumonitis & fibrosis: Drug-induced Radiation-indur oliomyelitis, anterior ostgastrectomy syndro ost-phlebitic syndrome ost-surgical residual resid	a loss of erectile power ans ans are also and a loss of erectile power ans are also	7, 7, 7, 7, 5, 6, 7, 6, 6, 6, 6, 6, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7, 7,
enis  Deformity, with Removal of gla Removal of pla Removal of hal Rericarditis Rericarditis Rericarditis Res cavus (Claw foot) a Reochromocytoma Rescavus (Claw foot) a Recordition of heochromocytoma Rescavus (Claw foot) a Recordition of hal Registration of hal Registrat	I loss of erectile power ans ans ans ans ans ans ans ans ans and ans ans ans ans and ans ans and ans ans and and ans a	7: 7: 7: 7: 5: 6: 6: 6: 6: 6: 6: 8: 8: 7: 7:
Penis  Deformity, with Removal of gla Removal of gla Removal of halfericardial adhesions dericarditis .	a loss of erectile power ans ans are also and a loss of erectile power ans are also	7: 7: 7: 7: 7: 5: 6: 6: 6: 7: 6: 6: 6: 8: 7: 7: 7: 7: 6: 8: 8: 7: 7: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8:
enis  Deformity, with Removal of gla Removal of gla Removal of ha ericardial adhesions ericarditis erichatitis erichatitis eritheau yestibular dis eritoneum, adhesions eritonitis es cavus (Claw foot) a heochromocytoma lague elural effusion or fibros luriglandular syndrome neumoconiosis neumonitis & fibrosis:  Drug-induced Radiation-inducoliomyelitis, anterior ostgastrectomy syndro ost-phlebitic syndrome ost-purgical residual regnancy, surgical con rogressive muscular ar rostate gland	loss of erectile power	7: 7: 7: 7: 7: 5: 6: 6: 6: 7: 6: 6: 6: 8: 7: 7: 7: 7: 8:
enis  Deformity, with Removal of gla Removal of gla Removal of hal ericardial adhesions ericarditis eriostitis eriostitis eriostitis eriostitis eriostitis erioneum, adhesions eritoneum, adhesions eritoneum, adhesions eritonitis es cavus (Claw foot) a heochromocytoma lague es cavus (Claw foot) a heochromocytoma lague es cavus (Ulaw foot) en eumoconiosis en	I loss of erectile power ans lif or more sorders sacquired sis sis sis sis sis sis sis sis sis si	7: 7: 7: 7: 7: 5: 6: 6: 6: 6: 6: 8: 8: 7: 7: 7: 8: 8: 7:
Penis  Deformity, with Removal of gla Removal of gla Removal of halfericardial adhesions Pericarditis Pericarditis Peripheral vestibular dis Peripheral vestibular dis Peripheral vestibular dis Peripheral vestibular dis Peripheral yestibular dis Penachronocytoma Penachronocytoma Pleural effusion or fibros Purugiandular syndrome Ineumoconiosis Preumonitis & fibrosis: Drug-induced Radiation-induced loilomyelitis, anterior Postgastrectomy syndrost-surgical residual Pergnancy, surgical con rogressive muscular a rostate gland Prosthetic Implants: Ankle replacem	loss of erectile power	7. 7. 7. 7. 5. 6. 6. 7. 6. 6. 6. 8. 7. 7. 7. 6. 8. 7. 7. 8. 8. 7. 7. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8.
Penis  Deformity, with Removal of gla Removal of gla Removal of ha rericardial adhesions Pericarditis	I loss of erectile power ans ans alf or more ans acquired	7: 7: 7: 7: 7: 5: 6: 6: 7: 6: 6: 6: 6: 6: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7:
Penis Deformity, with Removal of gla Removal of gla Removal of hat Pericarditis Pericarditis Pericarditis Pericarditis Pericheral vestibular dis Perichenum, adhesions Peritonitis Perichenum, adhesions Peritonitis Perichechromocytoma Pelaural effusion or fibros Pelaural effusion or fibros Penumoconiosis Pe	I loss of erectile power ans If or more  Sorders  Sorders	7: 7: 7: 7: 7: 5: 6: 6: 6: 6: 6: 6: 7: 7: 7: 6: 8: 8: 7: 7: 7: 8: 8: 7: 7: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8:
Penis  Deformity, with Removal of gla Removal of gla Removal of hat Pericardital adhesions Pericheral vestibular dis Periosititis	I loss of erectile power ans If or more  Sorders  Society  Society	7: 7: 7: 7: 7: 5: 6: 6: 6: 6: 6: 6: 6: 6: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7: 7:
Penis  Deformity, with Removal of gla Removal of gla Removal of ha Pericardial adhesions Pericarditis	I loss of erectile power ans If or more  Sorders  Society  Society	75 75 76 76 66 77 75 66 75 66 75 66 77 77 77 77 64 86 87 87 87 87 87 87 87 87 87 87 87 87 87
Penis  Deformity, with Removal of gla Removal of gla Removal of hall Removal o	I loss of erectile power ans  If or more  Sorders  Sorders  Social Socia	75 75 70 70 50 62 73
Penis  Deformity, with Removal of gla Removal of gla Removal of ha Pericardial adhesions Periostitis	I loss of erectile power ans If or more  Sorders  Society  Society	7: 7: 7: 7: 7: 7: 5: 6: 6: 6: 6: 6: 7: 7: 7: 8: 8: 7: 7: 7: 8: 7: 7: 8: 8: 7: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8: 8:

Alveolar proteinosis  Vascular diseases  Purtrius  Vascular diseases  Perploapphrist, chronic  Raynauxi syndrome  Ractum:  Rectum & anus, stricture  Prolapse  Removal:  Cardiage, semilunar  Coccyx  Gall bladder  Kidney  Penis glans  Resis  Vary  Ulerus  Ulerus  Ulerus  Ulerus  Ulerus  Ulerus  Ulerus  Horis  Arnyloid diseases  Diseases, chronic  Involvement in systemic diseases  Tubular disorders  Relain detachment of  Refinitis  Relain detachment of sessions  Resis		Diagnos code N
Vascular diseases Pyelonephrits, chronic Pyelonephrits, chronic Return & anus, stricture Prolapse Reroval  Cartilage, semilunar Cocoyx Gail bladder Kense Penis half or more Riss Riss Testis Ovary Ulterus and both ovaries  Tental  Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders  Vertinits Allergic or vasomotor Bacterial Grandomatous  Resection of intestinet Large Coraniomatous  Resection of intestinet Large Large Large Lorge Lorg	Alveolar proteinosis	
yelonephritis, chronic yanaud's syndrome ectum: Rectum & anus, stricture Prolapse emoval: Carliage, semilunar Coccyx Gall bladder Kddney Penis glans P		
synaud's syndrome ectum: Rectum & anus, stricture Prolapse emroval: Carillage, semilunar Coccyx Gall bladder Kidney Penis glans Penis glans Penis half or more Ribs Testis Ovary Uterus and both ovaries anal Uterus and both ovaries anal Amyloid disease Disease, chronic Involvement in systemic diseases Tutular disorders Tutular disorders Allergic or vasomotor Bacterial Granulomatous sesection of intestine: Large Small arcoidosis ararring alopecia arars: Deep, other than head, face or neck Other Retina Superficial, unstable Jeritial Jerit	uritus ani	
ectum: Rectum & anus, stricture Prolapse emoval: Cartilage, semilunar Coccyx Gall bladder Kidney Penis glans Penis half or more Ribs Testis Ovary Uterus Uterus and both ovaries enal: Anyloid disease Disease, chronic Involvement in systemic diseases Trubular disorders ettinitis Allergic or vasomotor Bacterial Allergic or vasomotor Bacterial Allergic or vasomotor Bacterial Allergic or half or more Bacterial Arocolosis Somali arocolosis Somali Allergic or half or more Cartillosis Cartillosis Superficial, unitable Clerris Superficial, unstable Clerris Cotoma Superficial, unstable Clerris Superficial, painful Super	/elonephritis, chronic	
Rectum & anus, stricture Prolapse emoval: Cartiage, semilunar Coccyx Gall bladder Kidney Penis glans Penis glans Penis half or more Ribs Testis Ovary Ulerus and both ovaries email: Anyloid disease Disease, chronic Tubular disorders  Tubular disorders  If under disor	aynaud's syndrome	
Prolapse emoval:  Cartilage, semilunar Cocoyx Gall bladder Kidney Penis glans — Penis half or more Ribs Testis Ovary Uterus Uterus Uterus Uterus Uterus Amyloid disease Disease, chronic Involvement in systemic diseases Trubular disorders  Allergic or vasomotor Bacterial Bact		
cardiage, semilunar Cococyx Call bladder Kidney Penis glans Penis half or more Ribs Testis Ovary Ulerus Ulerus and both ovaries enal: Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders ettina detachment of ettinitis Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia carris: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, or ther head, face, or neck Superficial, painful Superficial Superficial, painful Superficial Superficial Superficial Superficial Superficial Supe	Rectum & anus, stricture	
Cartiage, semiunar Cocoyx Gall bladder Kidney Penis glans Penis half or more Ribs Testis Ovary Uterus Uterus Uterus Uterus Uterus Uterus Blasses, chronic Involvement in systemic diseases Trubular disorders Trubular disorders Involvement of eteitinis shiritis: Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia cartis  Deep, other than head, face or neck Other Retina Superficial, unstable cleritis Superficial, painful Superincial, p	Prolapse	
Coccyx Gall bladder Kidney Penis glans Penis half or more Ribs Testis Ovary Ulerus Ulerus and both ovaries enal: Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders etina detachment of etinitis Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis cars: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable cleritis cotoma Superficial, unstable Cleritis Cotoma Superficial, unstable Cleritis Frontal Maxillary Pansinustis Sphenoid leep Apnea Syndrome oft issue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascurder Vascurder origin Varcurder origin V	emoval:	
Gall bladder Kidney Penis glans Penis half or more Ribs Testis Ovary Ulerus Ulerus Ulerus and both ovaries enal: Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders elina detachment of elinitis hinitis: Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoldosis Large Small Caranulomatous esection of intestine: Large Small Superficial, other than head, face or neck Other Retina Superficial, unstable cleritis cotiona Superficial, unstable cleritis Cotiona Superficial, instable cleritis Elthnoid Frontal Maxillary Pansinusitis Sphenoid Sphenoid Sphenoid Vascular origin Varioromes Chronic Fatigue Syndrome (CFS) Cushing's Cushing's	Cartilage, semilunar	
Kidney Penis palf or more Ribs Testis Ovary Uterus Uterus and both ovaries enal: Arryloid disease Disease, chronic Involvement in systemic diseases Trubular disorders atinuolyment in systemic diseases Trubular disorders etinitis Allergic or vasomotor Bacterial Arranulomatous esection of intestine: Large Small arrocidosis carring alopecia carris: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superfici	Coccyx	
Penis plans Penis half or more Ribs Testis Ovary Ulterus Ulterus and both ovaries enal:  Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders etinia detachment of etinitis hinitis: Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia cortina superficial, other than head, face or neck Superficial, other than head, face, or neck Superficial, other than head, face, or neck Superficial, other than head, face, or neck Superficial, unstable cottoma epturm, nasal, deviation of ciccide cell alore a	Gall bladder	
Penis half or more Ribs Testis Ovary Ulerus Ulerus and both ovaries enal:  Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders etinal detachment of etinitis Allergic or vasomotor Bacterial Granulomatous essection of intestine: Large Small arcoidosis acarring alopecia carring alopecia carring alopecia carris Deep, other than head, face or neck Other Retina Superficial, painful Superficial, painful Superficial, painful Superficial, unstable cleritis Celtritis Eltmoid Frontal Maxillary Pansinustis Sphenoid Isepa Anaica Sphenoid Isepa Anaica Isepa Sphenoid Isepa Apnea Syndrome of tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vasculer origin Vasculer origin Injury of, healed Iplenetomy pondylolisthesis or segmental Instability, spine tomach, stenosis of ymblepharon ymdromes: Chronic Fatigue Syndrome (CFS) Cushing's	Kidney	
Ribs Testis Ovary Uterus and both ovaries enal:  Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders etina detachment of etinitis Initis: Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia carring alopecia carring alopecia carris Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, other than head, face, or neck Superficial, instable cleritis cotoma eptum, nasal, deviation of citckel cell anemia inustitis: Ethmoid Frontal Maxillary Pansinustitis Sphenoid leep Apnea Syndrome of tissue sacroma: Muscle, fat, or fibrous connected Neurogenic origin pinal stenosis plenet, injury of, healed plenetcomy pondyloilsthesis or segmental instability, spine tomach, stenosis of ymblepharon ymblepha	Penis glans	
Testis Ovary Uterus Uterus Uterus and both ovaries  Lenal:  Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders  Etima detachment of Lettiniis Allergic or vasomotor Bacterial Granulomatous  Large Small Aaroidosis Large Small Aaroidosis Large Small Deep, other than head, face or neck Other Retina Superficial, paintul Superficial, pai		
Ovary Uterus and both ovaries enal: Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders etina detachment of etinitis hinitis: Allergic or vasomotor Bacterial Granulomatous escetion of intestine: Large Small arcoidosis carring alopecia cars: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, other than head, face, or neck superficial, unstable cleritis cotoma epitum, asaal, deviation of ickle cell anemia inussitis: Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome off tissue sacroma: Muscle, fat, or fibrous connected Neurogenic origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondyloitshesis or segmental instability, spine tomach, stenosis of ymitherparon ymitoremes: Chronic Fatigue Syndrome (CFS) Cushing's	Ribs	
Uterus and both ovaries enal:  Arnyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders  Involvement in systemic diseases Tubular disorders  Involvement in systemic diseases Tubular disorders  Involvement in systemic diseases Involvement in systemic diseases Involvement of ettinits Intitis:  Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, painful Superficial, painful Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinustits Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin pinal fusion pinal stenosis pleen, injury of, healed plenenctorny pondylolisthesis or segmental instability, spine tomach, stenosis of ymitolepharon ymidemens: Chronic Fatigue Syndrome (CFS) Cushing's	Testis	
Uterus and both ovaries enal: Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders etima detachment of etimitis Inititis: Allergic or vasomotor Bacterial Granulomatous escition of intestine: Large Small arcoidosis acarring alopedia cars: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, unstable cleritis cotoma Superficial, unstable cleritis Cotoma Peptum, nasal, deviation of cockle cell anemia Initustitis: Ethmoid Frontal Maxillary Pansinustitis Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin pinal fusion pinal stenosis plenen, injury of, healed plenen, injury of, healed plenen, injury of, healed plenen, injury of, healed plenen, or fortious consecutions comarch, stenosis of ymblepharon ymblephar	Ovary	
Amyloid disease Disease, chronic Involvement in systemic diseases Tubular disorders  etina detachment of etinitis hinitis:  Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia carring alopecia cars:  Dep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, other than head, face, or neck superficial, unstable cleritis cototoma eptum, nasal, deviation of ickle cell anemia inustits:  Ethmoid Frontal Maxillary Pansinustitis Sphenoid leep Apnea Syndrome oft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal flusion pinal stenosis plenen, injury of, healed pleneet, injury of, healed pleneet, injury of, healed pleneet, proposed promotines Cushing's Cushing's	Uterus	
Amyloid disease Disease, chronic Disease	Uterus and both ovaries	
Disease, chronic Involvement in systemic diseases Tubular disorders etina detachment of etinitis Finitis:  Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcordoosis acarring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable cleritis cottoma eptum, nasal, deviation of cikek cell anemia inusitis: Ethmoid Frontal Maxillary Panasinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma: Muscl, fat, or fibrous connected Neurogenic origin pinal flusion pinal stenosis pleen, injury of, healed pleneet, niyury of, healed pleneet, senosis of ymblepharon ymblepharon ymblepharon ymbromes Cushing's Cushing's	enal:	
Disease, chronic Involvement in systemic diseases Tubular disorders  teitina detachment of eteinitis  Allergic or vasomotor Bacterial Granulomatous esection of intestine:  Large Small arcoidosis carring alopecia carring alopecia carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck superficial, instable cleritis cottoma eptum, nasal, deviation of icickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma:  Muscl, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis plenen, injury of, healed plenectomy pondylolishesis or segmental instability, spine tomach, stenosis of ymblepharon ymblepharon ymblepharon ymblepharon ymblepharon ymbromes: Chronic Fatigue Syndrome (CFS) Cushing's	Amyloid disease	
Involvement in systemic diseases Tubular disorders  tetina detachment of etinitis Allergic or vasomotor Bacterial Granulomatous  tesection of intestine: Large Small arcoidosis carring alopecia cars: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis: Ethmoid Frontal Maxillary Pansinusitis Sphenoid Musce, fat, or fibrous connected Neurogenic origin Vascular origin pinal tusion pinal stenosis plenen, injury of, healed plenectiony pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon ymbleph		
tetinal detachment of tetinits:  Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia carris: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, unstable cleritis cotoma eptum, nasal, deviation of lickle cell anemia inusitis: Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal stenosis pienen, injury of, healed plenectiomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	Involvement in systemic diseases	
etinitis hinitis: Allergic or vasomotor Bacterial Granulomatous esection of intestine:  Large Small arcoidosis carring alopecia cars: Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable cleritis cotoma epitum, nasal, deviation of ickle cell anemia inusitis: Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscele, fat, or fibrous connected Neurogenic origin Vascular origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectory pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	Tubular disorders	
hinitis:  Allergic or vasomotor Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin vascular origin pinal fusion pinal fusion pinal stenosis pleen, injury of, healed plenectory pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	etina detachment of	
Allergic or vasomotor Bacterial Granulomatous esection of intestine:  Large Small arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	etinitis	
Bacterial Granulomatous esection of intestine: Large Small arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, inter than head, face, or neck Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	ninitis:	
Granulomatous desection of intestine:  Large Small arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable cotoma eptum, nasal, deviation of ickle cell anemia inustits:  Ethmoid Frontal Maxillary Pansinustits Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	Allergic or vasomotor	
lesection of intestine:  Large Small  arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, painful Superficial, instable cleritis cotoma eptum, nasal, deviation of ickle cell anemia insustits:  Ethmoid Frontal Maxillary Pansinustitis Sphenoid leep Apnea Syndrome oft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	Bacterial	
Large Small arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, paintul Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis: Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin vascular origin pinal fusion pinal stenosis pleend, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	Granulomatous	
Small araciolosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	esection of intestine:	
arcoidosis carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome off tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	Large	
carring alopecia cars:  Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable Cleritis cotoma eptum, nasal, deviation of ickle cell anemia inusitis: Ethmoid Frontal Maxillary Pansinusitis Sphenoid Leep Apnea Syndrome off tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal stenosis pleen, injury of, healed pleenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of yndromes: Chronic Fatigue Syndrome (CFS) Cushing's	Small	
Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable Superficial, unsta		
Deep, other than head, face or neck Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable Superficial, unsta	arring alopecia	
Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable  cleritis cotoma leptum, nasal, deviation of sickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid  leep Apnea Syndrome coft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pipinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of symblepharon lyndromes:  Chronic Fatigue Syndrome (CFS) Cushing's		
Other Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable  cleritis cotoma leptum, nasal, deviation of sickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid  leep Apnea Syndrome coft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pipinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of symblepharon lyndromes:  Chronic Fatigue Syndrome (CFS) Cushing's	Deep, other than head, face or neck	
Retina Superficial, other than head, face, or neck Superficial, painful Superficial, unstable Superficial, uns		
Superficial, other than head, face, or neck Superficial, painful Superficial, unstable  cotoma leptum, nasal, deviation of lickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid Maxillary Pansinusitis Sphenoid Interest of tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin Ipinal stenosis Ipieen, injury of, healed Iplenectomy Ipondylolisthesis or segmental instability, spine Itomach, stenosis of Iymblepharon Iymdromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Superficial, painful Superficial, unstable Scleritis Scotoma Septum, nasal, deviation of Sicikle cell anemia Sinusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid Sleep Apnea Syndrome Soft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin Spinal fusion Spinal stenosis Spleen, injury of, healed Splenectomy Spotomy of, healed Splenectomy Spotomy of, stenosis of Syndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Superficial, unstable cliceritis ciccitoma cleptum, nasal, deviation of clickle cell anemia clinusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid clicep Apnea Syndrome contitistus acroma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pipeen, injury of, healed pipenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of cymblepharon cyndromes:  Chronic Fatigue Syndrome (CFS) Cushing's		
icleritis cotoma cotoma septum, nasal, deviation of sickle cell anemia inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid sphenoid sphenoid seleep Apnea Syndrome soft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin spinal fusion spinal stenosis spleen, injury of, healed splenenctomy spondylolisthesis or segmental instability, spine tiomach, stenosis of symblepharon syndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
icotoma leptum, nasal, deviation of lickle cell anemia linusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome loft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin lipinal fusion lipinal fusion lipinal stenosis pleen, injury of, healed liplenectomy lipondylolisthesis or segmental instability, spine tomach, stenosis of lymblepharon lyndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
eptum, nasal, deviation of ickle cell anemia insusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid Ileep Apnea Syndrome oft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes:  Chronic Fatigue Syndrome (CFS) Cushing's		
inusitis:  Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
inusitis:  Ethmoid		
Ethmoid Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Frontal Maxillary Pansinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma: Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Maxillary Pansinusitis Sphenoid leep Apnea Syndrome oft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Pansinusitis Sphenoid  Ileep Apnea Syndrome Ioft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pipelen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of Symblepharon Syndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Sphenoid leep Apnea Syndrome oft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
leep Apnea Syndrome oft tissue sarcoma:  Muscle, fat, or fibrous connected Neurogenic origin  Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
off tissue sarcoma:  Muscle, fat, or fibrous connected  Neurogenic origin  Vascular origin  pinal fusion  pinal stenosis  pleen, injury of, healed plenectomy  pondylolisthesis or segmental instability, spine  tomach, stenosis of  ymblepharon yndromes:  Chronic Fatigue Syndrome (CFS)  Cushing's		
Muscle, fat, or fibrous connected Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenenctomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Neurogenic origin Vascular origin pinal fusion pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Vascular origin pinal fusion ————————————————————————————————————		
pinal fusion		
pinal stenosis pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
pleen, injury of, healed plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of yndromes:  Chronic Fatigue Syndrome (CFS) Cushing's		
plenectomy pondylolisthesis or segmental instability, spine tomach, stenosis of ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
condylolisthesis or segmental instability, spine		
tomach, stenosis of		
ymblepharon yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
yndromes: Chronic Fatigue Syndrome (CFS) Cushing's		
Chronic Fatigue Syndrome (CFS)		
Cushing's		
	• , , ,	
Meniere's		
MOTIOTO 5	Meniere's	
Raynaud's	Raynaud's	
Sleep Apnea	Sleep Apnea	
synovitis		

	Diagnost code No
Syphilis:	
Cerebrospinal	8
Meningovascular	8
Syphilitic heart disease	3
Syringomyelia	3
Fabes dorsalis	8
Tenosynovitis	į
Testis:	,
Atrophy, complete	7
Removal	· ·
Fhrombocytopenia	7
Fhrombosis, brain	8
Thyroid gland:	
Nontoxic adenoma	7
Toxic adenoma	7
Fig. convulsive	3
Finnitus, recurrent	6
Fourtie sheet well defeat	7
Fraumatic chest wall defect	
Adenitis	7
Bones and joints	5
Eye	ě
Kidney	7
Luposa (lupus vulgaris)	7
Miliary	6
Pleurisy, active or inactive	6
Pulmonary:	
Active, far advanced	6
Active, moderately advanced	6
Active, minimal	(
Active, advancement unspecified	(
Active, chronic	(
Inactive, far advanced	(
Inactive, not advanced	
Inactive, minimal	è
Inactive, advancement unspecified	ě
Tuberculosis luposa (lupus vulgaris)	7
Tympanic membrane	6
Typhus, scrub	6
Jlcer:	
Duodenal	7
Gastric	7
Marginal	7
Jreter, stricture of	7
Jreterolithiasis	7
Urethra.	7
Fistula	
Urticaria	-
Jterus:	
And both ovaries, removal	7
Disease or injury	7
Displacement	7
Prolapse	7
Removal	7
Jveitis	6
/agina, disease or injury	
/agotomy	
/alvular heart disease	
/aricose veins	-
/asculitis, primary cutaneous	7
/ertebral fracture or dislocation	
Visceroptosis	-
Vision: see also Blindness and Loss of	
One eye 5/200 (1.5/60), with visual acuity of other eye:	
5/200 (1.5/60)	6
10/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	ě
	6
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	,

#### Pt. 6

	Diagnostic code No.
One eye 10/200 (3/60), with visual acuity of other eye:	
10/200 (3/60); 15/200 (4.5/60); 20/200 (6/60)	607
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	607
20/40 (6/12)	607
One eye 15/200 (4.5/60), with visual acuity of other eye:	
15/200 (4.5/60) or 20/200 (6/60)	607
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	607
20/40 (6/12)	607
One eye 20/200 (6/60), with visual acuity of other eye:	
20/200 (6/60)	607
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	607
20/40 (6/12)	607
One eye 20/100 (6/30), with visual acuity of other eye: and other eye:	
20/100 (6/30); 20/70 (6/21); 20/50 (6/15)	60
20/40 (6/12)	60
One eye 20/70 (6/21), with visual acuity of other eye:	
20/70 (6/21) or 20/50 (6/15)	60
20/40 (6/12)	60
One eye 20/50 (6/15), with visual acuity of other eye:	
20/50 (6/15)	60
20/40 (6/12)	60
Each eye 20/40 (6/12)	60
iligo	78:
ılva disease or injury of	76
eak foot	52

[72 FR 13003, Mar. 20, 2007]

#### PART 5 [RESERVED]

# PART 6—UNITED STATES GOVERNMENT LIFE INSURANCE

Age

Sec.

6.1 Misstatement of age.

PREMIUMS

6.2 Premium rate.

POLICIES

6.3 Incontestability of United States Government life insurance.

BENEFICIARY OF UNITED STATES GOVERNMENT LIFE INSURANCE

- 6.4 Proof of age, relationship and marriage.
- 6.5 Conditional designation of beneficiary.
- 6.6 Change of beneficiary.
- 6.7 Claims of creditors, taxation.

### OPTIONAL SETTLEMENT

- $6.8\,\,$  Selection, revocation and election.
- 6.9 Election of optional settlement by beneficiary.
- 6.10 Options.

#### DIVIDENDS

- 6.11 How dividends are paid.
- 6.12 Special dividends.

#### LOANS

6.13 Policy loans.

#### CASH VALUE

- 6.14 Cash value; other than special endowment at age 96 plan policy.
- 6.15 Cash value; special endowment at age 96 plan policy.
- 6.16 Payment of cash value in monthly installments.

#### INDEBTEDNESS

6.17 Collection of any indebtedness.

TOTAL PERMANENT DISABILITY BENEFITS

6.18 Other disabilities deemed to be total and permanent.

#### DEATH BENEFITS

- 6.19 Evidence to establish death of the in-
- DETERMINATION OF LIABILITY UNDER SECTIONS 302 AND 313, WORLD WAR VETERANS' ACT, 1924, SECTIONS 607 AND 602(V)(2), NATIONAL SERVICE LIFE INSURANCE ACT, 1940, AS AMENDED, AND SECTIONS 1921 AND 1957 OF TITLE 38, UNITED STATES CODE
- 6.20 Jurisdiction.

#### APPEALS

6.21 Guardian: definition and authority.

AUTHORITY: 38 U.S.C. 501, 1940-1963, 1981-1988, unless otherwise noted.