ing

100

§4.124

§4.124 Neuralgia, cranial or peripheral.

Neuralgia, cranial or peripheral, characterized usually by a dull and intermittent pain, of typical distribution so as to identify the nerve, is to be rated on the same scale, with a maximum equal to moderate incomplete paralysis. See nerve involved for diagnostic code number and rating. Tic douloureux, or trifacial neuralgia, may be rated up to complete paralysis of the affected nerve.

§4.124a Schedule of ratings—neurological conditions and convulsive disorders.

[With the exceptions noted, disability from the following diseases and their residuals may be rated from 10 percent to 100 percent in proportion to the impairment of motor, sensory, or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, etc., referring to the appropriate bodily system of the schedule. With partial loss of use of one or more extremities from neurological lesions, rate by comparison with the mild, moderate, severe, or complete paralysis of peripheral nerves]

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM

	Rat- ing
8000 Encephalitis, epidemic, chronic:	
As active febrile disease	100
Rate residuals, minimum	10
Brain, new growth of:	400
8002 Malignant	100
for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized,	
the rating will be made on neurological residu-	
als according to symptomatology.	
Minimum rating	30
8003 Benign, minimum	60
Rate residuals, minimum	10
8004 Paralysis agitans:	
Minimum rating	30
8005 Bulbar palsy	100
8007 Brain, vessels, embolism of. 8008 Brain, vessels, thrombosis of.	
8009 Brain, vessels, hemorrhage from:	
Rate the vascular conditions under Codes 8007	
through 8009, for 6 months	100
Rate residuals, thereafter, minimum	10
8010 Myelitis:	
Minimum rating	10
8011 Poliomyelitis, anterior:	
As active febrile disease	100
Rate residuals, minimum	10

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

8012 Hematomyelia:

For 6 months

Rate residuals, minimum	10
8013 Syphilis, cerebrospinal.	
8014 Syphilis, meningovascular.	
8015 Tabes dorsalis.	
NOTE: Rate upon the severity of convulsions, pa- ralysis, visual impairment or psychotic involve- ment, etc.	
8017 Amyotrophic lateral sclerosis	100
NOTE: Consider the need for special monthly	100
compensation.	
8018 Multiple sclerosis:	
Minimum rating	30
8019 Meningitis, cerebrospinal, epidemic:	
As active febrile disease	100
Rate residuals, minimum	10
8020 Brain, abscess of:	
As active disease	100
Rate residuals, minimum	10
Spinal cord, new growths of:. 8021 Malignant	100
8021 Malignant	100
for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized,	
the rating will be made on neurological residu-	
als according to symptomatology.	
Minimum rating	30
8022 Benign, minimum rating	60
Rate residuals, minimum	10
8023 Progressive muscular atrophy: Minimum rating	30
8024 Syringomyelia:	00
Minimum rating	30
8025 Myasthenia gravis:	00
Minimum rating	30
NOTE: It is required for the minimum ratings for	
residuals under diagnostic codes 8000-8025,	
that there be ascertainable residuals. Deter-	
minations as to the presence of residuals not	
capable of objective verification, i.e., head-	
aches, dizziness, fatigability, must be approached on the basis of the diagnosis re-	
proached on the basis of the diagnosis re-	
corded; subjective residuals will be accepted	
when consistent with the disease and not more likely attributable to other disease or no	
disease. It is of exceptional importance that	
when ratings in excess of the prescribed min-	
imum ratings are assigned, the diagnostic	
codes utilized as bases of evaluation be cited,	
in addition to the codes identifying the diag-	
noses.	
8045 Residuals of traumatic brain injury (TBI):	
There are three main areas of dysfunction	
that may result from TBI and have pro-	
found effects on functioning: cognitive	
(which is common in varying degrees	
after TBI), emotional/behavioral, and physical. Each of these areas of dysfunc-	
tion may require evaluation	
non may require evaluation	

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

ing

Subjective symptoms may be the only residual of TBI or may be associated with cog-nitive impairment or other areas of dysfunction. Evaluate subjective symptoms that are residuals of TBI, whether or not they are part of cognitive impairment, under the subjective symptoms facet in the table titled "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified." However, separately evaluate any residual with a distinct diagnosis that may be evaluated under another diagnostic code, such as migraine headache or Meniere's disease. even if that diagnosis is based on subjective symptoms, rather than under the "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified" table.

Evaluate emotional/behavioral dysfunction under §4.130 (Schedule of ratings—mental disorders) when there is a diagnosis of a mental disorder. When there is no diagnosis of a mental disorder, evaluate emotional/behavioral symptoms under the criteria in the table titled "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified."

Evaluate physical (including neurological) dysfunction based on the following list, under an appropriate diagnostic code: Motor and sensory dysfunction, including pain, of the extremities and face; visual impairment; hearing loss and tinnitus; loss of sense of smell and taste; seizures; gait, coordination, and balance problems; speech and other communication difficulties, including aphasia and related disorders, and dysarthria; neurogenic bladder; neurogenic bowel; cranial nerve dysfunctions; autonomic nerve dysfunctions; and endocrine dysfunctions.

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

The preceding list of types of physical dysfunction does not encompass all possible residuals of TBI. For residuals not listed here that are reported on an examination, evaluate under the most appropriate diagnostic code. Evaluate each condition separately, as long as the same signs and symptoms are not used to support more than one evaluation, and combine under § 4.25 the evaluations for each separately rated condition. The evaluation assigned based on the "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classified" table will be considered the evaluation for a single condition for purposes of combining with other disability evaluations.

Consider the need for special monthly compensation for such problems as loss of use of an extremity, certain sensory impairments, erectile dysfunction, the need for aid and attendance (including for protection from hazards or dangers incident to the daily environment due to cognitive impairment), being housebound, etc.

Evaluation of Cognitive Impairment and Subjective Symptoms

The table titled "Evaluation of Cognitive Impairment and Other Residuals of TBI Not Otherwise Classi fied" contains 10 important facets of TBI related to cognitive impairment and subjective symptoms. It provides criteria for levels of impairment for each facet, as appropriate, ranging from 0 to 3, and a 5th level, the highest level of impairment, labeled "total." However, not every facet has every level of severity. The Consciousness facet, for example, does not provide for an impairment level other than "total," since any level of impaired consciousness would be totally disabling. Assign a 100-percent evaluation if "total" is the level of evaluation for one or more facets. If no facet is evaluated as "total," assign the overall percentage evaluation based on the level of the highest facet as follows: 0 = 0 percent; 1 = 10 percent; 2 = 40 percent; and 3 = 70 percent. For example, assign a 70 percent evaluation if 3 is the highest level of evaluation for any facet.

Note (1): There may be an overlap of manifestations of conditions evaluated under the table titled "Evaluation Of Cognitive Impairment And Other Residuals Of TBI Not Otherwise Classified" with manifestations of a comorbid mental or neurologic or other physical disorder that can be separately evaluated under another diagnostic code. In such cases, do not assign more than one evaluation based on the same manifestations. If the manifestations of two or more conditions cannot be clearly separated, assign a single evaluation under whichever set of diagnostic criteria allows the better assessment of overall impaired functioning due to both conditions. However, if the manifestations are clearly separable, assign a separate evaluation for each condition

ORGANIC DISEASES OF THE CENTRAL NERVOUS

SYSTEM—Continued

§4.124a

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

	Rat- ing		Rat- ing
Note (2): Symptoms listed as examples at certain evaluation levels in the table are only examples and are not symptoms that must be present in order to assign a particular evaluation Note (3): "Instrumental activities of daily living" refers to activities other than self-care that are needed for independent living, such as meal preparation, doing housework and other chores, shopping, traveling, doing laundry, being responsible for one's own medications, and using a telephone. These activities are distinguished from "Activities of daily living," which refers to basic self-care and includes bathing or showering, dressing, eating, getting in or out of bed or a chair, and using the toilet. Note (4): The terms "mild," "moderate," and "severe" TBI, which may appear in medical records, refer to a classification of TBI made at, or close to, the time of injury rather than to the current level of func-		Purely neurological disabilities, such as hemiplegia, cranial nerve paralysis, etc., due to cerebral arteriosclerosis will be rated under the diagnostic codes dealing with such specific disabilities, with citation of a hyphenated diagnostic code (e.g., 8046–8207). Purely subjective complaints such as headache, dizziness, tinnitus, insomnia and irritability, recognized as symptomatic of a properly diagnosed cerebral arteriosclerosis, will be rated 10 percent and no more under diagnostic code 9305. This 10 percent rating will not be combined with any other rating for a disability due to cerebral or generalized arteriosclerosis. Ratings in excess of 10 percent for cerebral arteriosclerosis under diagnostic code 9305 are not assignable in the absence of a diagnosis of multi-infarct dementia with cerebral arteriosclerosis. NOTE: The ratings under code 8046 apply only when the diagnosis of cerebral arteriosclerosis is substantiated by the entire clinical picture and not solely on findings of retinal arterio-	
tioning. This classification does not affect		sclerosis.	

ea Note ar me TE tioning. This classification does not affect the rating assigned under diagnostic code

Note (5): A veteran whose residuals of TBI are rated under a version of §4.124a, di-agnostic code 8045, in effect before Octo-ber 23, 2008 may request review under diagnostic code 8045, irrespective of whether his or her disability has worsened whether his or her disability has worsened since the last review. VA will review that veteran's disability rating to determine whether the veteran may be entitled to a higher disability rating under diagnostic code 8045. A request for review pursuant to this note will be treated as a claim for an increased rating for purposes of deter-mining the effective date of an increased rating awarded as a result of such review; however, in no case will the award be effective before October 23, 2008. For the purposes of determining the effective date of an increased rating awarded as a result of such review, VA will apply 38 CFR 3.114, if applicable..

8046 Cerebral arteriosclerosis:

8045..

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Memory, attention, con- centration, executive functions.	1	No complaints of impairment of memory, attention, concentration, or executive functions. A complaint of mild loss of memory (such as having difficulty following a conversation, recalling recent conversations, remembering names of new acquaintances, or find ing words, or often misplacing items), attention, concentration, or executive functions, but without objective evidence on testing.
	2	Objective evidence on testing of mild impairment of memory, attention, concentration, or executive functions resulting in mild functional impairment.
	3	Objective evidence on testing of moderate in pairment of memory, attention, concentration, or executive functions resulting in moderate functional impair ment.

§4.124a

Department of Veterans Affairs

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

sequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. 3 Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Total Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate	npairment and other esiduals of TBI not	evel of n- air- ent	Criteria	Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Judgment	7	otal	testing of severe im- pairment of memory, attention, concentra- tion, or executive func-	Orientation	3	Social interaction is frequently inappropriate. Social interaction is inappropriate most or all of the time. Always oriented to per-
Mildly impaired judgment. For complex or unfamiliar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. 2 Moderately impaired judgment. For complex or unfamiliar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. 3 Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decisions. Total Visual spatial orientation O Total Total		0	vere functional impair- ment.	Chornation		son, time, place, and situation.
and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. 2 Moderately impaired judgment. For complex or unfamiliar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. 3 Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Total	gment	-	Mildly impaired judg- ment. For complex or unfamiliar decisions,		1	Occasionally disoriented to one of the four as- pects (person, time, place, situation) of ori- entation.
judgment. For complex or unfamiliar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision, although has little difficulty with simple decisions. Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Total Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Visual spatial orientation 1 Total		2	identify, understand, and weigh the alter- natives, understand the consequences of choices, and make a reasonable decision.		2	Occasionally disoriented to two of the four as- pects (person, time, place, situation) of ori- entation or often dis- oriented to one aspect of orientation.
understand the con- sequences of choices, and make a reason- able decision, although has little difficulty with simple decisions. Moderately severely im- paired judgment. For even routine and familiar decisions, occa- sionally unable to iden- tify, understand, and weigh the alternatives, understand the con- sequences of choices, and make a reason- able decision. Total Total Total Motor activity (with intact motor and sensory sys- tem). 1 1 2 2 3 Total			judgment. For complex or unfamiliar decisions, usually unable to iden- tify, understand, and		3	Often disoriented to two or more of the four aspects (person, time, place, situation) of orientation.
Moderately severely impaired judgment. For even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total			understand the con- sequences of choices, and make a reason- able decision, although has little difficulty with		Total	Consistently disoriented to two or more of the four aspects (person, time, place, situation) of orientation.
even routine and familiar decisions, occasionally unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate		3	Moderately severely impaired judgment. For	motor and sensory sys-	0	Motor activity normal.
Total Severely impaired judgment. For even routine and familiar decisions, usually unable to identify, understand, and weigh the alternatives, understand the consequences of choices, and make a reasonable decision. For example, unable to determine appropriate Total			iar decisions, occa- sionally unable to iden- tify, understand, and weigh the alternatives, understand the con- sequences of choices, and make a reason-		1	Motor activity normal most of the time, but mildly slowed at times due to apraxia (inabil- ity to perform pre- viously learned motor activities, despite nor- mal motor function).
tify, understand, and weigh the alternatives, understand the con- sequences of choices, and make a reason- able decision. For ex- ample, unable to de- termine appropriate tify, understand, and Total Total	1	otal	Severely impaired judg- ment. For even routine and familiar decisions,		2	Motor activity mildly de- creased or with mod- erate slowing due to apraxia.
sequences of choices, and make a reason- able decision. For ex- ample, unable to de- termine appropriate Total Total Total Total			tify, understand, and weigh the alternatives,		3	Motor activity moderately decreased due to apraxia.
ample, unable to de- Visual spatial orientation 0 termine appropriate 1			sequences of choices, and make a reason-		Total	Motor activity severely decreased due to apraxia.
weather conditions or judge when to avoid dangerous situations or activities.			ample, unable to de- termine appropriate clothing for current weather conditions or judge when to avoid dangerous situations	Visual spatial orientation		Normal. Mildly impaired. Occasionally gets lost in unfamiliar surroundings, has difficulty reading maps or following directions. Is able to use
Social interaction	ial interaction		Social interaction is routinely appropriate.			assistive devices such as GPS (global positioning system).

§4.124a

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria	Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Subjective symptoms	3 Total 0	Moderately impaired. Usually gets lost in unfamiliar surroundings, has difficulty reading maps, following directions, and judging distance. Has difficulty using assistive devices such as GPS (global positioning system). Moderately severely impaired. Gets lost even in familiar surroundings, unable to use assistive devices such as GPS (global positioning system). Severely impaired. May be unable to touch or name own body parts when asked by the examiner, identify the relative position in space of two different objects, or find the way from one room to another in a familiar environment. Subjective symptoms that do not interfere with work; instrumental activities of daily living; or work, family, or other close relationships. Examples are: mild or occasional headaches, mild anxiety. Three or more subjective symptoms that mildly interfere with work; instrumental activities of daily living; or work, family, or other close relationships. Examples of findings that might be seen at this level of impairment are: intermittent dizziness, daily mild to moderate headaches, tinnitus, frequent insomnia, hypersensitivity to sound, hypersensitivity to light.	Neurobehavioral effects	0	Three or more subjective symptoms that moderately interfere with work; instrumental activities of daily living; or work, family, or other close relationships. Examples of findings that might be seen at this level of impairment are: marked fatigability, blurred or double vision, headaches requiring rest periods during most days. One or more neurobehavioral effects that do not interfere with workplace interaction or social interaction. Examples of neurobehavioral effects are: Irritability, impulsivity, unpredictability, lack of motivation, verbal aggression, belligerence, apathy, lack of empathy, lack of empathy, lack of empathy, and impaired awareness of disability. Any of these effects may range from slight to severe, although verbal and physical aggression are likely to have a more serious impact on workplace interaction and social interaction than some of the other effects. One or more neurobehavioral effects that occasionally interfere with workplace interaction, or both but do not preclude them. One or more neurobehavioral effects that frequently interfere with workplace interaction, or both but do not preclude them. One or more
					them.

Department of Veterans Affairs

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
	3	One or more neurobehavioral effects that interfere with or preclude workplace interaction, or both on most days or that occasionally require supervision for safety of self or others.
Communication	0	Able to communicate by spoken and written language (expressive communication), and to comprehend spoken and written language.
	1	Comprehension or ex- pression, or both, of either spoken lan- guage or written lan- guage is only occa- sionally impaired. Can communicate complex ideas.
	2	Inability to communicate either by spoken lan- guage, written lan- guage, or both, more than occasionally but less than half of the time, or to com- prehend spoken lan-
	3	guage, written lan- guage, or both, more than occasionally but less than half of the time. Can generally communicate complex ideas. Inability to communicate
		either by spoken lan- guage, written lan- guage, or both, at least half of the time but not all of the time, or to comprehend spo- ken language, written language, or both, at least half of the time
	Total	but not all of the time. May rely on gestures or other alternative modes of communication. Able to communicate basic needs. Complete inability to communicate either by spoken language, written language, or both, or to comprehend spoken language, written language, or both. Unable to communicate basic needs.

EVALUATION OF COGNITIVE IMPAIRMENT AND OTHER RESIDUALS OF TBI NOT OTHERWISE CLASSIFIED—Continued

Facets of cognitive impairment and other residuals of TBI not otherwise classified	Level of im- pair- ment	Criteria
Consciousness	Total	Persistently altered state of consciousness, such as vegetative state, minimally re- sponsive state, coma.

MISCELLANEOUS DISEASES

	Rat- ing
8100 Migraine:	
With very frequent completely prostrating and prolonged attacks productive of severe economic inadaptability	50
With characteristic prostrating attacks occurring on an average once a month over last several months	30
With characteristic prostrating attacks averaging one in 2 months over last several months	10
With less frequent attacks	0
8103 Tic, convulsive:	00
Severe Moderate	30 10
Mild	0
Note: Depending upon frequency, severity, mus-	U
cle groups involved.	
8104 Paramyoclonus multiplex (convulsive state,	
myoclonic type):	
Rate as tic; convulsive; severe cases	60
8105 Chorea, Sydenham's:	
Pronounced, progressive grave types	100
Severe	80
Moderately severe	50
Moderate	30
Mild	10
NOTE: Consider rheumatic etiology and com-	
plications.	
8106 Chorea, Huntington's.	
Rate as Sydenham's chorea. This, though a fa-	
milial disease, has its onset in late adult life,	
and is considered a ratable disability.	
8107 Athetosis, acquired.	
Rate as chorea.	
8108 Narcolepsy.	
Rate as for epilepsy, petit mal.	
DISEASES OF THE CRANIAL NERVES	

DISEASES OF THE CRANIAL NERVES

	Rat- ing
Disability from lesions of peripheral portions of first, second, third, fourth, sixth, and eighth nerves will be rated under the Organs of Special Sense. The ratings for the cranial nerves are for unilateral involvement; when bilateral, combine but without the bilateral factor. Fifth (trigeminal) cranial nerve 8205 Paralysis of:	
Complete	50
Incomplete, severe	30 10
incomplete, moderate	10

§4.124a

DISEASES OF THE CRANIAL NERVES—Continued

DISEASES OF THE PERIPHERAL NERVES

		Oshadada (::	Rating	
	ing Schedule of ratings		Major	Minor
NOTE: Dependent upon relative degree of sensory manifestation or motor loss. 8305 Neuritis. 8405 Neuralgia. NOTE: Tic douloureux may be rated in accordance with severity, up to complete paralysis. Seventh (facial) cranial nerve		The term "incomplete paralysis," with this and other peripheral nerve injuries, indicates a degree of lost or impaired function substantially less than the type picture for complete paralysis given with each nerve, whether due to varied level of the nerve lesion or to partial regenera-		
3207 Paralysis of: Complete	30 20 10	of the flere lesion of to partial regenera- tion. When the involvement is wholly sensory, the rating should be for the mild, or at most, the moderate degree. The ratings for the peripheral nerves are for unilateral involvement; when bilateral, combine with application of the bilateral factor.		
8407 Neuralgia. Ninth (glossopharyngeal) cranial nerve.		Upper radicular group (fifth and sixth cervicals)		
8209 Paralysis of: Complete Incomplete, severe Incomplete, moderate NOTE: Dependent upon relative loss of ordinary sensation in mucous membrane of the phar-	30 20 10	8510 Paralysis of: Complete; all shoulder and elbow movements lost or severely affected, hand and wrist movements not affected	70	6
ynx, fauces, and tonsils. 8309 Neuritis.		Severe	50	4
8409 Neuralgia.		Moderate Mild	40 20	3 2
Tenth (pneumogastric, vagus) cranial nerve. 8210 Paralysis of: Complete	50 30 10	8610 Neuritis. 8710 Neuralgia. Middle radicular group 8511 Paralysis of: Complete; adduction, abduction and rotation of arm, flexion of elbow, and extension of wrist lost or severely af-		
3310 Neuritis. 3410 Neuralgia.		fectedIncomplete:	70	6
Eleventh (spinal accessory, external branch) cra-		Severe	50	4
nial nerve.		Moderate	40	3
8211 Paralysis of: Complete	30 20 10	Mild	20	2
8311 Neuritis. 8411 Neuralgia. Twelfth (hypoglossal) cranial nerve.		Complete; all intrinsic muscles of hand, and some or all of flexors of wrist and fingers, paralyzed (sub-		
3212 Paralysis of: Complete	50	stantial loss of use of hand)	70	6
Incomplete, severe	30	Severe Moderate	50 40	3
Incomplete, moderate	10	Mild	20	2
8412 Neuralgia.		All radicular groups		
		8513 Paralysis of: Complete Incomplete:	90	8
		Severe	70	6
		Moderate Mild	40 20	30

Department of Veterans Affairs

DISEASES OF THE PERIPHERAL NERVES-Continued

		Rating	
	Schedule of ratings		Minor
8613 8713	Neuritis. Neuralgia.		
0/13	Neuraigia.		
The	musculospiral nerve (radial nerve)		
	Paralysis of: omplete; drop of hand and fingers, wrist and fingers perpetually flexed, the thumb adducted falling within the line of the outer border of the index finger; can not extend hand at wrist, extend proximal phalanges of fingers, extend thumb, or make lateral movement of wrist; supination of hand, extension and flexion of elbow weakened, the loss of synergic motion of extensors impairs the hand grip seriously; total paralysis of the triceps occurs only as the greatest		
le.	rarity	70	60
III	Severe	50	40
	Moderate	30	20
	Mild	20	20
8614 8714	Neuritis.		
N	OTE: Lesions involving only "dissocia communis digitorum" and "paralysis I sor communis digitorum," will not e	oelow the	exten-

erate rating under code 8514.

· ·		
The median nerve		
8515 Paralysis of: Complete; the hand inclined to the ulnar side, the index and middle fingers more extended than normally, considerable atrophy of the muscles of the thenar eminence, the thumb in the plane of the hand (ape hand); pronation incomplete and defective, absence of flexion of index finger and feeble flexion of middle finger, cannot make a fist, index and middle fingers remain extended; cannot flex distal phalanx of thumb, defective opposition and abduction of the thumb, at right angles to palm; flexion of wrist weakened; pain with trophic disturbances Incomplete: Severe	70 50 30 10	60 40 20 10
The ulnar nerve		
8516 Paralysis of: Complete; the "griffin claw" deformity, due to flexor contraction of ring and little fingers, atrophy very marked in dorsal interspace and thenar and hypothenar eminences; loss of extension of ring and little fingers cannot spread the fingers (or reverse), cannot adduct the thumb; flexion of wrist weakened	60	50
Incomplete:	40	
Severe Moderate	40 30	30 20

DISEASES OF THE PERIPHERAL NERVES-Continued

Continued			
	Rat	Rating	
Schedule of ratings	Major	Minor	
Mild 8616 Neuritis. 8716 Neuralgia.	10	10	
Musculocutaneous nerve			
8517 Paralysis of:			
Complete; weakness but not loss of flexion of elbow and supination of forearm	30	20	
Incomplete:			
Severe	20	20	
Moderate Mild	10	10	
8617 Neuritis.	0	1	
8717 Neuralgia.			
Circumflex nerve			
8518 Paralysis of:			
Complete; abduction of arm is impos- sible, outward rotation is weakened; muscles supplied are deltoid and teres minor	50	40	
Incomplete: Severe	30	21	
Moderate	10	1	
Mild	0		
8618 Neuritis. 8718 Neuralgia.			
Long thoracic nerve			
8519 Paralysis of: Complete; inability to raise arm above shoulder level, winged scapula de-			
formityIncomplete:	30	2	
Severe	20	2	
Moderate	10	10	
Mild	0	(
NOTE: Not to be combined with lost mo der level.	tion above	shoul-	
8619 Neuritis. 8719 Neuralgia.			
NOTE: Combined nerve injuries should	be rated	by ref-	

NOTE: Combined nerve injuries should be rated by reference to the major involvement, or if sufficient in extent, consider radicular group ratings.

	Rating
Sciatic nerve	
8520 Paralysis of:	
Complete; the foot dangles and drops, no active movement possible of muscles below the knee, flexion of	80
knee weakened or (very rarely) lost Incomplete:	80
Severe, with marked muscular at-	
rophy	60
Moderately severe	40
Moderate	20
Mild	10

§4.124a

	Rating		Rating
8620 Neuritis.		Incomplete:	
8720 Neuralgia.		Severe	20
		Moderate	10
External popliteal nerve (common		Mild	10
peroneal)		8625 Neuritis.	
8521 Paralysis of:		8725 Neuralgia.	
Complete; foot drop and slight droop		Anterior crural nerve (femoral)	
of first phalanges of all toes, cannot		8526 Paralysis of:	
dorsiflex the foot, extension (dorsal		Complete; paralysis of quadriceps ex-	
flexion) of proximal phalanges of		tensor muscles	40
toes lost; abduction of foot lost,		Incomplete:	
adduction weakened; anesthesia		Severe	30
covers entire dorsum of foot and	40	Moderate	20
toes	40	Mild	10
Incomplete:	00	8626 Neuritis.	
Severe	30	8726 Neuralgia.	
Moderate	20	Internal saphenous nerve	
Mild	10	-	
8621 Neuritis.		8527 Paralysis of: Severe to complete	10
8721 Neuralgia.		Mild to moderate	0
Musculocutaneous nerve (superficial		8627 Neuritis.	U
peroneal)		8727 Neuralgia.	
9500 Paralysis of:		-	
8522 Paralysis of:	20	Obturator nerve	
Complete; eversion of foot weakened	30	8528 Paralysis of:	
Incomplete: Severe	00	Severe to complete	10
	20	Mild or moderate	0
Moderate Mild	10 0	8628 Neuritis.	
8622 Neuritis.	U	8728 Neuralgia.	
8722 Neuralgia.		External cutaneous nerve of thigh	
0722 Neuraigia.		8529 Paralysis of:	
Anterior tibial nerve (deep peroneal)		Severe to complete	10
8523 Paralysis of:		Mild or moderate	0
Complete; dorsal flexion of foot lost	30	8629 Neuritis.	
Incomplete:		8729 Neuralgia.	
Severe	20	llio-inguinal nerve	
Moderate	10	-	
Mild	0	8530 Paralysis of:	40
8623 Neuritis.		Severe to complete	10
8723 Neuralgia.		Mild or moderate	0
Internal popliteal nerve (tibial)		8630 Neuritis. 8730 Neuralgia.	
		8540 Soft-tissue sarcoma (of neurogenic	
8524 Paralysis of:		origin)	100
Complete; plantar flexion lost, frank adduction of foot impossible, flexion		Note: The 100 percent rating will be co	
and separation of toes abolished; no		for 6 months following the cessation	
muscle in sole can move; in lesions		gical, X-ray, antineoplastic chemothe	
of the nerve high in popliteal fossa,		other therapeutic procedure. At this	
plantar flexion of foot is lost	40	there has been no local recurrence or	
Incomplete:	10	tases, the rating will be made on resid	
Severe	30		
Moderate	20		
Mild	10	THE EPILEPSIES	
8624 Neuritis.			Rat-
8724 Neuralgia.			ing
Posterior tibial nerve		A thorough study of all material in §§ 4.121 an	
8525 Paralysis of:		4.122 of the preface and under the ratings for	
Complete; paralysis of all muscles of		epilepsy is necessary prior to any rating ac	-
sole of foot, frequently with painful		tion. 8910 Epilepsy, grand mal.	
paralysis of a causalgic nature; toes		Rate under the general rating formula for major	or
cannot be flexed; adduction is weak-		seizures.	
ened; plantar flexion is impaired	30	8911 Epilepsy, petit mal.	1

THE EPILEPSIES—Continued

Rate under the general rating formula for minor seizures.
NOTE (1): A major seizure is characterized by the generalized tonic-clonic convulsion with unconsciousness.
NOTE (2): A minor seizure consists of a brief interruption in consciousness or conscious control associated with staring or rhythmic blinking of the eyes or nodding of the head ("pure" petit mal), or sudden jerking movements of the arms, trunk, or head (myoclonic type) or sudden loss of postural control
(akinetic type).
General Rating Formula for Major and Minor Epileptic Seizures:
Averaging at least 1 major seizure per
month over the last year
Averaging at least 1 major seizure in 3 months over the last year; or more than 10 minor seizures weekly
Averaging at least 1 major seizure in 4 months over the last year; or 9-10 minor
seizures per week
At least 1 major seizure in the last 6 months
or 2 in the last year; or averaging at least 5 to 8 minor seizures weekly
At least 1 major seizure in the last 2 years;
or at least 2 minor seizures in the last 6
months
A confirmed diagnosis of epilepsy with a
history of seizures
NOTE (1): When continuous medication is shown
necessary for the control of epilepsy, the min-
imum evaluation will be 10 percent. This rating
will not be combined with any other rating for

NOTE (2): In the presence of major and minor seizures, rate the predominating type.

NOTE (3): There will be no distinction between

diurnal and nocturnal major seizures

8912 Epilepsy, Jacksonian and focal motor or sensory.

8913 Epilepsy, diencephalic.

epilepsy.

Rate as minor seizures, except in the presence of major and minor seizures, rate the predominating type.

8914 Epilepsy, psychomotor.

Major seizures:

Psychomotor seizures will be rated as major seizures under the general rating formula when characterized by automatic states and/or generalized convulsions with unconsciousness

Minor seizures:

Psychomotor seizures will be rated as minor seizures under the general rating formula when characterized by brief transient episodes of random motor movements, hallucinations, perceptual illusions, abnormalities of thinking, memory or mood, or autonomic disturbances

Mental Disorders in Epilepsies: A nonpsychotic organic Mental Disorders in Epilepsies: A nonpsychotic organic brain syndrome will be rated separately under the appropriate diagnostic code (e.g., 9304 or 9326). In the absence of a diagnostic ropersychotic organic psychiatric disturbance (psychotic, psychoneurotic or personality disorder) if diagnosed and shown to be secondary to or directly associated with epilepsy will be rated separately. The psychotic or psychroneurotic disorder will be rated under the appropriate diagnostic code. The personality disorder will be rated as a diágnostic code. The personality disorder will be rated as a dementia (e.g., diagnostic code 9304 or 9326). Epilepsy and Unemployability: (1) Rating specialists must bear in mind that the epileptic, although his or her seizures are controlled, may find employment and rehabilitation difficult of attainment due to employer reluctance to the hiring of the

(2) Where a case is encountered with a definite history of unemployment, full and complete development should be un-dertaken to ascertain whether the epilepsy is the determining factor in his or her inability to obtain employment.

(3) The assent of the claimant should first be obtained for

(a) The assent of the challful should his be obtained by permission to conduct this economic and social survey. The purpose of this survey is to secure all the relevant facts and data necessary to permit of a true judgment as to the reason for his or her unemployment and should include information

Rat-

100

80

60

40

20

10

as to:

(a) Education;
(b) Occupations prior and subsequent to service;
(c) Places of employment and reasons for termination;
(d) Wages received;
(e) Number of seizures.
(4) Upon completion of this survey and current examination, the case should have rating board consideration. Where in the judgment of the rating board the veteran's unemployability is due to epilepsy and jurisdiction is not vested in that body by reason of schedular evaluations, the case should be submitted to the Compensation Service or the Director, Pension and Fiduciary Service. and Fiduciary Service.

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 40 FR 42540, Sept. 15, 1975; 41 FR 11302, Mar. 18, 1976; 43 FR 45362, Oct. 2, 1978; 54 FR 4282, Jan. 30, 1989; 54 FR 49755, Dec. 1, 1989; 55 FR 154, Jan. 3, 1990; 56 FR 51653, Oct. 15, 1991; 57 FR 24364, June 9, 1992; 70 FR 75399, Dec. 20, 2005; 73 FR 54705, Sept. 23, 2008; 73 FR 69554, Nov. 19, 2008; 76 FR 78824, Dec. 20, 2011; 79 FR 2100, Jan. 13, 2014]

MENTAL DISORDERS

§ 4.125 Diagnosis of mental disorders.

(a) If the diagnosis of a mental disorder does not conform to DSM-5 or is not supported by the findings on the examination report, the rating agency shall return the report to the examiner to substantiate the diagnosis. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), American Psychiatric Association (2013), is incorporated by reference into this section with the approval of the Director of the Federal Register under 5 U.S.C. 552(a) and 1 CFR part 51. To enforce any edition other than that specified in this section, the Department of Veterans Affairs must publish notice of change in the FEDERAL REGISTER and the material must be available to the public. All approved material is available from the American Psychiatric Association, 1000 Wilson Boulevard, Suite 1825, Arlington, VA 22209-3901, 703-907-7300, http://www.dsm5.org. It is also available for inspection at the Office of Regulation Policy and Management, Department of Veterans Affairs, 810 Vermont Avenue NW., Room 1068,